Dentistry and Intensive Care Unit: A Brief Report

Lisiane Cristina Bannwart¹ Clóvis Lamartine de Moraes Melo Neto¹ Daniela Micheline dos Santos^{1,2}
 André Luiz de Melo Moreno¹ Aldiéris Alves Pesqueira¹ Marcelo Coelho Goiato^{1,2}
 André Pinheiro de Magalhães Bertoz³

¹ Department of Dental Materials and Prosthodontics, School of Dentistry, São Paulo State University, Aracatuba, São Paulo, Brazil

² Oral Oncology Center, School of Dentistry, São Paulo State University, Aracatuba, São Paulo, Brazil

³ Department of Pediatric and Social Dentistry, School of Dentistry, São Paulo State University, Aracatuba, São Paulo, Brazil

Address for correspondence Marcelo Coelho Goiato, DDS, MS, PhD, Department of Dental Materials and Prosthodontics, School of Dentistry, São Paulo State University, José Bonifácio, 1193; Aracatuba, São Paulo, ZIP Code: 16050-050, Brazil (e-mail: m.goiato@unesp.br).

Eur J Dent 2022;16:449–453.

Abstract

Objective The aim of this study is to verify whether removable dentures of patients admitted to an intensive care unit (ICU) are niches of microorganisms that can cause pathologies (*Staphylococcus aureus*, *Candida spp.*, and enterobacteria).

Materials and Methods Fifteen patients who were denture wearers (removable partial denture and complete denture) were included in this study. Patients must wear their dentures daily, and these dentures must have acrylic parts. Microbial biofilm was collected from the acrylic part of one denture of each patient. Then, the biofilm was seeded on different culture media: Sabouraud agar, blood agar, MacConkey agar, and mannitol salt agar. In this study, biochemical evaluations of microorganisms were performed.

Statistical analysis The percentage of dentures with the microorganism identified by each culture medium was calculated.

Results In total, 100% of the dentures were positive for *Staphylococcus spp.* (blood agar) and *Candida spp.* (Sabouraud agar); 33.3% of the dentures were positive for *S. aureus* (Mannitol salt agar); and 13.3% of the dentures were positive for *Shigella spp.* (MacConkey agar).

Conclusion Removable dentures of patients (removable partial dentures and complete dentures) admitted to an ICU are niches of microorganisms that can cause pathologies.

Keywords

- complete denture
- intensive care units
- removable partial denture
- dental prosthesis
- cross infection
- acrylic resin

Introduction

Intensive care units (ICUs) have contributed to the survival of patients with trauma and other life-threatening conditions.¹ Despite this, ICUs are associated with a high risk of nosocomial infections.^{1,2}

Nosocomial infections are infections that were not present in the patient at the time of admission to a health facility and appear 48 hours after the patient's hospital

admission.^{3,4} It is estimated that the incidence of these infections is two to five times higher in ICUs than in the general hospitalized population.² According to Ak et al, ICU patients are at increased risk of developing nosocomial infections for a variety of reasons, including presence of underlying diseases, prolonged ICU stay, invasive diagnostic and monitoring procedures performed, impaired host defenses, and colonization by resistant microorganisms.²

published online
December 1, 2021

DOI <https://doi.org/10.1055/s-0041-1735797>.
ISSN 1305-7456.

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

Nosocomial infections can affect any part of the body; however, the respiratory tract is the region most often affected (nosocomial pneumonia).^{2,5} Nosocomial pneumonia can occur when the patient aspirates microorganisms (respiratory pathogens), and these microorganisms reach the lower respiratory tract.^{6,7} Microbial plaque can be a source of respiratory pathogens, and it can be found on teeth and removable dentures.^{6,7}

Based on the fact that elderly patients admitted to an ICU may be wearers of removable dentures (removable partial denture and complete denture),⁸ and that aging is associated with an increased risk of hospital-acquired infections⁹; the aim of this study is to verify whether removable dentures of patients admitted to an ICU are niches of microorganisms that can cause pathologies (*S. aureus*, *Candida spp.*, and enterobacteria).

Materials and Methods

Ethics Committee

This study was approved by the ethics committee of the Paulista University (number: 45973115.9.0000.5512). This study included patients admitted to the ICU of the Hospital 28 de Agosto in the city of Manaus (Brazil). This study was conducted between September and December 2017.

Initially, 68 patients admitted to the ICU were selected. After applying the inclusion and exclusion criteria, 15 patients (11 men and 4 women) were included in this study.

Inclusion Criteria

- Men and women over 18 years of age.
- Each patient must wear at least one removable denture (complete denture or removable partial denture).
- Removable dentures must have acrylic parts.
- Patients must wear their removable dentures in the ICU.
- Patients must remain in the ICU for at least 6 days.
- Patients must wear their removable dentures daily.
- No use of antibiotics.

Exclusion Criteria

- Patients with bronchospasm, altered intracranial pressure, or hemodynamic instability.
- Patients in terminal stage.
- Patients who refused to participate in this study.

Sample Collection

The team of nurses performed the cleaning of the oral tissue and removable dentures of patients. The patients' oral tissue was cleaned with a sterile gauze moistened with 0.12% chlorhexidine. Removable dentures were cleaned with sterile water and sterile gauze. There were no dentists working at the hospital.

After cleaning one denture of each patient, the biofilm was collected on the acrylic part of the denture. A sterile cotton swab moistened with sterile saline was rubbed over the acrylic base of the denture for 1 minute. After biofilm collection, the swab was placed in a test tube (16 × 160 mm) with 3 mL of sterile buffered saline solution (phosphate buffered

saline), and this set was stored in a thermobox at a temperature between 2°C and 8°C. Then, the collected material was sent to the microbiology laboratory of the Paulista University.

The biofilm was seeded on different culture media: Sabouraud agar, blood agar, MacConkey agar, and Mannitol salt agar. Petri dishes were placed in a laboratory oven at 35°C, and after 48 hours, the presence or absence of colonies was checked. In this study, biochemical evaluations of microorganisms were performed.

The percentage of dentures with the microorganism identified by each culture medium was calculated.

Results

The mean age of the participants was approximately 71 years, and the mean time of wearing the same denture or pair of dentures was 10.5 years.

All partially edentulous patients had dental calculus and gingival inflammation. In addition, in 100% of patients the following characteristics were observed: (1) removable dentures in poor condition (bacterial calculus adhesion on dentures, wear of acrylic teeth, and loss of retention and stability of dentures); (2) patients slept with their dentures removable; (3) tongue coating; and (4) oral tissue with reddened areas, suggesting the presence of candidiasis.

► **Table 1** shows that 100% of the dentures were positive for *Staphylococcus spp.* (blood agar) and *Candida spp.* (Sabouraud agar). It was also observed that 33.3% of the dentures were positive for *S. aureus* (Mannitol salt agar); and 13.3% of dentures were positive for *Shigella spp.* (MacConkey agar).

Discussion

In the present study, teeth and mucosa were cleaned with 0.12% chlorhexidine. Chlorhexidine can destroy bacteria by breaking their membranes.¹⁰ Another method that can be used in conjunction with chemical cleaning is mechanical brushing of the teeth, mucosa and tongue. Thus, for ICU patients, it is important that brushing (mechanical cleaning) be performed before chemical cleaning (0.12% chlorhexidine) to remove the tongue coating, bacterial plaque and food remains.⁸ In the present study, tongue coating, dental calculus, gingival inflammation and oral tissue with reddened areas were observed in patients, therefore, the presence of a dentist in an ICU could help to prevent or control these situations. According to Miranda et al, biofilm on tooth surfaces, coated tongues, and periodontal disease tend to aggravate patient clinical conditions because they offer an optimal environment for growth of Gram-negative bacteria.⁸ This growth results in more virulent oral microflora.⁸

In this study, it was possible to identify pathogens on removable dentures (► **Table 1**). The poor hygiene of a removable denture allows the accumulation of biofilm on it, and this can contribute to the development of several oral and systemic infections.¹⁰ Thus, cleaning removable

Table 1 Results of different culture media

Sex	Denture type	Nutrition	Primary disease	Blood Agar	Sabouraud Agar	Mannitol Salt Agar	MacConkey Agar	Systemic complications
1	Male ^a	Parenteral	Cardiorespiratory arrest	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	Negative	
2	Male ^a	Oral	Cerebral vascular disease	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	Negative	Respiratory insufficiency
3	Male ^a	Oral	Acute congestive heart failure and respiratory failure	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	Negative	Pneumonia
4	Female	Oral	Diabetic ketoacidosis	Positive	<i>Candida spp.</i>	<i>Staphylococcus aureus</i>	Negative	Pneumonia/death
5	Male ^a	Parenteral	Acute myocardial infarction	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	Negative	
6	Male	Oral	Unstable angina	Positive	<i>Candida spp.</i>	<i>Staphylococcus aureus</i>	Negative	Pneumonia
7	Male	Oral	Cardiac tamponade	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	Negative	Respiratory insufficiency/death
8	Female ^a	Oral	Hypophosphatemia and muscle weakness	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	Positive	Diarrhea and death
9	Female	Parenteral	Severe acidosis	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	Negative	
10	Female	Oral	Acute myocardial infarction	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	negative	Diarrhea
11	Male	Oral	Acute myocardial infarction	Positive	<i>Candida spp.</i>	<i>Staphylococcus aureus</i>	negative	Pneumonia
12	Male	Oral	Diabetic ketoacidosis	Positive	<i>Candida spp.</i>	<i>Staphylococcus aureus</i>	negative	Pneumonia
13	Male	Oral	Acute myocardial infarction	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	Positive	Diarrhea
14	Male	Oral	Diabetic ketoacidosis	Positive	<i>Candida spp.</i>	<i>Staphylococcus spp.</i>	Negative	
15	Male	Oral	Cardiorespiratory arrest	Positive	<i>Candida spp.</i>	<i>Staphylococcus aureus</i>	Negative	Pneumonia
Total	66.6% Male			100% Positive	100% <i>Candida spp.</i>	66.7% <i>Staphylococcus spp.</i>	86.7% Negative	
	33.4% Female			0% Negative		33.3% <i>Staphylococcus aureus</i>	13.3% Positive	

Abbreviations: CD, complete denture; RPD, removable partial denture.

^aPresence of natural teeth in the lower dental arch.

dentures is important to avoid these problems.¹⁰ Literature reports that brushing removable dentures with soap, associated with chemical disinfection (e.g., 2% chlorhexidine) is effective in reducing the colonization of microorganisms (*Candida spp.* and *S. aureus*) on them.^{11–15}

Patients evaluated in this study slept with their dentures removable. Individuals who sleep with their dentures removable tend to have poor oral hygiene practices, high rates of microbial plaque on their tongues and dentures, oral candidiasis, and visit their dentists less often.¹⁶ Thus, these individuals may have an increased risk of aspirating microorganisms that can cause pneumonia.¹⁶ Therefore, this is particularly dangerous for patients admitted to an ICU because their immunity is usually reduced.² This may explain the pneumonia events observed in ►Table 1.

Removable dentures (removable partial dentures and complete dentures) should normally be replaced after 5 years of constant use.^{17,18} Over time, the roughness of removable dentures can increase due to their cleaning and daily use, favoring the adhesion of microorganisms.^{14,19} In this study, the mean time of wearing the same denture or pair of dentures was 10.5 years. This suggests that the surface roughness of the evaluated dentures was high.

Patients aged 65 and over are more prone to hospital infections due to generally low immunity, impaired physiological function, chronic illness, and medication use (e.g., immunosuppressants, anticholinergics, sedatives, and gastric acid inhibitors).⁹ Therefore, these patients have a high mortality rate.⁹ In the present study, the mean age of patients was approximately 71 years.⁹ Therefore, advanced age of the evaluated patients was a risk factor.

The most frequent etiologic agents that cause hospital infections are *S. aureus*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Acinetobacter spp.*, *Escherichia coli*, *Enterobacter spp.*, and *Candida spp.*^{20–22} In this study, two of these microorganisms (*S. aureus* and *Candida spp.*) were identified on removable dentures of patients included in this study (►Table 1).

Staphylococci aureus are responsible for skin and soft tissue infections, surgical site infections, endocarditis, pneumonia, and hospital-acquired bacteremia.^{23,24} In this study, 33.3% of the dentures were positive for *S. aureus* (Mannitol salt agar). Coincidentally, the wearers of these removable dentures had nosocomial pneumonia (►Table 1). Thus, it is possible that they have aspirated this microorganism causing this systemic complication.⁸

In the ICUs, *Candida spp.* may be associated with nosocomial infection,²⁵ morbidity, and mortality.²⁶ All dentures evaluated in this study were positive for *Candida spp.* It is worth noting that *Candida albicans* is the most common species found in the mouth.^{25,27} According to Akpan et al, the wearing of dentures produces a microenvironment with low oxygen and low pH, and this favors the growth of *Candida spp.*²⁷ In addition, the ease of adhesion of *Candida spp.* to acrylic, reduced flow of saliva under the surfaces of the dentures, and lack of oral hygiene can contribute to the development of candidiasis.²⁷ Thus, all of this information may explain the presence of *Candida spp.* in all samples evaluated (►Table 1).

According to Formal et al, invasive pathogens such as the *Shigella spp.* damage the intestinal mucosa.²⁸ This mucosal damage, which may range from a mild inflammatory response to gross ulceration, is a consequence of this pathogen's ability to invade the intestinal epithelium.²⁸ In this study, 13.3% of dentures were positive for *Shigella spp.*, and their wearers have had diarrhea (systemic complication). Therefore, it is possible that this pathogen caused this systemic complication.

Thus, based on the present study, the dentist is extremely important in an ICU.

Conclusion

Removable dentures of patients (removable partial dentures and complete dentures) admitted to an ICU are niches of microorganisms that can cause pathologies.

Funding
None.

Conflict of interest
None declared.

References

- 1 Maki DG, Crnich CJ, Safdar N. Nosocomial infection in the intensive care unit. *Crit Care Med* 2008;1003–1069
- 2 Ak O, Batirel A, Ozer S, Çolakoğlu S. Nosocomial infections and risk factors in the intensive care unit of a teaching and research hospital: a prospective cohort study. *Med Sci Monit* 2011;17(05):PH29–PH34
- 3 Wang L, Zhou KH, Chen W, Yu Y, Feng SF. Epidemiology and risk factors for nosocomial infection in the respiratory intensive care unit of a teaching hospital in China: a prospective surveillance during 2013 and 2015. *BMC Infect Dis* 2019;19(01):145
- 4 Divatia JV, Pulinkunnathil JG, Myatra SN. Nosocomial Infections and ventilator-associated pneumonia in cancer patients. *Oncologic Critical Care* 2019;1419–1439
- 5 Vincent JL. Nosocomial infections in adult intensive-care units. *Lancet* 2003;361(9374):2068–2077
- 6 Scannapieco FA. Pneumonia in nonambulatory patients. The role of oral bacteria and oral hygiene. *J Am Dent Assoc* 2006;137(Suppl):21S–25S
- 7 Yoneyama T, Yoshida M, Ohnishi T, et al; Oral Care Working Group. Oral care reduces pneumonia in older patients in nursing homes. *J Am Geriatr Soc* 2002;50(03):430–433
- 8 Miranda AF, de Paula RM, de Castro Piau CG, Costa PP, Bezerra ACB. Oral care practices for patients in intensive care units: a pilot survey. *Indian J Crit Care Med* 2016;20(05):267–273
- 9 Avci M, Ozgenc O, Coskuner SA, Olut AI. Hospital acquired infections (HAI) in the elderly: comparison with the younger patients. *Arch Gerontol Geriatr* 2012;54(01):247–250
- 10 de Andrade IM, Cruz PC, Silva-Lovato CH, de Souza RF, Souza-Gugelmin MC, Paranhos HdeF. Effect of chlorhexidine on denture biofilm accumulation. *J Prosthodont* 2012;21(01):2–6
- 11 Salles AE, Macedo LD, Fernandes RA, Silva-Lovato CH, Paranhos HdeF. Comparative analysis of biofilm levels in complete upper and lower dentures after brushing associated with specific denture paste and neutral soap. *Gerodontology* 2007;24(04):217–223
- 12 Moreno A, Dos Santos DM, Lamartine de Moraes Melo Neto C, Luiz de Melo Moreno A, de Magalhães Bertoz AP, Goiato MC. In vitro

- evaluation of the effect of different disinfectants on the biofilm of *Staphylococcus epidermidis* and *Staphylococcus aureus* formed on acrylic ocular prostheses. *PLoS One* 2020;15(10):e0240116
- 13 Bell JA, Brockmann SL, Feil P, Sackuvich DA. The effectiveness of two disinfectants on denture base acrylic resin with an organic load. *J Prosthet Dent* 1989;61(05):580–583
 - 14 Felipucci DN, Davi LR, Paranhos HF, Bezzon OL, Silva RF, Pagnano VO. Effect of different cleansers on the surface of removable partial denture. *Braz Dent J* 2011;22(05):392–397
 - 15 Mima EG, Pavarina AC, Vargas FdaS, Giampaolo ET, Machado AL, Vergani CE. Effectiveness of chlorhexidine on the disinfection of complete dentures colonised with fluconazole-resistant *Candida albicans*: in vitro study. *Mycoses* 2011;54(05):e506–e512
 - 16 Iinuma T, Arai Y, Abe Y, et al. Denture wearing during sleep doubles the risk of pneumonia in the very elderly. *J Dent Res* 2015; 94(3, Suppl):28S–36S
 - 17 Goiato MC, Dos Santos DM, Baptista GT, et al. Effect of thermal cycling and disinfection on colour stability of denture base acrylic resin. *Gerodontology* 2013;30(04):276–282
 - 18 Vermeulen AH, Keltjens HM, van't Hof MA, Kayser AF. Ten-year evaluation of removable partial dentures: survival rates based on retreatment, not wearing and replacement. *J Prosthet Dent* 1996; 76(03):267–272
 - 19 Dantas LC, da Silva-Neto JP, Dantas TS, Naves LZ, das Neves FD, da Mota AS. Bacterial adhesion and surface roughness for different clinical techniques for acrylic polymethyl methacrylate. *Int J Dent* 2016;2016:8685796
 - 20 Vandecandelaere I, Matthijs N, Van Nieuwerburgh F, et al. Assessment of microbial diversity in biofilms recovered from endotracheal tubes using culture dependent and independent approaches. *PLoS One* 2012;7(06):e38401
 - 21 Kusahara DM, Friedlander LT, Peterlini MA, Pedreira MLG. Oral care and oropharyngeal and tracheal colonization by Gram-negative pathogens in children. *Nurs Crit Care* 2012;17(03): 115–122
 - 22 Chacko R, Rajan A, Lionel P, Thilagavathi M, Yadav B, Premkumar J. Oral decontamination techniques and ventilator-associated pneumonia. *Br J Nurs* 2017;26(11):594–599
 - 23 Casey AL, Lambert PA, Elliott TS. *Staphylococci*. *Int J Antimicrob Agents* 2007;29(Suppl 3):S23–S32
 - 24 Sampedro GR, Bubeck Wardenburg J. *Staphylococcus aureus* in the intensive care unit: are these golden grapes ripe for a new approach? *J Infect Dis* 2017;215(Suppl 1):S64–S70
 - 25 Cliff PR, Sandoe JA, Heritage J, Barton RC. Use of multilocus sequence typing for the investigation of colonisation by *Candida albicans* in intensive care unit patients. *J Hosp Infect* 2008;69(01): 24–32
 - 26 Al-Dorzi HM, Sakkiha H, Khan R, et al. Invasive candidiasis in critically ill patients: a prospective cohort study in two tertiary care centers. *J Intensive Care Med* 2020;35(06):542–553
 - 27 Akpan A, Morgan R. Oral candidiasis. *Postgrad Med J* 2002;78 (922):455–459
 - 28 Formal SB, Hale TL, Sansonetti PJ. Invasive enteric pathogens. *Rev Infect Dis* 1983;5(Suppl 4):S702–S707