Evaluation of the Efficacy of Coenzyme Q₁₀ in the Management of Chronic Periodontitis: A Clinical Study

Gaurav Pandav¹ Sakshi Pandav² Sanjeev Jain¹ Divya Saxena¹ Ridhi Aggarwal¹ Prerna Gulati³

¹Department of Periodontology, Guru Nanak Dev Dental College and Research Institute, Sunam, Punjab, India

²Department of Orthodontics, Guru Nanak Dev Dental College and Research Institute, Sunam, Punjab, India

³Department of Public Health Dentistry, Guru Nanak Dev Dental College and Research Institute, Sunam, Punjab, India

Address for correspondence Divya Saxena, MDS, H. No. 29/5-B, Sodhian Street, Malgodham Road, Dhuri, Punjab 148024, India (e-mail: drdivyasaxena2254@gmail.com).

Dent J Adv Stud 2021;9:138–146.

Abstract

Aim The present study was aimed to clinically evaluate the effectiveness of coenzyme Q₁₀ (CoQ₁₀) in the management of chronic periodontitis.

Materials and Methods A total of 60 patients aged between 30 and 60 years with bleeding on probing and probing pocket depth (PPD) of 3 to 5 mm were selected and divided into three groups, with group I receiving scaling and root planing, group II CoQ₁₀ formulation for 6 weeks, and group III receiving both scaling and root planning, followed by coenzyme Q₁₀ administration for 6 weeks. PPD, relative attachment level (RAL), and gingival index were recorded in all the groups at baseline, 6 weeks, and 3 months, respectively. The data was statistically analyzed using Kruskal–Wallis, Mann–Whitney, and Wilcoxon signed rank tests.

Result Intragroup comparison showed statistically significant difference ($p \leq 0.05$) between the clinical parameters of all the groups at all time intervals, whereas intergroup comparison of all the parameters showed high statistically significant difference ($p \leq 0.001$) in group III at various time intervals followed by group I and group II.

Conclusion It was concluded from the study that CoQ₁₀ is a useful adjunct in treating chronic periodontitis by boosting the host resistance to periodontal disease.

Keywords

- antioxidants
- chronic periodontitis
- coenzyme Q₁₀

Introduction

Chronic periodontitis is an inflammatory disease resulting from the interaction of bacteria and the host inflammatory response.¹ The neutrophils play a vital role in host defense, causing destruction of periodontal pathogens by both oxygen-dependent and oxygen-independent mechanisms.² Excessive production of reactive oxygen species (ROS) by

neutrophils results in tissue damage due to oxidation of DNA, lipids, and proteins, stimulating proinflammatory cytokine release by monocytes and macrophages.³

Physiologically, an equilibrium exists between ROS and antioxidant defense mechanism. Elevated intracellular levels of ROS results in oxidative stress.²

Most of the periodontal therapies attempt to arrest the progression of periodontal destruction by causing an

published online
October 18, 2021

DOI <https://doi.org/10.1055/s-0041-1736267>
ISSN 2321-1482.

© 2021. Bhojia Dental College and Hospital affiliated to Himachal Pradesh University. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

alteration of the subgingival environment, which in most instances is sufficient to improve the periodontal health. Systemic pharmacological agents such as antibiotics, antioxidants, and chemical plaque control agents tend to reinforce the mechanical periodontal treatment and enhance the host defense mechanism.⁴ Provision of antioxidants, either through endogenous or exogenous sources, causes significant reduction in periodontal disease.⁵ Antioxidants such as coenzyme Q₁₀ (CoQ₁₀), vitamin C, vitamin E, carotenoids, and reduced glutathione protect cells and tissues from damage caused by free radicals.

CoQ₁₀ is a naturally occurring lipid-soluble antioxidant discovered by Frederick Crane and colleagues in 1957. Human cells synthesize CoQ₁₀ through a cascade of eight aromatic precursors, which require adequate levels of vitamins such as folic acid, niacin, riboflavin, vitamins B6, B2 and B12, pyridoxine, and pantothenic acid.⁴

CoQ₁₀ plays an important role in energy coupling by oxidative phosphorylation within mitochondria and also acts as a primary scavenger of free radicals. It also has an anti-inflammatory role because of its ability to repress inflammatory gene expression. CoQ₁₀ is an important lipid-soluble antioxidant which helps in protecting cellular membrane and lipoproteins from free radical-induced oxidative damage.^{4,6}

Deficient levels of CoQ₁₀ can result due to various reasons such as nutritional deficiencies, genetic or acquired defects, increased tissue needs, and declining levels of CoQ₁₀ in advancing age.⁷ Deficiency in CoQ₁₀ levels are associated with development and progression of periodontal disease. Hence, ROS-induced tissue destruction has led to the search of an appropriate complimentary antioxidant therapy, useful in the treatment of numerous diseases including chronic periodontitis.

Keeping these views in mind, the present study was aimed to clinically evaluate the effectiveness of CoQ₁₀ in the management of chronic periodontitis.

Materials and Methods

A total of 60 systemically healthy individuals aged between 30 and 60 years with bleeding on probing and probing pocket depth (PPD) of 3 to 5 mm were selected among the patients visiting the Department of Periodontology and Oral Implantology. Subjects who were on antibiotics for last 3 months; smokers and pregnant or lactating women were not included in the study.

Test Material

- Ubidecarenone (CoQ₁₀) capsules (Recharje Forte manufactured by Troikaa Pharmaceuticals Ltd. Uttarakhand, India) containing 30 mg of CoQ₁₀ was given in the dosage of one capsule two times daily for 6 weeks.

Method

Patients were selected by simple random sampling technique (using table of random numbers) with 20 patients in each group. Selected patients were divided into three groups with group I receiving scaling and root planing, group II CoQ₁₀ formulation for 6 weeks, and group III receiving both scaling and root planing, followed by CoQ₁₀ administration for 6 weeks.

Ethical clearance was obtained from the Institutional Review Board, and all the participants were informed about the study procedure by taking verbal and written informed consent.

Parameters like PPD, relative attachment level (RAL) and gingival index (GI)⁸ were recorded in all the groups at baseline, 6 weeks, and 3 months, respectively. At baseline, GI, PPD, and RAL were recorded among all the patients in three groups.

All patients were instructed to use only tooth brushing with toothpaste, and no use of mouth rinse was permitted.

Statistical Analysis

Statistical analysis was done using statistical package for social sciences (SPSS) software version 17. The quantitative data was presented as mean and standard deviation (SD). As the data was not normally distributed, nonparametric tests (Kruskal-Wallis, Mann-Whitney, and Wilcoxon signed rank) were applied to represent qualitative data. A *p*-value of ≤ 0.001 was considered to be statistically highly significant.

Results

The results are presented in the ►Tables (1–7). ►Table 1 and ►Fig. 1 show the baseline comparisons of various parameters in different groups. ►Tables 2–4 and ►Figs. 2–4 show the intragroup comparisons of all the parameters (PPD, RAL, and GI) at various time intervals in groups I, II, and III, respectively, using Wilcoxon signed rank test. ►Tables 5–7 and ►Figs. 5–7 shows the intergroup comparisons between group I and group II, group I and group III, and between group II and group III, respectively, for mean change in PPD, RAL, and GI at various time intervals.

Table 1 Comparison of various parameters in different groups at baseline

Parameters	Group I	Group II	Group III	<i>p</i> -Value	Significance
PPD	3.5 ± 0.36	3.5 ± 0.62	3.6 ± 0.68	0.853	NS
RAL	6.72 ± 0.92	6.77 ± 1.05	6.70 ± 0.96	0.779	NS
GI	1.81 ± 0.20	1.69 ± 0.21	1.74 ± 0.30	0.218	NS

Abbreviations: GI, gingival index; NS, nonsignificant; PPD, probing pocket depth; RAL, relative attachment level.

Table 2 Intragroup comparison of the parameters at various time intervals in group I (Wilcoxon signed rank test)

Parameters	Time interval	Mean $\pm$ SD	p-Value	Significance
PPD	Baseline–6 weeks	0.50 $\pm$ 0.16	< 0.001	HS
	Baseline–3 months	1.05 $\pm$ 0.27	< 0.001	HS
	6 weeks–3 months	0.55 $\pm$ 0.27	< 0.001	HS
RAL	Baseline–6 weeks	0.50 $\pm$ 0.28	< 0.001	HS
	Baseline–3 months	0.87 $\pm$ 0.48	< 0.001	HS
	6 weeks–3 months	0.37 $\pm$ 0.35	< 0.001	HS
GI	Baseline–6 weeks	0.61 $\pm$ 0.32	< 0.001	HS
	Baseline–3 months	0.64 $\pm$ 0.33	< 0.001	HS
	6 weeks–3 months	0.03 $\pm$ 0.10	0.157	NS

Abbreviations: GI, gingival index; HS, highly significant; NS, not significant; PPD, probing pocket depth; SD, standard deviation; RAL, relative attachment level.

Table 3 Intragroup comparison of the parameters at various time intervals in group II (Wilcoxon signed rank test)

Parameters	Time interval	Mean $\pm$ SD	p-Value	Significance
PPD	Baseline–6 weeks	0.20 $\pm$ 0.25	0.005	HS
	Baseline–3 months	0.40 $\pm$ 0.30	0.005	HS
	6 weeks–3 months	0.20 $\pm$ 0.25	< 0.001	HS
RAL	Baseline–6 weeks	0.12 $\pm$ 0.22	0.025	S
	Baseline–3 months	0.40 $\pm$ 0.38	0.001	HS
	6 weeks–3 months	0.27 $\pm$ 0.25	0.001	HS
GI	Baseline–6 weeks	0.18 $\pm$ 0.20	0.004	HS
	Baseline–3 months	0.38 $\pm$ 0.24	0.001	HS
	6 weeks–3 months	0.19 $\pm$ 0.16	< 0.001	HS

Abbreviations: GI, gingival index; HS, highly significant; PPD, probing pocket depth; RAL, relative attachment level; S, significant; SD, standard deviation.

Table 4 Intragroup comparison of the parameters at various time intervals in group III (Wilcoxon signed rank test)

Parameters	Time interval	Mean $\pm$ SD	p-Value	Significance
PPD	Baseline–6 weeks	0.65 $\pm$ 0.28	< 0.001	HS
	Baseline–3 months	1.35 $\pm$ 0.43	< 0.001	HS
	6 weeks–3 months	0.70 $\pm$ 0.34	< 0.001	HS
RAL	Baseline–6 weeks	0.52 $\pm$ 0.44	< 0.001	HS
	Baseline–3 months	1.22 $\pm$ 0.65	< 0.001	HS
	6 weeks–3 month	0.67 $\pm$ 0.37	< 0.001	HS
GI	Baseline–6 weeks	0.53 $\pm$ 0.36	< 0.001	HS
	Baseline–3 months	0.71 $\pm$ 0.31	0.014	S
	6 weeks–3 months	0.18 $\pm$ 0.31	< 0.001	HS

Abbreviations: GI, gingival index; HS, highly significant; PPD, probing pocket depth; RAL, relative attachment level; S, significant; SD, standard deviation.

Discussion

Chronic periodontitis is the result of abnormal host response and alterations in the complex microflora of subgingival plaque biofilm. This microflora causes tissue destruction

directly by the formation of toxic products and indirectly by activating the host defense mechanisms, that is, inflammation. Inflammation results in tissue infiltration by polymorphonuclear leukocytes and monocytes, subsequently followed by

Table 5 Intergroup comparison of mean change in PPD at various time intervals (Mann–Whitney test)

Time interval	Group	Mean $\pm$ SD	p-Value	Significance
Baseline–6 weeks	Group I	0.50 $\pm$ 0.16	< 0.001	HS
	Group II	0.20 $\pm$ 0.25		
	Group I	0.50 $\pm$ 0.16	0.049	S
	Group III	0.65 $\pm$ 0.28		
	Group II	0.20 $\pm$ 0.25	< 0.001	HS
	Group III	0.65 $\pm$ 0.28		
Baseline–3 months	Group I	1.05 $\pm$ 0.27	< 0.001	HS
	Group II	0.40 $\pm$ 0.30		
	Group I	1.05 $\pm$ 0.27	0.018	S
	Group III	1.35 $\pm$ 0.43		
	Group II	0.40 $\pm$ 0.30	< 0.001	HS
	Group III	1.35 $\pm$ 0.43		
6 weeks–3 months	Group I	0.55 $\pm$ 0.27	< 0.001	HS
	Group II	0.20 $\pm$ 0.25		
	Group I	0.55 $\pm$ 0.27	0.147	NS
	Group III	0.70 $\pm$ 0.34		
	Group II	0.20 $\pm$ 0.25	< 0.001	HS
	Group III	0.70 $\pm$ 0.34		

Abbreviations: GI, gingival index; HS, highly significant; NS, not significant; PPD, probing pocket depth; RAL, relative attachment level; S, significant; SD, standard deviation.

Table 6 Intergroup comparison of mean change in RAL at various time intervals (Mann–Whitney test)

Time interval	Group	Mean $\pm$ SD	p-Value	Significance
Baseline–6 weeks	Group I	0.50 $\pm$ 0.28	< 0.001	HS
	Group II	0.12 $\pm$ 0.22		
	Group I	0.50 $\pm$ 0.28	0.964	NS
	Group III	0.52 $\pm$ 0.44		
	Group II	0.12 $\pm$ 0.22	0.002	HS
	Group III	0.52 $\pm$ 0.44		
Baseline–3 months	Group I	0.87 $\pm$ 0.48	0.003	HS
	Group II	0.40 $\pm$ 0.38		
	Group I	0.87 $\pm$ 0.48	0.153	NS
	Group III	1.20 $\pm$ 0.65		
	Group II	0.40 $\pm$ 0.38	< 0.001	HS
	Group III	1.20 $\pm$ 0.65		
6 weeks–3 months	Group I	0.37 $\pm$ 0.35	0.422	NS
	Group II	0.27 $\pm$ 0.25		
	Group I	0.37 $\pm$ 0.35	0.016	S
	Group III	0.67 $\pm$ 0.37		
	Group II	0.27 $\pm$ 0.25	0.001	HS
	Group III	0.67 $\pm$ 0.37		

Abbreviations: GI, gingival index; HS, highly significant; NS, not significant; PPD, probing pocket depth; RAL, relative attachment level; S, significant; SD, standard deviation.

Table 7 Intergroup comparison of mean change in GI at various time intervals (Mann–Whitney test)

Time interval	Group	Mean $\pm$ SD	p-Value	Significance
Baseline–6 weeks	Group I	0.61 $\pm$ 0.32	< 0.001	HS
	Group II	0.18 $\pm$ 0.20		
	Group I	0.61 $\pm$ 0.32	0.532	NS
	Group III	0.53 $\pm$ 0.36		
	Group II	0.18 $\pm$ 0.20	0.003	HS
	Group III	0.53 $\pm$ 0.36		
Baseline–3 months	Group I	0.64 $\pm$ 0.33	0.010	HS
	Group II	0.38 $\pm$ 0.24		
	Group I	0.64 $\pm$ 0.33	0.489	NS
	Group III	0.71 $\pm$ 0.31		
	Group II	0.38 $\pm$ 0.24	0.001	HS
	Group III	0.71 $\pm$ 0.31		
6 weeks–3 months	Group I	0.03 $\pm$ 0.10	0.001	HS
	Group II	0.19 $\pm$ 0.16		
	Group I	0.03 $\pm$ 0.10	0.053	S
	Group III	0.18 $\pm$ 0.31		
	Group II	0.19 $\pm$ 0.16	0.194	NS
	Group III	0.18 $\pm$ 0.31		

Abbreviations: GI, gingival index; HS, highly significant; NS, not significant; PPD, probing pocket depth; RAL, relative attachment level; S, significant; SD, standard deviation.

phagocytosis, leading to nonmitochondrial O₂ consumption. This phenomenon, in turn, generates free radicals (FR) and ROS capable of damaging cellular membranes.

The oxidative damage is controlled by the antioxidant defense mechanisms of the surrounding tissues but plaque microflora responsible for periodontitis tend to alter this equilibrium. These findings, therefore, led to a search for an efficient “antioxidant therapy” in inflammatory periodontal disease.⁹

CoQ₁₀ was evaluated first in animal models of periodontitis and was shown to be capable of producing significant reduction in PPD and GI.¹⁰

Various clinical trials of Coenzyme Q₁₀ have been conducted so far and the results showed that it is an effective

intracellular antioxidant and its increased concentration in the diseased gingiva effectively suppresses advanced periodontal inflammation.^{4,11–14}

On the basis of new concepts of synergism with nutritional supplements and host response, the present study is intended to compare the following three treatment modalities: scaling and root planing, administration of CoQ₁₀ orally, and scaling and root planing combined with CoQ₁₀ in patients suffering from chronic periodontitis.

In our study, CoQ₁₀ was given in the dosage of 30 mg twice daily for 6 weeks, which was in accordance with the study conducted by Brzozowska et al.¹⁵ Different authors have used different dosages of the drug for different time periods.^{12,13,16,17} In our study, the time period for drug

Fig. 1 Baseline comparison of the different groups for various parameters. **Fig. 2** Intragroup comparison of the parameters at various time intervals in group I (Wilcoxon signed rank test).

Fig. 3 Intragroup comparison of the parameters at various time intervals in group II (Wilcoxon signed rank test).

Fig. 6 Intergroup comparison of mean change in relative attachment level (RAL) at various time intervals (Mann-Whitney test).

Fig. 4 Intragroup comparison of the parameters at various time intervals in group III (Wilcoxon signed rank test).

Fig. 7 Intergroup comparison of mean change in gingival index (GI) at various time intervals (Mann-Whitney test).

Fig. 5 Intergroup comparison of mean change in probing pocket depth (PPD) at various time intervals (Mann-Whitney test).

administration was 6 weeks, owing to the fact that a complete healing of the soft tissues takes at least 6 weeks, whereas maturation and remodeling can continue up to 6 months.¹⁸

The intragroup comparison of all the parameters at various time intervals showed significant improvement in all the three groups. Improvement in group I can be attributed to the fact that scaling and root planing causes gradual reduction in inflammatory cells and crevicular fluid flow, causing

repair of connective tissue, which results in decreased inflammatory response in the periodontal tissues.¹⁹ A similar improvement in the parameters was seen in group II, which can be attributed to the fact that CoQ₁₀ decreases the bacterial peptidase activity specific for periodontopathic bacteria like *Porphyromonas gingivalis*, *Bacteroides forsythus* and *Treponema denticola*, as proposed in the study conducted by Ishihara et al.²⁰ Wilson et al.²¹ and McRee et al.¹³ in their study showed that topical application of a redox agent to the periodontal pocket shifted subgingival microbiota from Gram-negative anaerobes, spirochetes and motile bacteria to facultative anaerobes and cocci, causing improvement in periodontal health. Similarly, significant improvement was also seen in our study in group III due to the combined effect of scaling and root planing along with CoQ₁₀ administration. The results of our study are also in accordance with the study conducted by Hanioka et al.²²

Intergroup comparison of the mean change in PPD was significantly higher in group III, followed by group I and group II. Hence, it was seen that scaling and root planing along with CoQ₁₀ in group III caused greatest reduction in PPD. These findings were in conjunction with study conducted by Sale et al.²³ and Saini.²⁴ Scaling and root planing reduces the number of subgingival plaque microorganisms significantly and produces a shift in its composition, with increase of Gram-negative anaerobes to one dominated by

Gram-positive facultative bacteria.^{25–31} It also leads to profound reduction in the number of spirochetes, motile rods and putative pathogens such as *Actinobacillus actinomyces*, *Porphyromonas gingivalis* and *Prevotella intermedia* and an increase in coccoid cells.^{26,28,30,32–34} These changes are accompanied by either reduction or elimination of the signs of clinical inflammation.^{35–42} CoQ₁₀, used as an adjunct to scaling and root planing, tends to lower the gingival lipid peroxidation at an advanced rate than scaling and root planing alone, thus causing greater improvement in periodontal health, as shown in the study conducted by Kuru et al,¹⁰ which can be the possible reason for improvement seen in PPD in group II and group III in our study. Also, the oral administration of CoQ₁₀ increases the succinate dehydrogenase-coenzyme Q₁₀ reductase enzyme activity in gingiva, which contributes to the extraordinary effective healing and could be the reason for the results seen in groups II and III.¹¹

The intergroup comparison of mean change in RAL showed significantly higher improvement in group III followed by group I and group II. Studies conducted by Hans et al, Sale et al, and Saini showed similar results.^{23,24,43} Possible reason for such an improvement, according to Matsumura et al,⁴⁴ is that the administration of CoQ₁₀ restores the elevated blood-citrate and serum-calcium levels, leading to significant improvement in alveolar bone porosity, myelofibrosis, and atrophy of periodontal membrane. It also profoundly effects citric acid metabolism and consequently prevents the decreased activity of osteoblasts and fibroblasts. According to Schmelzer et al,^{45,46} CoQ₁₀ is an efficient anti-inflammatory molecule that decreases the lipopolysaccharide (LPS)-induced secretion of proinflammatory cytokine tumor necrosis factor- α (TNF- α). CoQ₁₀ regulates the functioning of interleukin-5 (IL-5), thrombin, vitronectin, vitronectin receptor, and C-reactive proteins (CRP) via transcription factor NFkappaB1. The study concluded that CoQ₁₀ exerts an anti-inflammatory property via NFkappaB1-dependent gene expression. Fischer et al⁴⁷ concluded in their study that CoQ₁₀ played a significant role in the reduction of the expression of proinflammatory genes CXCL2, PMAIP1, and MMD, thus limiting the periodontal disease process. These can be the other possible reasons for the improvement seen in the parameters in our study.

On comparing the three groups for the GI at baseline, 6 weeks, and 3 months, group I and III showed similar results and group II showed least reduction in the GI. At 6 weeks to 3 months, highest mean change in GI was seen in group II. Studies conducted by Iwamoto et al,⁴⁸ Denny et al,⁴⁹ Brzozowska et al,¹⁵ Babbush et al,⁵⁰ Chatterjee et al,⁵¹ Hans et al,⁴³ and CoQ₁₀⁵² were in accordance with the results shown in our study. Elevated levels of CoQ₁₀ are significantly responsible for improvement in GI scores. The oral administration of CoQ₁₀ enhanced the restoration of the metabolic energy required for the diseased tissue and also led to increased activity of CoQ₁₀-dependent enzymes and oxygen utilization in inflamed human gingiva.^{12,53} CoQ₁₀, used as an adjunct to scaling and root planing, tends to lower the gingival lipid peroxidation, causing greater improvement,

which can be the possible reason for improvement seen in GI score in groups II and III in our study.⁸ Hansen et al⁵⁴ and Shizukuishi et al⁵⁵ concluded in their studies that CoQ₁₀ deficiencies in gingiva and leucocytes could predispose the tissues to periodontitis which, in turn, further augment the deficiency, as inflamed gingiva needs increased bioenergetics for proliferative and reparative changes. The present study showed improvement in clinical parameters like PPD, RAL, and GI score, which can be attributed to the fact that once the deficiency of CoQ₁₀ is met with its oral administration, significant improvement in periodontal health results.

In the light of our study and available literature, it can be concluded that use of CoQ₁₀ as an adjunct to scaling and root planing gives promising results in the treatment of plaque-induced gingivitis, significantly reduces PPD, and also causes improvement in clinical attachment level.

Conclusion

It was concluded from the study that CoQ₁₀ is a significant factor in boosting host resistance to periodontal disease, not only improving the periodontal status but also enhancing the quality of life. Further studies are required with other antioxidant agents for their use in clinical practice and opening new treatment options by strengthening the host side of the disease interaction. Additional studies are needed to define the precise role of CoQ₁₀, keeping in mind its appropriate dose, effectiveness, and bioavailability in the treatment of periodontal diseases. Further role of CoQ₁₀ also needed to be substantiated not only as an adjunct but also as a major therapeutic agent in the treatment of various diseases.

Conflict of Interest

None declared.

References

- Bhardwaj A, Bhardwaj SV. Role of antioxidants in periodontal disease and its therapy. *Asian J Med Res* 2012;1(02):62–67
- Sharma A, Sharma S. Reactive oxygen species and antioxidants in periodontics: a review. *Int J Dent Clin* 2011;3(02):44–47
- Ritchie CS, Kinane DF. Nutrition, inflammation, and periodontal disease. *Nutrition* 2003;19(05):475–476
- Prakash S, Sunitha J, Hans M. Role of coenzyme Q(10) as an antioxidant and bioenergizer in periodontal diseases. *Indian J Pharmacol* 2010;42(06):334–337
- Chandra RV, Srinivas G, Reddy AA, et al. Locally delivered antioxidant gel as an adjunct to nonsurgical therapy improves measures of oxidative stress and periodontal disease. *J Periodontal Implant Sci* 2013;43(03):121–129
- Pepping J. Coenzyme Q10. *Am J Health Syst Pharm* 1999;56(06):519–521
- Gupta HL, Soni S, Chaturvedi V, Dev V. Periodontal diseases and coenzyme Q₁₀: a review. *Indian J Dent Sci* 2012;2(04):124–127
- Loe H, Silness J. Periodontal disease in pregnancy II. Correlation between oral hygiene and periodontal conditions. *Acta Odontol Scand* 1964;22:112–135
- Battino M, Bullon P, Wilson M, Newman H. Oxidative injury and inflammatory periodontal diseases: the challenge of anti-oxidants to free radicals and reactive oxygen species. *Crit Rev Oral Biol Med* 1999;10(04):458–476

- 10 Kuru B, Yildiz D, Kayali R, Akcay T. Gingival lipid peroxidation and glutathione redox cycle before and after periodontal treatment with and without adjunctive Coenzyme Q₁₀. *Turk Klin J Dent Sci* 2006;12:1–8
- 11 Wilkinson EG, Arnold RM, Folkers K, Hansen I, Kishi H. Bioenergetics in clinical medicine. II. Adjunctive treatment with coenzyme Q in periodontal therapy. *Res Commun Chem Pathol Pharmacol* 1975;12(01):111–123
- 12 Shizukuishi S, Hanioka T, Tsunemitsu A, Fukunaga Y, Kishi T, Sato N. Clinical effect of Coenzyme Q₁₀ on periodontal disease; evaluation of oxygen utilisation in gingiva by tissue reflectance spectrophotometry. In: *Biomedical and Clinical Aspects of Coenzyme Q*. Amsterdam: Elsevier; 1986:359–368
- 13 McRee JT, Hanioka T, Shizukuishi S, Folkers K. Therapy with coenzyme Q₁₀ for patients with periodontal disease. Effect of coenzyme Q₁₀ on subgingival micro-organisms. *J Dent Health* 1993;43:659–666
- 14 Manthena S, Rao MV, Penubolu LP, Putcha M, Harsha AV. Effectiveness of CoQ₁₀ oral supplements as an adjunct to scaling and root planing in improving periodontal health. *J Clin Diagn Res* 2015;9(08):ZC26–ZC28
- 15 Brzozowska TM, Flisykowska AK, Swiatkowska MW, Stopa J. Healing of periodontal tissue assisted by coenzyme Q₁₀ with vitamin E: clinical and laboratory evaluation. *Pharmacol Rep* 2007;59(01):257–260
- 16 Iwamoto Y, Watanabe T, Okamoto H, Ohata N, Folkers K. Clinical effect of coenzyme Q₁₀ on periodontal disease. In: Folkers K, Yamamura Y, eds. *Biomedical and Clinical Aspects of Coenzyme Q*, Vol. 3. New York: Elsevier Science Publishers; 1981
- 17 Wilkinson EG, Arnold RM, Folkers K. Bioenergetics in clinical medicine. VI. adjunctive treatment of periodontal disease with coenzyme Q₁₀. *Res Commun Chem Pathol Pharmacol* 1976;14(04):715–719
- 18 Carranza FA, Newman MG, Takei H. *Carranza's Clinical Periodontology*. 10th edition. St. Louis, MO: Elsevier Saunders; 2006: 950–967
- 19 Cobb CM. Non-surgical pocket therapy: mechanical. *Ann Periodontol* 1996;1(01):443–490
- 20 Ishihara K, Naito Y, Kato T, et al. A sensitive enzymatic method (SK-013) for detection and quantification of specific periodontopathogens. *J Periodontal Res* 1992;27(02):81–85
- 21 Wilson M, Gibson M, Strahan D, Harvey W. A preliminary evaluation of the use of a redox agent in the treatment of chronic periodontitis. *J Periodontal Res* 1992;27(05):522–527
- 22 Hanioka T, Tanaka M, Ojima M, Shizukuishi S, Folkers K. Effect of topical application of coenzyme Q₁₀ on adult periodontitis. *Mol Aspects Med* 1994;15(Suppl):s241–s248
- 23 Sale ST, Parvez H, Yeltiwar RK, Vivekanandan G, Pundir AJ, Jain P. A comparative evaluation of topical and intrasulcular application of coenzyme Q₁₀ (Perio Q™) gel in chronic periodontitis patients: A clinical study. *J Indian Soc Periodontol* 2014;18(04):461–465
- 24 Saini R. A clinical and microbiological study to evaluate the effect of dietary supplement of coenzyme Q₁₀ in nonsurgical outcome of chronic periodontitis patients after phase I periodontal therapy. *Eur J Gen Dent* 2014;3(03):194–198
- 25 Listgarten MA, Lindhe J, Hellden L. Effect of tetracycline and/or scaling on human periodontal disease. Clinical, microbiological, and histological observations. *J Clin Periodontol* 1978;5(04): 246–271
- 26 Magnusson I, Lindhe J, Yoneyama T, Liljenberg B. Recolonization of a subgingival microbiota following scaling in deep pockets. *J Clin Periodontol* 1984;11(03):193–207
- 27 Mousquès T, Listgarten MA, Phillips RW. Effect of scaling and root planing on the composition of the human subgingival microbial flora. *J Periodontal Res* 1980;15(02):144–151
- 28 Renvert S, Wikström M, Dahlén G, Slots J, Egelberg J. Effect of root debridement on the elimination of *Actinobacillus actinomyce-*temcomitans and *Bacteroides gingivalis* from periodontal pockets. *J Clin Periodontol* 1990;17(06):345–350
- 29 Rosenberg ES, Evian CI, Listgarten MA. The composition of the subgingival microbiota after periodontal therapy. *J Periodontol* 1981;52(08):435–441
- 30 Shiloah J, Patters MR. Repopulation of periodontal pockets by microbial pathogens in the absence of supportive therapy. *J Periodontol* 1996;67(02):130–139
- 31 Slots J, Mashimo P, Levine MJ, Genco RJ. Periodontal therapy in humans. I. Microbiological and clinical effects of a single course of periodontal scaling and root planing, and of adjunctive tetracycline therapy. *J Periodontol* 1979;50(10):495–509
- 32 Sbordone L, Ramaglia L, Gulletta E, Iacono V. Recolonization of the subgingival microflora after scaling and root planing in human periodontitis. *J Periodontol* 1990;61(09):579–584
- 33 van Winkelhoff AJ, van der Velden U, de Graaff J. Microbial succession in recolonizing deep periodontal pockets after a single course of supra- and subgingival debridement. *J Clin Periodontol* 1988;15(02):116–122
- 34 Ximénez-Fyvie LA, Haffajee AD, Socransky SS. Comparison of the microbiota of supra- and subgingival plaque in health and periodontitis. *J Clin Periodontol* 2000;27(09):648–657
- 35 Badersten A, Nilvéus R, Egelberg J. Scores of plaque, bleeding, suppuration and probing depth to predict probing attachment loss. 5 years of observation following nonsurgical periodontal therapy. *J Clin Periodontol* 1990;17(02):102–107
- 36 Garrett JS. Effects of nonsurgical periodontal therapy on periodontitis in humans. A review. *J Clin Periodontol* 1983;10(05): 515–523
- 37 Greenstein G. Nonsurgical periodontal therapy in 2000: a literature review. *J Am Dent Assoc* 2000;131(11):1580–1592
- 38 Hughes TP, Caffesse RG. Gingival changes following scaling, root planning and oral hygiene. A biometric evaluation. *J Periodontol* 1978;49(05):245–252
- 39 Kaldahl WB, Kalkwarf KL, Patil KD, Molvar MP, Dyer JK. Long-term evaluation of periodontal therapy: I. Response to 4 therapeutic modalities. *J Periodontol* 1996;67(02):93–102
- 40 Lowenguth RA, Greenstein G. Clinical and microbiological response to nonsurgical mechanical periodontal therapy. *Periodontol* 2000 1995;9:14–22
- 41 Pihlstrom BL, McHugh RB, Oliphant TH, Ortiz-Campos C. Comparison of surgical and nonsurgical treatment of periodontal disease. A review of current studies and additional results after 61/2 years. *J Clin Periodontol* 1983;10(05):524–541
- 42 Quirynen M, Mongardini C, de Soete M, et al. The rôle of chlorhexidine in the one-stage full-mouth disinfection treatment of patients with advanced adult periodontitis. Long-term clinical and microbiological observations. *J Clin Periodontol* 2000;27(08): 578–589
- 43 Hans M, Prakash S, Gupta S. Clinical evaluation of topical application of perio-Q gel (Coenzyme Q₁₀) in chronic periodontitis patients. *J Indian Soc Periodontol* 2012;16(02):193–199
- 44 Matsumura T, Kimata M, Tsunemitsu A, Akiyoshi M. Effect of coenzyme Q on blood-citrate and calcium levels and on the periodontium of citrated rats. *J Dent Res* 1969;48(06):1264–1267
- 45 Schmelzer C, Lindner I, Rimbach G, Niklowitz P, Menke T, Döring F. Functions of coenzyme Q₁₀ in inflammation and gene expression. *Biofactors* 2008;32(1-4):179–183
- 46 Schmelzer C, Lorenz G, Lindner I, et al. Effects of coenzyme Q₁₀ on TNF-alpha secretion in human and murine monocytic cell lines. *Biofactors* 2007;31(01):35–41
- 47 Fischer A, Onur S, Schmelzer C, Döring F. Ubiquinol decreases monocytic expression and DNA methylation of the pro-inflammatory chemokine ligand 2 gene in humans. *BMC Res Notes* 2012;5:540
- 48 Iwamoto Y, Nakamura R, Folkers K, Morrison RF. Study of periodontal disease and coenzyme Q. *Res Commun Chem Pathol Pharmacol* 1975;11(02):265–271

- 49 Denny N, Chapple ILC, Matthews JB. Antioxidant and anti-inflammatory effects of coenzyme Q₁₀—a preliminary study. *J Dent Res* 1999;78:543
- 50 Babbush CA, Kagan D, Madhavi D, Rubido A. The efficacy of perfect smile toothpaste containing Coenzyme Q₁₀– β -Cyclodextrin inclusion complex in reducing mild to moderate gingivitis. *Natural Med J* 2010;2(05):16–20
- 51 Chatterjee A, Kandwal A, Singh N, Singh A. Evaluation of Co-Q10 anti-gingivitis effect on plaque induced gingivitis: A randomized controlled clinical trial. *J Indian Soc Periodontol* 2012;16(04): 539–542
- 52 Pitale U, Khetarpal S, Peter K, Pal V, Verma E, Gupta P. Evaluation of efficacy of coenzyme Q₁₀ in management of gingivitis and slight periodontitis- A clinical study. *Int J Curr Pharm Res* 2012;4(04):33–38
- 53 Shizukuishi S, Inoshita E, Tsunemitsu A, Nagae K, Kishi T, Folkers K. Effect of Coenzyme Q₁₀ on experimental periodontitis in dogs. *Biomed Res* 1983;4:33–40
- 54 Hansen IL, Iwamoto Y, Kishi T, Folkers K, Thompson LE. Bioenergetics in clinical medicine. IX. Gingival and leucocytic deficiencies of coenzyme Q10 in patients with periodontal disease. *Res Commun Chem Pathol Pharmacol* 1976;14(04):729–738
- 55 Shizukuishi S, Inoshita E, Tsunemitsu A, Takahashi K, Kishi T, Folkers K. Therapy by Coenzyme Q10 of experimental periodontitis in a dog-model supports results of human periodontitis therapy. In: *Biomedical and Clinical Aspects of Coenzyme Q*. Amsterdam: Elsevier; 1984:153–162