Survey-Based Research of Knowledge and Perception of Endodontic Treatment on an Indian Subpopulation

Shalin Ann Saji¹ Chitharanjan Shetty¹ Aditya Shetty¹ Mithra N. Hegde¹

¹Department of Conservative Dentistry and Endodontics, AB Shetty Memorial Institute of Dental Sciences, Mangalore, Karnataka, India

J Health Allied Sci^{NU} 2022;12:162–167.

Address for correspondence Shalin Ann Saji, Department of Conservative Dentistry and Endodontics, AB Shetty Memorial Institute of Dental Sciences, Mangalore 575018, Karnataka, India (e-mail: shalinannsaji91@gmail.com).

Abstract

Objective To assess the patients' knowledge and perceptions regarding endodontic treatment.

Materials and Methods A questionnaire-based survey was conducted in a group of 400 patients, consisting of questions based on sociodemographic details as well as their knowledge about endodontic treatment. The collected data was tabulated and analyzed using methods of descriptive analysis.

Results The majority of the respondents stated their level of knowledge regarding endodontic treatment as average.

Conclusion Within the limitations of the study, an improvement in the knowledge regarding endodontic treatment among patients was observed.

Keywords

- Fear
- Pain
- Questionnaire
- Root canal treatment

Clinical Significance Population-based surveys are useful for evaluating the community and patients' mentality toward endodontic treatment. Proper information regarding the treatment modalities can be explained to the patient based on their lack of information, which is of utmost importance to develop a positive approach to root canal treatment.

Introduction

Root canal treatment (RCT) is a procedure that involves the removal of damaged and diseased dental pulp and replacement by a sterile and inert substance.¹ One of the most common endodontic procedures for which a patient visits the dentist is RCT. In routine dental practice, patients' awareness and knowledge regarding endodontic treatment play a significant role, as it influences the course and effects of treatment.² Patients tend to delay or avoid dental treatments due to the fear of previously experienced pain. Apart from this, other reasons affect the choice of treatment plans like the patient's socioeconomic status, attitude, and prefer-

ences. These patients are more prone to dental decay, have more missing teeth, and have less functioning teeth, according to studies, owing to inadequate oral health maintenance.³ A survey conducted by Janczarek et al found that patients' knowledge and awareness of endodontic treatment has improved and that the majority of patients surveyed are aware of the options available to them during endodontic treatment, including different types of anesthesia, RCT, modern techniques, and treatment outcome prediction.⁴

Another survey conducted by Sisodia et al reported a moderate level of awareness of endodontic treatment among the selected population. It also emphasized the significance of providing patients more information about the benefits of

DOI <https://doi.org/10.1055/s-0041-1736451>.
ISSN 2582-4287.

© 2021. Nitte (Deemed to be University). All rights reserved.
This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

retaining teeth through endodontic therapy.¹ Dental apprehension and expectation of pain had an imperative role on the patient's ability to comprehend the information given.² Recognizing the reasons that result in distress among patients, consequently hindering them from undertaking RCT, is necessary. The population-based survey is a key step to adequately address the issue.⁵ This survey was conducted to evaluate the patients' knowledge and perceptions regarding endodontic treatment in an Indian subpopulation.

Materials and Methods

Institutional ethical clearance was obtained (ABSM/EC 22/2019 dated 28/06/2019). All the participants signed an informed consent declaration.

A self-administered questionnaire was prepared and distributed among 400 patients who had visited the department of conservative dentistry and endodontics. The inclusion criteria for the study were patients belonging to the age group of 18 to 65 years, who agreed to participate in the study. People associated with the field of dentistry were excluded. A pilot study was conducted to validate the questionnaire. A multiple-choice questionnaire consisting of sociodemographic details as well as their knowledge and perceptions about endodontic treatment was distributed (► **Table 1**). The questionnaire was prepared in Kannada, Malayalam, and English. Most of the respondents belonged to the south Canara district of Karnataka and the northern districts of Kerala.

Statistical Analysis

Data analysis was performed using methods of descriptive statistics like frequency and percentages.

Results

This is a simple observational study. The findings were tabulated, and the results were analyzed to find out patients' awareness about RCT. Results obtained are summarized in ► **Tables 2 and 3**.

Discussion

The study group comprised 400 subjects in the age group of 18 to 65 years. Out of those, 224 (56%) participants were females and 176 (44%) were males. Out of the total participants, 30% were from urban areas, 46% from semiurban, and 24% from rural areas. As much as 12% of them had primary education, 52% with secondary education, and 36% with higher education. People who were employed accounted for 50% of the participants, 34% were students, and 16% were unemployed (► **Table 2**).

Responses to questions regarding knowledge and perceptions about endodontic treatment are summarized in ► **Table 3**. While considering the experiences of patients regarding endodontic treatment, 68% of the participants had a history of RCT. Out of the total participants, 52% had given

the response that RCT is not painful. Majority of the patients (56%) report to the dentist instead of trying home remedies in case of toothache. Out of those who have tried home remedies, 62% report that home remedies did not help in relieving toothache.

The major concern associated with RCT accounted for pain (42%), followed by the need for extraction despite undertaken treatment (26%). As much as 20% of the participants were concerned about the need for multiple X-rays and 10% about long treatment time. When participants were asked to identify the symptoms that require RCT, majority of the responses were strong spontaneous toothache (62%), followed by pain during biting (16%). As much as 10% of the participants drew attention to pain during chewing and 8% reported discolored tooth. The majority of the respondents define their level of knowledge about RCT as average and ask for details. (32%). When asked if RCT is indicated for any toothache, only 36% agreed, while the remaining did not agree. A total of 88% of the participants agreed that medicines alone cannot resolve endodontic infections without the need for RCT.

The participants were asked about the factor that determines them to choose the place for RCT. The majority of the responses were professional staff (54%), followed by painless treatment (20%). Both free of cost treatment and easy access to the place accounted for 10% of the responses, followed by 6% of the respondents favoring friendly service (► **Table 3**).

Endodontics has been one of the fastest-growing branches of dentistry. The predictable outcome of the treatment has been significantly influenced by the use of advanced methods.⁴ RCT is performed primarily for irreversible diseases of the pulp, which is the most common reason for seeking dental treatment.² A study conducted by Klages et al demonstrates that patients visiting the dental office expect pain usually rather than experiencing during treatment.⁶

People who are related to the field of dentistry have been excluded to make sure that the responses were either based on their previous experiences or expectations based on what the public hears about RCT. While analyzing the results of this survey, the majority of the respondents had undergone RCT. Most of them are aware that RCT is not painful. Patients can benefit from the use of local anesthesia and nitrous oxide, based on various recent reports, enhancing its effect. Preoperative intravenous (IV) sedatives and analgesics can be used as another form of premedication. General anesthesia is the alternative in extreme cases.⁷

A study conducted by Armfield et al reveals that patients avoid visits to the dentist, due to fear.⁸ Most of the surveyed patients admit that they immediately report to the dentist in case of spontaneous ailments in the mouth, and others tried home remedies. Respondents who had tried home remedies mostly opt for self-prescribed medicines. Natural home remedies like the use of warm saline rinses, clove oil dressings, and the use of cold compresses were also followed by some patients. In this survey, most of the patients' concerns regarding RCT were pain followed by the need for extraction despite treatment undertaken. Patients were also concerned

Table 1 Detailed questionnaire

Sl. no.	Questions	Options	
1.	Age	a) 18–24 years	b) 25–34 years
		c) 35–44 years	d) 45–54 years
		e) 55–65 years	
2.	Gender	a) Male	b) Female
3.	Location	a) Urban	b) Semi-urban
		c) Rural	
4.	Qualification	a) Primary education b) Secondary education c) Higher education	
5.	Occupation	a) Employed	b) Unemployed
		c) Student	
6.	Have you ever undergone RCT?	a) Yes	b) No
7.	Do you think root canal treatment is painful?	a) Yes	b) No
8.	Have you ever tried any home remedies to relieve toothache?	a) Yes	b) No
9.	What home remedies have you tried?	a) Warm saltwater rinses b) Use of cold compresses c) Dressings with clove oil d) Self-prescribed medicines e) Others	
10.	Have you got any relief after trying home remedies?	a) Yes	b) No
11.	What concerns do you have regarding RCT?	a) Pain b) Need for multiple X-rays c) Need for extraction despite undertaken treatment d) Long treatment time e) Others	
12.	What symptoms indicate the need to undergo RCT?	a) Pain during chewing b) Pain during biting c) Strong spontaneous toothache d) Discolored tooth e) Others	
13.	Do you know anything about RCT?	a) I know a lot b) Average knowledge, I ask for details c) Average knowledge, am not interested in the course d) I do not know anything e) Knowledge from media	
14.	Does any toothache indicate RCT to be done?	a) Yes	b) No
15.	Can medicines alone resolve root canal infections without the need for RCT?	a) Yes	b) No
16.	What factor determines you to choose the place for RCT?	a) Free of cost treatment b) Easy access to the place c) Professional staff d) Friendly service e) Painless treatment	

Abbreviation: RCT, root canal treatment.

about the need for multiple X-rays and a long treatment time.

The majority of the respondents agreed that strong spontaneous toothache indicates the need for RCT. In the present study, only 18% of the respondents did not know RCT, whereas a study conducted by Habib et al stated that 25.3% of the respondents did not know RCT.⁹ Patients are

also aware that RCT is not indicated for all cases of toothache. While considering the patients' awareness about the adequacy of medicines in treating endodontic infections, the majority of them agreed that they are not enough in treating endodontic infections. This indicates that the majority of the studied sample recognize the importance of endodontic treatment, as antibiotics should be prescribed only in

Table 2 Responses for sociodemographic details

Question	Response	Frequency	Percent
Age	a) 18–24 years	120	30.0
	b) 25–34 years	144	36.0
	c) 35–44 years	56	14.0
	d) 45–54 years	40	10.0
	e) 55–65 years	40	10.0
Gender	a) Male	176	44.0
	b) Female	224	56.0
Location	a) Urban	120	30.0
	b) Semiurban	184	46.0
	c) Rural	96	24.0
Qualification	a) Primary education	48	12.0
	b) Secondary education	208	52.0
	c) Higher education	144	36.0
Occupation	a) Employed	200	50.0
	b) Unemployed	64	16.0
	c) Student	136	34.0

Table 3 Responses to questions regarding knowledge and perceptions about endodontic treatment

Question	Response	Frequency	Percent
Have you ever undergone RCT?	a) Yes	272	68.0
	b) No	128	32.0
Do you think RCT is painful?	a) Yes	192	48.0
	b) No	208	52.0
Have you ever tried any home remedies to relieve toothache?	a) Yes	176	44.0
	b) No	224	56.0
What home remedies have you tried?	a) Warm saltwater rinses	120	30.0
	b) Use of cold compresses	40	10.0
	c) Dressings with clove oil	56	14.0
	d) Self-prescribed medicines	160	40.0
	e) Others	24	6.0
Have you got any relief after trying home remedies?	a) Yes	152	38.0
	b) No	248	62.0
What concerns do you have regarding RCT?	a) Pain	168	42.0
	b) Need for multiple X-rays	80	20.0
	c) Need for extraction despite undertaken treatment	104	26.0
	d) Long treatment time	40	10.0
	e) Others	8	2.0
What symptoms indicate the need to undergo RCT?	a) Pain during chewing	40	10.0
	b) Pain during biting	64	16.0
	c) Strong spontaneous toothache	248	62.0
	d) Discolored tooth	32	8.0
	e) Others	16	4.0

(Continued)

Table 3 (Continued)

Question	Response	Frequency	Percent
Do you know anything about RCT?	a) I know a lot	96	24.0
	b) Average knowledge, I ask for details	128	32.0
	c) Average knowledge, am not interested in the course	56	14.0
	d) I do not know anything	72	18.0
	e) Knowledge from media	48	12.0
Does any toothache indicate RCT to be done?	a) Yes	144	36.0
	b) No	256	64.0
Can medicines alone resolve root canal infections without the need for RCT?	a) Yes	48	12.0
	b) No	352	88.0
What factor determines you to choose the place for RCT?	a) Free of cost treatment	40	10.0
	b) Easy access to the place	40	10.0
	c) Professional staff	216	54.0
	d) Friendly service	24	6.0
	e) Painless treatment	80	20.0

Abbreviation: RCT, root canal treatment.

specific cases with systemic infection and as prophylaxis for other medical conditions.¹⁰ The knowledge regarding endodontic treatment is not only due to school education but also due to previous visits to the dental clinic, communication with peers, and the influence of mass media—television, newspapers, and Internet.^{2,4} Majority of the surveyed patients choose the place for RCT based on the services provided by professional staff. Iqbal et al in their study concluded that patients choose the dental service provider based on professional experience, staff courtesy, and friendly environment.¹¹

Over the last years, the knowledge regarding endodontic treatment among patients has been significantly increased. Patients are not just aware of the immediate pain relief but also about the prospects offered by the modern clinics, carrying out of the treatment at the highest level and adequate postoperative care by ensuring long-term radiological surveillance, allowing observation of results attained, aiding immediate intervention if required and monitoring the treatment effects.^{12,13}

Conclusion

Within the limitations of the study, an improvement in the knowledge regarding endodontic treatment among patients was observed. Knowledge and awareness of patients about endodontic treatment differ among races and populations. The majority of the surveyed patients are aware of the advanced techniques available for endodontic treatment.

Clinical Significance

Questionnaire-based surveys in the population help evaluate the community and assess the patient's mentality

toward RCT. It is of utmost importance, especially in developing countries like India, where oral health has been neglected by the people, mainly due to inadequate dental health knowledge or myths about the treatment. Proper information regarding the treatment modalities can be explained to the patient based on their lack of information, which is of utmost importance to develop a positive approach to RCT. The importance of saving natural teeth should be emphasized and misperceptions about endodontic treatment should be dissipated. Improving awareness in patients regarding endodontic treatment will change the patients' attitude and make them accept the treatment.

Conflict of interest

None declared.

References

- 1 Sisodia N, Yadav S, Nangia T, et al. Dental patients' knowledge and attitude towards endodontics – A survey. *Pharm J Sci Biomed* 2015;5:80–83
- 2 Purra AR, Sajad M, Ahangar FA, et al. Patient's awareness and knowledge of the root canal treatment in Kashmiri Population: a survey- based original research. *International Journal of Contemporary Medical Research*. 2018;5(07):G12–G15
- 3 Aldawsari M, Alamri HM. Public knowledge and perception regarding endodontic treatment in a Saudi population. *J Int Oral Health* 2017;9:255–257
- 4 Janczarek M, Cieszko-Buk M, Bachanek T, et al. Survey-based research on patients' knowledge about endodontic treatment. *Zdrow Publiczne* 2014;124:134–137
- 5 Bansal R, Jain A. An insight into patient's perceptions regarding root canal treatment: A questionnaire-based survey. *J Family Med Prim Care* 2020;9(02):1020–1027
- 6 Klages U, Ulusoy O, Kianifard S, Wehrbein H. Dental trait anxiety and pain sensitivity as predictors of expected and experienced

- pain in stressful dental procedures. *Eur J Oral Sci* 2004;112(06): 477–483
- 7 Montagnese TA. Why intravenous moderate sedation should be taught in graduate endodontic programs. *J Dent Educ* 2012;76(03):288–290
 - 8 Armfield JM, Heaton LJ. Management of fear and anxiety in the dental clinic: a review. *Aust Dent J* 2013;58(04):390–407
 - 9 Habib AA, Dormant MD, Al Saysd T, et al. Dental patient's knowledge and awareness about root canal treatment in Syrian population: Survey-based research. *IJRSR* 2017;8:20583–20586
 - 10 Aminoshariae A, Kulild JC. Evidence-based recommendations for antibiotic usage to treat endodontic infections and pain: a systematic review of randomized controlled trials. *J Am Dent Assoc* 2016;147(03):186–191
 - 11 Iqbal M, Jameel A, Girach MM, et al. Factors affecting patient's choice of dental services. *PODJ* 2014;34:691–695
 - 12 Parirokh M, V Abbott P. Various strategies for pain-free root canal treatment. *Iran Endod J* 2014;9(01):1–14
 - 13 Murray CA, Saunders WP. Root canal treatment and general health: a review of the literature. *Int Endod J* 2000;33(01):1–18