Y. FUKUDA, K. UTIMOTO* (KYOTO UNIVERSITY, JAPAN)

Effective Transformation of Unactivated Alkynes into Ketones or Acetals with a Gold(III) Catalyst

J. Org. Chem. **1991**, 56, 3729–3731, DOI: 10.1021/jo00011a058.

Gold(III)-Catalyzed Hydration of Alkynes

Significance: In 1991, Fukuda and Utimoto disclosed the gold(III)-catalyzed hydration of alkynes. In contrast to traditional Hg(II)-mediated hydrations that require strongly acidic conditions, the reported Au(III)-catalyzed protocol is much milder and does not require any acid, allowing for increased functional group tolerance.

Comment: The hydration reaction is operationally simple, requiring only 2 mol% of Au(III) catalyst in a refluxing mixture of methanol and water. When strictly anhydrous methanol is used, the corresponding dimethyl acetal product is obtained in place of the ketone.

Review: Z. Li, C. Brouwer, C. He *Chem. Rev.* **2008**, 108, 3239–3265.

Category

Metals in Synthesis

Key words

hydration

alkynes

gold catalysis

Synfact
Classic

SYNFACTS Contributors: Mark Lautens, Jeanne Masson-Makdissi

Synfacts 2022, 18(01), 0063 Published online: 17.12.2021

DOI: 10.1055/s-0041-1737317; Reg-No.: L16022SF

© 2022, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany