Sharp Needle Recanalization: A Salvage Procedure for Failing Arteriovenous Fistula Due to Tight Uncrossable Venous Stenosis

Sultan R. Alharbi¹

¹Department of Radiology and Medical Imaging, College of Medicine, King Saud University, Riyadh, Kingdom of Saudi Arabia

Arab J Intervent Radiol 2021;5:114–115.

Address for correspondence Sultan R. Alharbi, MD, Department of Radiology and Medical Imaging, College of Medicine, King Saud University, P.O. Box 7805, Riyadh 11472, Kingdom of Saudi Arabia (e-mail: drsultan000@gmail.com).

A 33-year-old male presented with malfunctioning brachiocephalic arteriovenous fistula (AVF) due to short tight juxta anastomotic stenosis (►Fig. 1). Access was gained into the AVF draining vein, and multiple trials to cross the

tight stenosis using catheter and wire failed (►Fig. 2). Sharp needle recanalization was performed under ultrasound guidance, using 21G needle and wire to cross this tight stenosis (►Fig. 3a,b). Wire passed into the AVF

Fig. 1 Ultrasound of arteriovenous fistula (AVF) showing a juxta anastomotic tight stenosis and aneurysmal dilatation.

Fig. 2 Fistulogram showing patent arteriovenous fistula (AVF) with uncrossable tight stenosis.

Fig. 3 (a and b) Ultrasound-guided sharp needle recanalization using 21G needle and wire.

published online
December 22, 2021

DOI <https://doi.org/10.1055/s-0041-1739304>.
ISSN 2542-7075.

© 2021. The Pan Arab Interventional Radiology Society. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

Fig. 4 Postballoon angioplasty fistulogram showing successful angioplasty without contrast extravasation.

Fig. 5 Follow-up fistulogram after 2 years showing patent arteriovenous fistula (AVF).

draining vein and snared through the already placed vascular sheath. Balloon angioplasty was performed using 6-mm, high-pressure balloon and inflated for 3 minutes. Postangioplasty venogram showed successful result (► **Fig. 4**). Patient had follow-up fistulogram and angio-

plasty every 6 months for 2 years, and fistula remained patent (► **Fig. 5**).

Sharp needle recanalization is a well-known technique for central venous occlusion.¹ This technique is rarely used in peripheral veins.² Jump graft is a salvage technique used to bridge stenotic/thrombotic venous segment of AVF. Jump graft is created surgically or percutaneously by a mean of stent graft for failing AVF.³ Sharp needle recanalization is a minimally invasive alternative salvage technique to surgical jump graft in tight uncrossable AVF peripheral venous stenosis.⁴

Financial Support and Sponsorship

Nil.

Conflicts of Interest

None declared.

References

- 1 Arabi M, Ahmed I, Mat'hami A, Ahmed D, Aslam N. Sharp central venous recanalization in hemodialysis patients: a single-institution experience. *Cardiovasc Intervent Radiol* 2016;39(06): 927–934
- 2 Miller GA, Koh E, Khariton A, et al. Sharp needle recanalization for salvaging hemodialysis accesses with chronically occluded peripheral outflow. *J Vasc Access* 2012;13(01):22–28
- 3 Timpone VM, Gover DD. Percutaneous creation of a jump bypass graft in a native arteriovenous hemodialysis fistula. *J Vasc Interv Radiol* 2011;22(05):734–736
- 4 Noh SY, Goo DE, Kim YJ, Yang SB, Lee JM, Lee WH. Sharp needle recanalization technique for peripheral hemodialysis arteriovenous fistula occlusion. *Cardiovasc Intervent Radiol* 2021 (e-pub ahead of print). Doi: 10.1007/s00270-021-02809-1