



Correlation between Noncompliance Behavior and Biochemical Parameters of Chronic Kidney Disease Patients Undergoing Hemodialysis

Sarita Telma Fernandes¹ Sushma Marita Dsouza²

¹Department of Community Health Nursing, Nitte Usha Institute of Nursing Sciences, Deralakatte, Mangalore, Karnataka, India

²Access Health International, Delhi, India

Address for correspondence Sushma Marita Dsouza, MPH, Access Health International, House No. 275, Behind Bank of India, Shri Ram Nagar, G.T. Road, Shahdara, East Delhi, Delhi 110032, India (e-mail: sushmadsouza22@gmail.com).

J Health Allied Sci^{NU} 2022;12:258–262.

Abstract

Background and Objectives Compliance with dietary, fluid, and medication instructions is a critically significant factor for the health and well-being of patients undergoing hemodialysis for an extended period. The current study assessed the correlation between noncompliance behavior and biochemical parameters of chronic kidney disease (CKD) patients undergoing hemodialysis. The study's objectives were 1) to assess patients' noncompliance behavior using a questionnaire, 2) to determine the biochemical parameters of CKD patients having noncompliance behavior, and 3) to find the correlation between biological parameters and noncompliance behavior.

Methods A cross-sectional study was conducted from July 12, 2018, to February 10, 2019. A purposive sampling technique was used to recruit the participants, and 100 participants were included in the study. The End-Stage Renal Disease–Adherence Questionnaire was used to assess the non-compliance behavior.

Results Most dialysis patients (70%) were compliant with the therapeutic regime, while 30% failed to comply. The study revealed a significant positive correlation between noncompliance behavior and biochemical parameters of CKD patients undergoing hemodialysis ($r = 0.578$, $p < 0.05$).

Conclusion Noncompliance behavior is significantly correlated with the biochemical parameters among patients with CKD on dialysis. Optimal health is possible through fluid and dietary compliance in these patients. Hence, adherence behavior plays a significant role in the health and recovery of dialysis patients with CKD.

Keywords

- biological parameter
- chronic kidney disease
- correlation
- hemodialysis
- noncompliance behavior

Introduction

In the body, healthy kidneys purify the blood and eliminate the extra fluid and the waste in the form of urine. Moreover, they keep our bodies healthy. Chronic kidney disease (CKD) is a slowly progressive disease of kidney function, character-

ized by a low glomerular filtration rate (GFR).¹ When kidneys fail to function, dialysis performs its function as replacement therapy for kidney failure. It involves eliminating waste products such as creatinine and urea and free water from the blood when the kidneys do not work regularly.²

published online
December 31, 2021

DOI <https://doi.org/10.1055/s-0041-1740023>.
ISSN 2582-4287.

© 2021. Nitte (Deemed to be University). All rights reserved.
This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

Hemodialysis is an intensive and expensive therapy and requires strict fluid and dietary restrictions. Consequently, repeated dialysis leads to a personal and financial burden to the person and their families.³ Even with the unceasing development of dialysis technology and pharmacological treatment, mortality rates are still high for dialysis patients. In India, nearly 9 to 13% of patients undergoing hemodialysis die within 1 year. Along with dialysis adequacy, the noncompliance behavior also affects the patient's survival in hemodialysis.⁴

Noncompliance is common among patients undergoing hemodialysis. Furthermore, noncompliance behavior varies from patient to patient with different treatment regimens which includes adherence to drugs, fluid, treatment, and dietary restrictions.^{5–7} Failure to adhere to dietary restriction leads to faster progression of kidney disease into kidney failure and eventually results in kidney transplants. The shreds of evidence from studies show that failure to comply with dietary restrictions relates to renal disease progression.⁸

When CKD patients start hemodialysis, their entire life changes; they should regularly travel to dialysis centers to take prescribed medications without fail, and they have to modify their dietary patterns. The successful treatment of renal disease will largely depend on the patient's adherence behavior.^{9,10}

The most common data sources that indicate adherence to medications and restrictions on diet and fluid intake are serial values of the biochemical index, which have straight implications on health outcomes.¹¹ The often-used parameters are interdialytic weight gain (IWG), as an indicator of fluid compliance, predialysis blood urea nitrogen (BUN), and serum potassium (K) as measures of dietary compliance and serum phosphate (PO₄) as a parameter for medication compliance.^{12,13} As compared with other compliance measures, such as patient self-reports and staff reports, these parameters are generally considered to be more consistent, less biased, and objective.^{14,15}

The current study was conducted to assess patients' noncompliance behavior with regard to medication, fluid, diet, and dialysis and determine the biochemical variations in patients with noncompliance behavior and its correlation. Here, noncompliance is referred to as failure to maintain the dietary, drugs, and fluid restrictions prescribed for the CKD patients undergoing hemodialysis.

Materials and Methods

Study Design and Participants

A cross-sectional study was conducted among 100 dialysis patients (universal samples) who fulfilled the inclusion criteria. Subjects were selected from three different settings namely a tertiary care hospital in Mangaluru comprising 12 dialysis beds, dialysis centers in Karkala (6 beds) and Puttur (5 beds). The study's inclusion criteria were CKD patients aged 18 years and above, undergoing hemodialysis for more than 3 months, and screened for noncompliance behavior. The study's exclusion criteria were patients undergoing hemodialysis for the first time and critically ill patients.

Sample Size Calculation

The sample size was calculated using the statistical formula for estimating a single proportion. It was expected that 50% of CKD patients undergoing dialysis were noncompliant with dietary and fluid restriction with 10% relative precision with a 95% confidence interval. The estimated sample size was $n = 100$.

Ethical Consideration

The study was reviewed and approved on April 20, 2018, by the Institutional Review and Ethics Committee. The registration number was NUINS/CON/NU/IEC/2018–19. A subject information sheet was provided with a clear explanation about the study purpose and written informed consent was obtained before including the study participants. Confidentiality was assured to the entire subjects. They were instructed to sign the consent form, fill the demographic proforma, and End-Stage Renal Disease–Adherence Questionnaire (ESRD–AQ).

Instruments

The following tools were used to collect the data: (1) demographic proforma, which comprised 10 items such as age, gender, religion, educational status, marital status, family income, diagnosis of the sample, duration of dialysis, and the number of times he/she has undergone hemodialysis; (2) ESRD–AQ questionnaire, which refers to the End-Stage Renal Disease–Adherence Questionnaire. It is a standardized tool developed to test the treatment adherence of people undergoing hemodialysis; hemodialysis attendance, medications, fluid restrictions, and diet prescription.¹⁶

Data Collection

A purposive sampling technique was used to recruit the participants to the study. Formal written permission was obtained from the Medical Superintendent of the respective hospital and dialysis centers. The data collection period was from July 12, 2018, to February 10, 2019. Homogenous time sessions were followed for the data collection.

Data Analysis

The data were analyzed using Statistical Package for Social Sciences 20 (SPSS 20) software. The data were analyzed in terms of objectives and hypotheses of the study using both descriptive and inferential statistics. Data on noncompliance behavior and biochemical parameters are expressed in frequency and percentage. Pearson's correlation coefficient (r) was used to find the relationship between biochemical parameters and noncompliance behavior. A p -value < 0.05 was considered significant.

Results

A total of 100 patients were selected for the study. The mean age of the patients was 54.15 ± 14.29 (standard deviation [SD]). The distribution of subjects based on gender revealed that most dialysis patients were males (77%). The majority of dialysis patients had high school education (63%), 15% had

Table 1 Distribution of demographic characteristics

Variables (n = 100)	Frequency (n)	Percentage (%)
Age (y)		
<20	1	1
20–50	42	42
>50	57	57
Gender		
Male	77	77
Female	23	23
Religion		
Hindu	79	79
Muslim	11	11
Christian	10	10
Education		
Primary school	11	11
High school	63	63
Pre-degree	15	15
Diploma	6	6
Graduate	6	6
Marital status		
Single	6	6
Married	84	84
Divorced	1	1
Widow	9	9
Family income		
<5000	17	17
5000–10000	50	50
>10000	33	33
Duration of hemodialysis		
<1 y	33	33
1–5 y	67	67
Diagnosis		
CKD	100	100
No of dialysis per week	100	100

Abbreviation: n = number.

pre-degree education, 11% had primary school education, 6% had diploma education, and 6% had graduate education.

The distribution of subjects based on marital status revealed that most dialysis patients were married (84%), 9% were widows, and 6% were single. Based on family income, the distribution revealed that the majority (50%) had an income between 5,000 to 10,000. With regard to dialysis treatment duration, most dialysis patients (67%) underwent dialysis for a period of 1 to 5 years. The distribution of subjects based on the number of dialysis per week revealed that 100% of dialysis patients were undergoing dialysis twice a week (→Table 1).

Among the 100 subjects, 30% had noncompliance behavior (→Table 2).

→Table 3 shows the distribution of biochemical parameters of dialysis patients showing noncompliance behavior. The noncompliance behavior was assessed, and their biochemical parameters indicated that (43%) had a high BUN level. All patients had low hemoglobin (Hb) levels. More than half of the subjects (67%) had elevated serum potassium, (80%) had high serum phosphate levels, and (93%) of patients

Table 2 Distribution of noncompliance behavior score of dialysis patients

Behavior (n = 100)	Score range	Frequency (n)	Percentage (%)
Compliance behavior	More than 850	70	70
Noncompliance behavior	Less than 850	30	30

Table 3 Biological parameters of dialysis patients showing noncompliance behavior

Biological parameters (n = 30)	Frequency (n)	Percentage (%)
BUN		
2.5–7.1 mg/dL	17	57
>7.1 mg/dL	13	43
Hb		
<12 g/dL	30	100
Potassium		
3.5–5 mEq/L	10	33
>5 mEq/L	20	67
Phosphorous		
2.5–4.5 mg/dL	6	20
>4.5 mg/dL	24	80
Albumin		
<3.4 g/dL	2	7
3.4–5.4 g/dL	28	93
Weight gain		
2–4 Kg	27	90
>4 Kg	3	10

had shown average serum albumin (Alb) levels. Among the 100 subjects, (7%) had low serum albumin levels and (90%) of the patients had interdialytic weight gain (IWG) between 2 and 4 kg and 10% had IWG more than 4 kg.

The Karl Pearson correlation coefficient (*r*) value of non-compliance behavior and biochemical parameters of CKD was 0.578 ($p < 0.05$). A significant positive correlation between noncompliance behavior and biochemical parameters was observed among the dialysis patients with CKD (→Table 4).

Discussion

The present study showed that 30% of the subjects had noncompliance behavior. A study conducted by Lee et al

Table 4 Correlation between noncompliance and biological parameters

Variables (n = 30)	Correlation (r value)	Level of significance
Noncompliance behavior	0.578	0.01 ($p < 0.05$)
Biological parameters		

revealed that dietary and fluid compliance was observed among 35.5% and 40.3% of patients, respectively.¹⁷ Our study reported that 37.8% had compliance with diet and 42.6% had compliance with fluid. Kugler et al revealed that non-adherence to diet and fluid restrictions were 81.4% and 74.6%, respectively.¹⁸ Ahrari et al reported that 41.1% of patients reported moderate nonadherence to diet, 45.2% reported moderate nonadherence to fluid.⁹ Contrary to the above studies, Bame et al reported that compliance with the diet regimen was 90% and 49.5% of patients were compliant with the fluid-restriction regimen in a hospital-based study.¹⁹ Such variations in the prevalence of dietary non-compliance or compliance may be accredited to the variations in the indicators and standards used for measuring compliance. In most of the studies in the past, dietary compliance was measured using serum K as the sole indicator.¹⁷ However, we used serum K, Alb, Hb, and BUN parameters concurrently as an indicator of dietary compliance. Lee et al used serum K and phosphate levels as an indicator of dietary compliance.¹⁷

The serial values of biochemical measures serve as familiar sources of data to assess adherence to medications, diet and fluid restrictions, directly impacting the health outcomes.¹¹ The current study showed that around 43% of patients had a high BUN level, 7% had low albumin levels, 67% had elevated serum potassium, and 90% had IWG between 2 kg and 4 kg. A study has predicted that serum albumin levels as one of the sturdiest predictors in laboratories for mortality in dialysis patients as it indicates the nutritional status and the level of systemic inflammation.²⁰ In the present study, out of 30% of dialysis patients showing noncompliance behavior, 28% of patients had serum albumin levels between 3.4 and 5.4 g/dL, which was slightly elevated. There was a strong association between comorbidity and hypoalbuminemia among dialysis patients. Furthermore, a tendency of decreased albumin concentration within 6 months prior to the patient's death was present.²¹

There was a positive correlation observed between non-compliance behavior and biochemical parameters among dialysis patients. Rambod et al had done a study to assess the factors of adherence in uremic patients on hemodialysis. BUN, serum potassium, serum phosphate levels, and IWG were collected and analyzed. There was a significant relationship between the biochemical parameters and IWG with educational status ($p \leq 0.05$).²²

The study concludes that dietary compliance may be improved using nutritional counseling techniques and education, and motivating patients to modify and comply with dietary recommendations. The study findings have suggestions to all the healthcare workers, including clinicians, nurses, and all other medical staff in the dialysis unit, to promote counseling to the dialysis patients. This study was limited to hemodialysis patients, and the purposive sampling technique was used to select participants, which imposed limits in a larger generalization.

The present study shows that there is a significant relationship between noncompliance behavior and biochemical

parameters. To improve the compliance behavior among these patients, nurses have to be empowered as counselors, providing health education and counseling the patients on proper diet, fluid intake, and maintaining health-compliance behavior. Based on the current study, a researcher can develop further studies to progress toward new problem-solving and preventive strategies.

Conclusion

Noncompliance among dialysis patients is a significant problem even though it is said to be inversely related to survival rates. Hemodialysis requires patients to commit considerable time to their treatment, comply with strict dietary and fluid restrictions, and take medication regularly. This study reveals a significant relationship between the noncompliance behavior and biochemical parameters of CKD patients. Periodic counseling and constant motivation can improve patients' confidence in fluid, dietary, and medication compliance leading to a healthy life.

Funding

This study received funding from Nitte (Deemed to be University) and the grant number was NUSR2/2018/10/70.

Conflict of Interest

None declared.

Acknowledgments

The authors would like to thank all the CKD patients undergoing hemodialysis who participated in this study. We want to thank K. S. Hegde Medical Academy (KSHEMA) hospital, dialysis centers in Karkala and Puttur, Nitte Usha Institute of Nursing Sciences, and Nitte (Deemed to be University) for their support.

References

1. Weis L, Metzger M, Haymann J-P, et al; NephroTest Study Group. Renal function can improve at any stage of chronic kidney disease. *PLoS One* 2013;8(12):e81835. Doi: 10.1371/journal.pone.0081833
2. Krans B. Dialysis. 2019. Accessed November 1, 2021 at: www.healthline.com/health/dialysis
3. Mollaoğlu M. Quality of life in patients undergoing hemodialysis. Chapter 37. Intech Open; 2013:823–843. Available at: <http://dx.doi.org/10.5772/45929>
4. Chandrashekar A, Ramakrishnan S, Rangarajan D. Survival analysis of patients on maintenance hemodialysis. *Indian J Nephrol* 2014;24(04):206–213
5. Wolcott DL, Maida CA, Diamond R, Nissenson AR. Treatment compliance in end-stage renal disease patients on dialysis. *Am J Nephrol* 1986;6(05):329–338
6. Lohman-Adham M. Medication noncompliance in patients with chronic disease: issues in dialysis and renal transplantation. *Am J Manag Care* 2003;9(02):155–171
7. Leggat JE Jr, Orzol SM, Hulbert-Shearon TE, et al. Noncompliance in hemodialysis: predictors and survival analysis. *Am J Kidney Dis* 1998;32(01):139–145
8. Lentine K, Wrone EM. New insights into protein intake and progression of renal disease. *Curr Opin Nephrol Hypertens* 2004;13(03):333–336

- 9 Ahrari S, Moshki M, Bahrami M. The relationship between social support and adherence of dietary and fluids restrictions among hemodialysis patients in Iran. *J Caring Sci* 2014;3(01):11–19
- 10 Beerendrakumar N, Ramamoorthy L, Haridasan S. Dietary and fluid regime adherence in chronic kidney disease patients. *J Caring Sci* 2018;7(01):17–20
- 11 Arici M, Altun B, Usalan C, et al. Compliance in hemodialysis patients: unanticipated monitoring of biochemical indices. *Blood Purif* 1998;16(05):275–280
- 12 Ferraro KF, Dixon RD, Kinlaw BJR. Measuring compliance among in-center hemodialysis patients. *Dial Transplant* 1986; 15:226–236
- 13 Simmens S, Kimmel PL, Kobrin S, Reiss D. Multidimensional assessment of compliance in two dialysis units. *J Am Soc Nephrol* 1991;2:351
- 14 Blackburn SL. Dietary compliance of chronic hemodialysis patients. *J Am Diet Assoc* 1977;70(01):31–37
- 15 Hitchcock PB, Brantley PJ, Jones GN, McKnight GT. Stress and social support as predictors of dietary compliance in hemodialysis patients. *Behav Med* 1992;18(01):13–20
- 16 Kim Y, Evangelista LS, Phillips LR, Pavlish C, Kopple JD. The End-Stage Renal Disease Adherence Questionnaire (ESRD-AQ): testing the psychometric properties in patients receiving in-center hemodialysis. *Nephrol Nurs J*. 2010;37(04):377–393
- 17 Lee SH, Molassiotis A. Dietary and fluid compliance in Chinese hemodialysis patients. *Int J Nurs Stud* 2002;39(07):695–704
- 18 Kugler C, Vlaminc H, Haverich A, Maes B. Nonadherence with diet and fluid restrictions among adults having hemodialysis. *J Nurs Scholarsh* 2005;37(01):25–29
- 19 Bame SI, Petersen N, Wray NP. Variation in hemodialysis patient compliance according to demographic characteristics. *Soc Sci Med* 1993;37(08):1035–1043
- 20 Yeun JY, Kaysen GA. Factors influencing serum albumin in dialysis patients. *Am J Kidney Dis* 1998;32(6, Suppl 4):S118–S125
- 21 Kaysen GA. Serum albumin concentration in dialysis patients: why does it remain resistant to therapy? *Kidney Int Suppl* 2003; 64(87):S92–S98
- 22 Rambod M, Peyravi H, Shokrpour N, Sareban MT. Dietary and fluid adherence in Iranian hemodialysis patients. *Health Care Manag (Frederick)* 2010;29(04):359–364