

# Does the Chemical Diversity of the Order Haplosclerida (Phylum Porifera: Class Demospongia) Fit with Current Taxonomic Classification?

## Authors

Marie-Aude Tribalat<sup>1</sup>, Maria V. Marra<sup>2</sup>, Grace P. McCormack<sup>2</sup>, Olivier P. Thomas<sup>1,3</sup>

## Affiliations

<sup>1</sup> Géoazur UMR Université Nice Sophia Antipolis-CNRS-IRD-OCA, Valbonne, France

<sup>2</sup> National University of Ireland Galway, Zoology, School of Natural Sciences, University Road, Galway, Ireland

<sup>3</sup> National University of Ireland Galway, Marine Biodiscovery, School of Chemistry, University Road, Galway, Ireland

## Key words

- sponges
- Haplosclerida
- alkaloids
- polyacetylenic
- chemotaxonomy

received Nov. 29, 2015  
revised March 8, 2016  
accepted March 22, 2016

## Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-105879>  
Published online May 2, 2016  
Planta Med 2016; 82: 843–856  
© Georg Thieme Verlag KG  
Stuttgart · New York ·  
ISSN 0032-0943

## Correspondence

**Prof. Grace P. McCormack**  
Ryan Institute  
School of Natural Sciences  
National University of Ireland  
University Road  
Galway  
Ireland  
Phone: + 35 3 91 49 23 21  
Fax: + 35 3 91 49 45 35  
[grace.mccormack@nuigalway.ie](mailto:grace.mccormack@nuigalway.ie)

## Correspondence

**Prof. Olivier P. Thomas**  
School of Chemistry  
National University of Ireland  
University Road  
Galway  
Ireland  
Phone: + 35 3 91 49 35 63  
Fax: + 35 3 91 49 55 76  
[olivier.thomas@nuigalway.ie](mailto:olivier.thomas@nuigalway.ie)

## Abstract

Sponges and their associated microbiota are well known to produce a large diversity of natural products, also called specialized metabolites. In addition to their potential use in the pharmaceutical industry, these rather species-specific compounds may help in the classification of some particular sponge groups. We review herein compounds isolated from haplosclerid sponges (Class Demospongia, Order Haplosclerida) in order to

help in the revision of this large group of marine invertebrates. We focus only on 3-alkylpyridine derivatives and polyacetylenic compounds, as these two groups of natural products are characteristic of haplosclerid species and are highly diverse. A close collaboration between chemists and biologists is required in order to fully apply chemotaxonomical approaches, and whenever possible biological data should include morphological and molecular data and some insight into their microbial abundance.

## Introduction

Sponges (Phylum Porifera) are sessile invertebrates distributed in most aquatic ecosystems. In marine areas like the Caribbean Sea, they may represent the largest substrate cover and/or the largest organic biomass of living organisms, thus contributing significantly to several nutrient cycles due to their outstanding filtering capabilities [1,2]. In addition, during their long evolutionary history, most, if not all, of the diverse species of this group (> 8000 described species to date) together with their associated microbiota have developed unique metabolic pathways leading to a huge diversity of natural products, also called specialized metabolites [3]. Taxonomic classification in this particular group of marine invertebrates is still highly challenging due to a paucity of morphological characters and a discrepancy between molecular and morphological data in many cases. For this reason, biochemical information has been recently used as a complementary tool (particularly within the framework of targeted or untargeted metabolomic approaches), leading to the recent concept of integrative systematics [4].

Within the Class Demospongiae, sponges belonging to the Order Haplosclerida are considered among the most prolific sources of bioactive

marine natural products, including alkaloids, polyacetylenes, or terpene derivatives. This group is also one of the most diverse of the sponge groups in terms of numbers of species and habitats, and its members also have few distinguishing morphological characteristics. The current classification of the order as outlined in Systema Porifera, based primarily on morphology, is comprised of three suborders [two marine (Haplosclerina and Petrosina) and one freshwater (Spongillina)]. The marine suborders together comprise six families; Callyspongiidae, Chalinidae, Niphatiidae, Petrosiidae, Phloeodictyidae, and Calcifibrospongiidae [5]. Analysis of sterol chemistry had indicated possible difficulties with their classification [6], suggesting patterns of relatedness that did not agree with morphological data, while in contrast, a review of 3-alkylpiperidine alkaloids appeared to agree with the current classification [7]. Subsequent molecular phylogenetic studies reveal an evolutionary history that is not completely compatible with Systema Porifera, indicating that the freshwater sponges belong elsewhere in the Demospongiae, and that while the marine Haplosclerida do form a clade, the suborders and the families (where there is enough data) are polyphyletic [8]. For this reason, a reassessment of the Order Haplosclerida is underway by multiple research groups, as an assessment of chemical

diversity may help in the development of a robust integrative classification of the group and, in return, current systematic studies in this group may also direct the search for related compounds of interest.

This review will focus on two major classes of specialized metabolites found in the Order Haplosclerida (3-alkylpyridine derivatives and polyacetylene derivatives) and will discuss them in view of the currently accepted classification for five of the six families described in *Systema Porifera* [5], with no compound being reported so far from sponges of the sixth family Calcifibrospongiidae [9]. Sterols and fatty acids have been reported for some members of this group in 1994 [6], and the large chemical diversity produced by this prolific sponge group was last collated in 1996 for 3-alkylpiperidine derivatives [7]. This review is not aimed to be exhaustive but discusses representatives of 3-alkylpyridine derivatives and polyacetylene derivatives. Given that they are both widely distributed amongst haplosclerid species, are largely restricted to this group, and are rather unique in the field of natural products, we consider that focusing our review on these two chemical families provides ample information to help identify discrepancies in the classification, areas for focus in the construction of a revised integrative classification and valuable avenues for future research focus.

Major issues with publishing descriptions of compounds from species that have not been fully identified will become apparent. This issue is relevant not only for the group of marine sponges included here, but for any organism of interest for bioprospecting. Accompanying morphological identification with a DNA sequence is a highly recommended way forward (in addition to deposition of a specimen voucher in a respected institute/museum). We intend this review to encourage better connections between chemists and biologists in an effort to identify novel compounds of interest while also striving to understand the origins and evolution of the same. A solid taxonomic description and classification will help in the rational collection of closely related species that will produce a large diversity of analogues of a targeted natural product family. Greater collaborative efforts will therefore increase avenues for identification of bioactive compounds, increase the likelihood of finding exploitable sources and perhaps even a biotechnological route for production. Knowledge of the complex comparative chemistry will assist in unravelling the controversial systematics specifically within the Order Haplosclerida, but will also lead to fruitful considerations in terms of chemotaxonomy within the phylum itself (Porifera, the sponges).

### Distribution of Simple 3-Alkylpyridine Derivatives in the Haplosclerida

Several 3-alkylpyridine derivatives have been isolated from the Haplosclerida, including in some cases the coupling of one or more similar moieties. This group of compounds is differentially distributed across Niphatidae, Callyspongiidae, Chalinidae, and Petrosiidae, but they are not yet reported from confirmed members of the Phloeodictyidae. The presence/absence of these classes of compounds in haplosclerid sponges are summarized in **Table 1**.

### Family Niphatidae

Within the family Niphatidae, an outstanding diversity of alkaloids has been isolated from members of the genus *Amphimedon*. Monomers of 3-alkylpyridines named hachijodines E–G as well as glycosylated monomers named amphimedosides A–E were isolated from *Amphimedon* sp. collected off Hachijo-Jima Island (Japan; **Fig. 1**) [10,11]. Additional mono- and dimers, pyrinodemins B–I, were reported from *Amphimedon* sp. found in the same area (Nakijin, Okinawa) along with a high number of dimers bearing a hydroxylamine moiety, starting with the bis-3-alkylpyridine derivative pyrinodemin A and nakinadines A–F (with a  $\beta$ -aminoacid) [12–16]. All of the compounds mentioned were found to exhibit cytotoxicity. *Niphates* sp. yielded some mono-3-alkylpyridine derivatives named niphatynes A–B, again from the Northwestern Pacific (Fiji; **Fig. 2**) [17], and Kobayashi's group more recently described some very close analogues named niphatines A–H from a Japanese sponge of the same genus (**Fig. 2**) [18,19]. Pyrinadines A–G have been isolated from *Cribrochalina* sp., also from the Japanese coast (**Fig. 2**) [20,21]. It is not known if the two *Amphimedons* or *Niphates* sp. mentioned here are the same species or two separate species in each case given that the specimens were collected in the same geographical area. Comparison of the specimen vouchers from the various studies will have to be performed to confirm if this is the case and preferably also comparison of DNA sequences.

It would appear that simple mono- and bis-3-alkylpyridine derivatives are restricted to species belonging to the genera *Amphimedon*, *Cribrochalina*, and *Niphates*, all belonging to the family Niphatidae (**Table 1**). While molecular data confirms that certain *Niphates* and *Amphimedon* species are related to each other, the position of *Cribrochalina* and *Pachychalina* in relation to other members of the Niphatidae family remains unresolved [22]. As more focus is placed on resolving haplosclerid taxonomy, and compounds are isolated from more fully described species, it will be interesting to confirm if this pattern remains.

Also, within the family Niphatidae, a dimeric cyclostelletamine (3-alkylpyridiniums) was isolated from the Northwestern Pacific sponge *Amphimedon compressa* [23]. Complex mixtures of polymeric halitoxins were isolated from several species present in the Northern Caribbean (*Haliclona rubens*, *H. viridis* and *H. erina*) that later were classified as *A. compressa*, *Amphimedon viridis*, and *Amphimedon erina* (**Fig. 3**) [24]. Later, analogous and bioactive polymers named amphitoxins were isolated as major constituents of *A. compressa* from the Caribbean [25,26]. Some trimeric 3-alkylpyridinium salts named niphatoxins A–B were isolated from *Niphates* sp. collected in the Red Sea by Talpir et al. (**Fig. 3**) [27]. No DNA sequence has been generated yet for *A. erina*, and while molecular data does not cluster *A. compressa* and *A. viridis* directly together, they are both placed in the same subgroup (Clade C), and are thus related to each other and with *Niphates* species [8,22]. The chemical content of the Southwestern Atlantic (Brazil) sponge *Pachychalina alcaloidifera* also includes the presence of antimicrobial and antimycobacterial cyclostelletamines A–K (**Fig. 3**) [28,29]. In addition, C–C and C–N bis- and tris-3-alkylpyridinium derivatives named pachychalines A–C were isolated from *Pachychalina* sp. collected in the Northwestern Atlantic (Caribbean; **Fig. 3**) [30]. Relationships of these *Pachychalina* species to each other and to other Haplosclerida have not yet been resolved using morphological and molecular data.

**Table 1** Distribution of 3-alkylpyridine derivatives and polyacetylenic compounds within Haplosclerida sponges.


	3-Alkyl-pyridine	3-Alkyl-pyridiniums	Simple C4–C5' bis-3-alkylpiperidine	Complex C4–C5' bis-3-alkylpiperidine	Manzamines	Polyacetylenes
<b>Callyspongiidae</b>						
<i>Arenosclera brasiliensis</i>			X			
<i>Callyspongia fistularis</i>						X
<i>Callyspongia pseudoreticulata</i>						X
<i>Callyspongia ridleyi</i>		X				
<i>Callyspongia siphonella</i>						X
<i>Callyspongia truncata</i>						X
<i>Callyspongia</i> sp.		X				X
<i>Siphonochalina</i> sp.						X
<b>Chalinidae</b>						
<i>Haliclona densaspicula</i>				X		
<i>Haliclona fulva</i>						X
<i>Haliclona osiris</i>						X
<i>Haliclona sarai</i>				X		
<i>Haliclona viscosa</i>		X	X			
<i>Haliclona</i> ( <i>Haliclona</i> )						X
<i>Haliclona</i> sp.	X	X	X		X	X
<i>Haliclona</i> ( <i>Reniera</i> )		X		X		
<b>Niphatidae</b>						
<i>Amphimedon</i> sp.	X	X	X		X	
<i>Cribrachalina dura</i>						X
<i>Cribrachalina vasculum</i>						X
<i>Cribrachalina</i> sp.	X					
<i>Niphates lunisimisis</i>						X
<i>Niphates</i> sp.	X	X				X
<i>Pachychalina alcaloidifera</i>		X	X	X		
<i>Pachychalina</i> sp.		X				
<b>Petrosiidae</b>						
<i>Acanthostrongylophora ingens</i>			X		X	
<i>Acanthostrongylophora</i> sp.					X	
<i>Neopetrosia exigua</i>			X			
<i>Neopetrosia proxima</i>			X			
<i>Neopetrosia seriata</i>			X			
<i>Petrosia corticata</i>						X
<i>Petrosia ficiformis</i>						X
<i>Petrosia solida</i>						X
<i>Petrosia strongylata</i>						X
<i>Petrosia volcano</i>						X
<i>Petrosia</i> ( <i>Strongylophora</i> )						X
<i>Petrosia</i> sp.						X
<i>Xestospongia muta</i>						X
<i>Xestospongia testudinaria</i>						X
<i>Xestospongia</i> sp.					X	X
<b>Phloeodictyidae</b>						
<i>Oceanapia triangulata</i>						X
<i>Oceanapia</i> sp.						X

### Family Callyspongiidae


Alkylation at the nucleophilic nitrogen N-1 of the pyridine leading to 3-alkylpyridinium salts is observed in some species of this family. A tris-3-alkylpyridine derivative called niphatoxin C was found in *Callyspongia* sp. from the Western Pacific (Northwestern Australia), while Buchanan et al. showed the presence of 3-alkylpyridinium polymeric salts from *Callyspongia* (*Toxochalina*) *ridleyi* also collected in the Western Pacific (Papua New Guinea; ● Fig. 4) [31]. These compounds with MW between 5 and 6 kDa cause an irreversible membrane potential depolarization.

In some particular cases, 3-alkylpyridiniums are functionalized through reduction of the aromatic ring leading to highly reactive iminium/enamines intermediates that most commonly lead to the coupling of two units through a C-4/C-5' bond. This function

to connect the two chemical moieties is highly specific and the capability is present only in Haplosclerida and within this group in only a small number of species. The presence of this enzymatic capacity is likely to have phylogenetic significance and thus represents a link between chemistry and taxonomy. Within the family Callyspongiidae, Torres et al. [32] described the arenosclerins A, B, and C but also haliclonyclamine E, macrocyclic bis-3-alkylpiperidine derivatives from the Southwestern Pacific sponge *Arenosclera brasiliensis* (● Fig. 5) [32]. Inversion of configurations at several asymmetric centers is intriguing and suggests a non-specificity of the enzymes involved. Later, the same authors reported some antibacterial activities of arenosclerins but also some cytotoxic activities against tumoral cell lines [33]. *Arenosclera* as a member of the family Callyspongiidae is a genus well




**Fig. 1** 3-Alkylpyridine derivatives isolated from the species *Amphimedon* sp. (Niphatidae)



**Fig. 2** 3-Alkylpyridine derivatives isolated from the species *Cribrochalina* sp. and *Niphates* sp. (Niphatidae).

separated from *Haliclona* (Chalinidae) where bis-3-alkylpiperidines are more commonly found. However, recent molecular studies suggest that members of the Callyspongiidae family are closely related to the type species of *Haliclona* including the only *Arenosclera* species so far sequenced (*Arenosclera heroni*) [8, 22, 34].



**Fig. 3** 3-Alkylpyridinium salts isolated from Niphatidae sponges.



amines A–D were isolated from another *Haliclona* sp. in the Western Pacific (Australia; ● Fig. 5) [58,59]. These compounds were found to be cytotoxic, but also displayed antibacterial and antifungal properties [54].

There are some taxonomic connections between the families Chalinidae and Niphatidae that are currently under investigation, but are relevant to the organization of this review. Very few C-4/C-5' connected bis-3-alkylpiperidine derivatives have been isolated from the Niphatidae family, but include oxidized analogues of halicyclamines identified in *Amphimedon* sp. from the North-western Pacific (Japan; ● Fig. 5) [60] and some bis-3-alkylpiperidine alkaloids identified from *Pachychalina alcaloidifera*, namely haliclonacyclamine F and arenosclerins D and E (● Fig. 5) [61]. However, *H. viscosa* mentioned above is more related via molecular data to species of *Niphates* than it is to the type species of *Haliclona* (Stephens, unpublished PhD thesis, National University of Ireland, Galway). Some of the Australian *Haliclona* also cluster with *Niphates* species rather than the type *Haliclona* species, meaning that this group of compounds may end up being more characteristic of Niphatidae than the Chalinidae in time, and make better sense of the distribution of these specific derivatives.

### Family Petrosiidae

A large diversity of alkaloids has been described from sponges of the genus *Acanthostrongylophora*, but only a small number of simple bis-3-alkylpiperidines. Garson's group isolated a derivative named acanthocyclamine A from *Acanthostrongylophora ingens* collected from the Western Pacific (Indonesia; ● Fig. 7) [62]. Nakagawa et al. [63] first isolated Xestospongins A–D from the Northwestern Pacific (Japan) sponge *Xestospongia exigua* that was subsequently renamed *Neopetrosia exigua* (● Fig. 7). Araguspungines A–H are stereoisomers of these compounds isolated from *Xestospongia* sp. collected in the same area [64]. An additional xestospongine analogue was further isolated from *Xestospongia* sp. collected in the Southwestern Pacific (New Caledonia) [65]. Some macrocyclic bis-3-alkylpiperidine named araguspongines C and D, but also xestospongine D, were also isolated from *N. exigua* collected in the Northeastern Indian Ocean (India; ● Fig. 7), while Araguspungine M was isolated from this species collected in Palau and a dimethyl analogue of xestospongine C was also isolated from this species from the Northwestern Pacific [66–68]. Wei et al. [69] isolated the C-4/C-5' connected neopetrosiamine A from a related species from the Caribbean, i.e., *Neopetrosia proxima* (● Fig. 7). This compound was shown to exhibit cytotoxic bioactivities, and the relatedness of the two *Neopetrosia* species is evident also from molecular data [22]. Later, xestoproxamines were described from the same Caribbean sponge (*N. proxima*) [70]. Petrosins A and B isolated from *Petrosia seriata* collected in the Western Pacific (Indonesia) are unique reports of 3-alkylpiperidine derivatives from this genus (● Fig. 7) [71,72]. This sponge has been renamed as *Neopetrosia seriata* and two separate pieces of DNA evidence (a short piece of the 28 S ribosomal RNA gene and a region of the mitochondrial cytochrome oxidase 1 gene) place this species with another but unnamed *Petrosia* species [8]. The relationships of both species to other *Petrosia* or *Neopetrosia* or indeed any other Haplosclerida currently remain unresolved.




Fig. 6 Polyamines from Chalinidae sponges.




Fig. 7 Simple 3-alkylpiperidines isolated from Petrosiidae sponges.

### Distribution of Complex 3-Alkylpiperidine Derivatives in the Haplosclerida

▼ The complexity of 3-alkylpiperidine derivatives culminates with the biosynthesis of outstanding complex molecules, most of them including the C-4/C-5' but also additional C-C connections and then becoming real cage compounds. Many such compounds are present in members of Chalinidae and Niphatidae families, which, as mentioned above, require reassessment, e.g., some *Haliclona* should be renamed and are more closely related to members of the Niphatidae family. Hence, they are treated here in a separate section. Saraines 1–3 and A–C were isolated from the Northern Mediterranean sponge *Haliclona (Rhizoniera) sarai*

(● Fig. 5) [73,74]. Both piperidine cycles are also connected through the C-4/C-5' bond, but in the case of saraines A–C, other connections were established between both cycles. A large array of biological activities have been described for these compounds [75]. Finally a bis-3-alkylpiperidine derivative called haliclonein A was isolated from *Haliclona* sp. in the Northwestern Pacific (Korea) [76]. Just like saraines A–C, this compound exhibits a C-2/C-3' bond and one of the piperidines has been opened, maybe after oxidation/hydrolysis processes.

An unusual connection with a quinoline aromatic ring on one alkyl chain is characteristic of the njaoamines A–H isolated from the same *Reniera* sp. as the njaoaminiums in the Western Indian Ocean (● Fig. 8) [77,78]. In these compounds, both piperidine cycles are connected through C-4/C-5' as before but also through a second C-3/C-2' connection. Densanins were isolated from *Haliclona densaspicula* in the shallow waters of the Northwestern Pacific (Japan) [79]. They are highly complex pyrrole alkaloids, but their biosynthesis may originate from a common bis-3-alkylpiperidine group through contraction of a piperidine. We still observe the usual C-4/C-5' connection between both cycles. Complex bis-3-alkylpiperidine alkaloids ingenamine G and madangamine F have been isolated from the Southwestern Atlantic sponge *Pachychalina alcaloidifera*, a species currently placed in the family Niphatidae via morphology (● Fig. 8) [28,61,80]. Preliminary molecular data suggest that *H. sarai* is actually more closely related to species of *Niphatidae* than to the *Haliclona* species (*H. oculata*). Much more work relating to morphology, DNA, and chemical evidence is again needed to understand the origin and development of such compounds. Despite the complexity of the chemistry in *P. alcaloidifera*, the authors were not able to isolate more advanced derivatives belonging to the manzamine alkaloids from this sponge.

In some cases, the opening of one piperidine ring can lead to condensation of aromatic rings through Pictet–Spengler condensation. This is the case for the well-known manzamines A and B that maintain the C-4/C-5' connection and which were first isolated from *Haliclona* sp. in the Northwestern Pacific (Japan; ● Fig. 9) [81,82]. Manzamine C, isolated in the same study, lacks the 3-alkylpiperidine parts but keeps the  $\beta$ -carboline. These compounds are among the most promising natural antiparasitic and antitumoral compounds to date [83]. A close analogue called manzamine Y was isolated later from a similar sponge (● Fig. 9) [84]. At the same time, similar compounds named keramamimes A and B were isolated from *Pellina* sp. [85]. However, its taxonomic status has not been clearly confirmed and it may be the same *Haliclona* species mentioned above. The structure of keramamine B has subsequently been revised and corresponds to manzamine F, a structure much more consistent with biosynthetic hypotheses (● Fig. 9) [86].

A large diversity of manzamine alkaloids has also been isolated from the family Niphatidae and especially from one *Amphimedon* species. First, 6-hydroxymanzamine and 3,4-dihydromanzamine were isolated from *Amphimedon* sp. in the Northwestern Pacific (Japan) (● Fig. 9) [87]. Keramaphidin B is a plausible biogenetic precursor of the bis-3-alkylpiperidine parts of manzamine [88], while keramaphidin C and keramamine C [89] are precursors of manzamine C (● Fig. 9), and icanols are antipode of the putative biosynthetic precursor of the manzamines [90]. Tetrahydro- $\beta$ -carbolines manzamines H and L were isolated from the same species of *Amphimedon* (● Fig. 9) [91] and additional manzamines were isolated later: 3,4-dihydromanzamine J, 3,4-dihydro-6-hydroxymanzamine A and manzamine M [92]. Finally, manz-




Fig. 8 Complex bis-3-alkylpiperidines from Chaliniidae sponges.




Fig. 9 Manzamine alkaloids isolated from Chaliniidae and Niphatidae sponges.


amine derivatives condensed with a second  $\beta$ -carboline unit gave a name to the zamamidines, but also 3,4-dihydropyridazine J *N*-oxide and 3,4-dihydro-6-hydroxymanzamine A [93]. Finally a derivative of keramaphidin B named zamamiphidin A was isolated from the same *Amphimedon* sp. (Fig. 9) [94].

Isolated from *Acanthostrongylophora* sp., a sponge currently placed in the family Petrosiidae and collected in the Northwestern Pacific, are quite a number of manzamines, i.e., ircinal A, ircinal A, 12,28-oxaircinal A, manzamine A, manzamine A *N*-oxide, 3,4-dihydropyridazine A *N*-oxide, and 8-hydroxymanzamine A as well as manzamine B, 8-hydroxymanzamine B, manzamine E, 12,34-oxamanzamine E and both 6-hydroxy derivatives, 12,28-oxamanzamine E, manzamine F, manzamine J, 8-hydroxymanzamine J, manzamine X, 6-deoxymanzamine X, and acantholactone (Fig. 10) [95–97]. It is important to notice here that the extraordinary complex dimer named *neo*-kauluamine was isolated from this sponge but had also been isolated previously from a species belonging to a new genus (and as yet undescribed) within the Petrosiidae family [98]. The huge pharmacological potential of these compounds showing antimicrobial and antialzheimer activities has been demonstrated [91]. Other researchers demonstrated the presence of manzamine A and its 8-hydroxy derivative as well as a highly original series of acanthomanzamines A–E and additional manzamine derivatives like acantholactam and pre-*neo*-kauluamine (Fig. 11) from *A. ingens* collected in the same area [99–101]. Two precursors of manzamine alkaloids were isolated from *Xestospongia* sp. collected in the Western Pacific (Papua New Guinea) and the structure of xestocyclamine A was later revised (Fig. 11) [102, 103]. Finally, a large chemical diversity of complex bis-3-alkylpiperidines was isolated from *Xestospongia ingens* (Petrosiidae) collected in the Western Pacific (Papua New Guinea). It is worth noting here that the name of this sponge is no longer accepted and instead refers to *A. ingens* already mentioned previously. Ingenamine A was first isolated [104] and then ingamines A and B [105], madangamine A [106], and other analogues (Fig. 11) [107]. The unique alkaloid 8-hydroxymanzamine A was isolated from *Pachypellina* sp. (Phloedictyidae) but this species was later assigned to the same unnamed genus from the Petrosiidae family mentioned above [98, 108].


### Distribution of Polyacetylenic Derivatives across the Haplosclerida

It has been impossible to find a chemical and logical classification of the polyacetylenic compounds produced by members of the Haplosclerida and the chemical structures do not give valuable information for identification of key enzymes leading to their synthesis. A pattern, however, does emerge linking *either* alkaloid or polyacetylene pathways in each species but never both (Table 1). Therefore, we report on a selection of compounds from this group in view of the taxonomic classification of the species in question as we did for the alkaloids.

A large diversity of polyacetylenes, mostly from the aikupikanyne group, was isolated from *Callyspongia* species, including *Callyspongia truncata*, *Callyspongia pseudoreticulata*, and *Callyspongia fistularis* (Callyspongiidae) [109–115]. Previously classified as *Siphonochalina* (also Callyspongiidae), *C. truncata* and *Callyspongia siphonella* uniquely produce polyacetylenic derivatives [116–118]. The change of genus indicates the difficulty with identifying haplosclerid sponges via morphological methods alone.



**Fig. 10** Manzamine alkaloids from the sponge *Acanthostrongylophora* sp. (Petrosiidae).



**Fig. 11** Manzamine alkaloids from sponges *A. ingens* and *Xestospongia* (Petrosiidae).



From Chalinidae sponges, several renierins followed by fulvinol and the polyhydroxylated fulvynes were isolated from the Mediterranean sponge *Haliclona (Halichocona) fulva* [119–121]. Some polyacetylenes were also isolated from *Haliclona* sp. collected in the Western Pacific (Palau) while polyhydroxylated osirisynes were isolated from *H. (Reniera) osiris* collected in the Northwestern Pacific (China) [122, 123]. Other compounds from unidentified *Haliclona* species include Haliclonyne [124], lembehynes (Northwestern Pacific, Japan) [125, 126], Halycinones A and B [127] (Western Pacific, Micronesia), and brominated derivative [128]. *Adocia* sp. provided adocycetylenes A–D, and while this genus is now accepted as *Haliclona (Haliclona)* [129], it is likely that the genus *Adocia* will be re-erected in the near future (McCormack, unpublished data). How many different species are included in the above list and what their actual final classification will be is therefore as yet unknown and needs attention.

Among the family Niphatidae, the genus *Cribrochalina* has provided a large diversity of polyacetylenic compounds. Duryne was first isolated from *C. dura* collected in the Northwestern Atlantic (Bahamas) [130], while the Caribbean sponge *C. vasculum* afforded several simple acetylenic alcohols, including vasculyne [131–134]. Nepheliosyne B was isolated from *Niphates* sp. collected in the Western Pacific (New Caledonia) [135], and chlorinated acetylenic compounds were isolated from *Haliclona lunisimisis*, later renamed *Niphates lunisimisis* de Jesus & Faulkner [136].

The family Petrosiidae has yielded a large diversity of polyacetylenic products, mostly from sponges of the genus *Petrosia*, a genus that, unfortunately, also needs taxonomic revision. The Mediterranean sponge *Petrosia ficiformis* has afforded a large panel of polyacetylenic derivatives, including petroformynes 1–10 [137–144], while petrosolic acid was later isolated from *Petrosia* sp. from the red sea [145]. Aztequynols A and B were isolated from a Caribbean *Petrosia* sp. [146]. Northwestern Pacific species from this genus have generated many compounds, including Petrosynol and petrosynone isolated from *Petrosia* sp., [147], brominated derivatives from *P. (Petrosia) volcano* [148], corticatic acids A–E from *P. (Strongylophora) corticata* [149, 150], petrocortynes A–H and petrosiacetylenes A–D and other analogues from *Petrosia* sp. collected in Korea [151–157], strongylodiols from *P. (Strongylophora)* sp. [158, 159], polyacetylene carboxylic acids [160], neopetroformynes A–D and later miyakosynes A–F [161, 162], petroacetylene from *P. (Petrosia) solida* [163], and finally, petrosiols A–E from *P. (Strongylophora) strongylata* [164].

Research focused on another petrosid genus led to the isolation of some polyacetylenic compounds from the Caribbean sponge *Xestospongia muta* [165], while several brominated derivatives were isolated from *Xestospongia testudinaria* from the Southwestern Pacific (Australia) [166–168]. Other compounds isolated from *Xestospongia* sp. include unsaturated fatty acids [169], melynones A–C [170], a polyhydroxylated derivative named nepheliosyne A analogous to fulvynes and osirisynes [171], and brominated unsaturated fatty acids [172].

With regard to the family Phloedictyidae, their unique isolated compounds were polyacetylenic derivatives, triangulynes A–H and pelynnols A–D from the Western Pacific sponge (New Caledonia) *Oceanapia triangulate* [173, 174], and another polyacetylene from *Oceanapia* sp. [175]

## Integrating Chemistry and Biology



A large diversity of novel compounds of interest have been isolated from haplosclerid sponges and this group is likely to be a focus of continued efforts in this area for the foreseeable future given the diversity of species and habitats available providing a huge diversity of chemicals in turn. This review provides tantalizing insights into possible patterns of shared structures and biosynthetic pathways between species. However, real insights are hampered by the fact that most species included have not been identified to species and have not been included in any modern phylogenetic study. It is important to convey to chemists about the ever-changing world of taxonomic classification given the new methods being employed, including phylogenetic analysis of DNA sequence data. This is never truer than for haplosclerid sponges. There is an urgent need to revisit the studied sponges to place them in the new phylogenetic classification (including patterns of ancestry and descent between species) being built for demosponges. Only an approach of fully identifying sponges at the species level for analysis of chemistry will provide real additional value to the usual tools of systematics, thus contributing significantly to integrative taxonomy. Marine natural chemists should work tightly with taxonomic specialists (including specifically a phylogenetic approach) of each group in order to publish relevant data. For example, some species groups, i.e., of unidentified *Haliclona* and *Xestospongia* species, produce alkaloids but also polyacetylenic compounds, which prevents us from reaching any clear conclusion regarding patterns of distribution of specific compounds across taxonomic groups. However, once understood, the presence of each family of compounds could help significantly in a proper classification and avoid misinterpretation of species identification/classification.

From **Table 1**, several clues are provided for assisting sponge classification. First of all, the presence of both polyacetylenic and alkaloid compounds have never been reported in a single species. Both biosynthetic genes/pathways cannot, therefore, be present in a species. This observation leads us to suggest a clear separation between species producing alkaloids and species producing polyacetylenes. For example, *C. ridleyi* is the only known species of the genus *Callyspongia* that does not produce polyacetylenic compounds. In the same way, only one species of *Cribrochalina* sp. was found to produce alkaloids. This could lead to a clear revision of the taxonomy of these sponges. Finally, *Neopetrosia* species were found to produce alkaloids, while *Petrosia* species produce only polyacetylenic compounds. However, these two genera are currently placed in the same family, Petrosiidae, indicating that revision is required at different levels of the taxonomic hierarchy.

One undefined species of *Cribrochalina* (Niphatidae) produces only 3-alkylpyridine derivatives. Simple alkylation at the nitrogen atom of the pyridine leads to 3-alkylpyridiniums that can polymerize. Only the species of *Callyspongia* (Callyspongiidae), *Niphates*, and *Pachychalina* (Niphatidae) seem to stop the biosynthesis at this step for this group of alkaloids. Because these genera are distinct, revisiting the voucher samples could help to identify similarities between these species. The subsequent ability of some species to reduce the aromatic rings enabling a C-4/C-5' connection between two cycles is shared by several species, including *Arenosclera brasiliensis* (Callyspongiidae) and *H. viscosa* (Chalinidae). Contrary to other families, members of Callyspongiidae seem unable to develop more complex alkaloids. In the same way, *Neopetrosia* (Petrosiidae) species only produce simple

bis-3-alkylpiperidine derivatives. *Neopetrosia* could be hypothesized then to be closer to some *Callyspongia* sponges, and recent phylogenetic classification suggests that some *Neopetrosia* are placed within the same major clade as *Callyspongia* (Clade A) [8, 21]. Much more complex alkaloids are produced by few species distributed in several families, which also raises questions regarding the reliability of the classification. *H. densaspicula*, *H. sarai*, a *Reniera* sp. (Chalinidae), and *Pachychalina alcaloidifera* (Niphatidae) are able to produce complex alkaloids, but no manzamines have been isolated from these species. It appears that clearly this ability should gather some *Haliclona* species. As for the extremely complex alkaloid manzamines, members of the genera *Acanthostrongylophora* (Petrosiidae), *Haliclona* (Chalinidae), and *Amphimedon* (Niphatidae) were found to produce them. These compounds are not characteristic of the Petrosiidae, as most species of this family produce only polyacetylenes.

A diversity of polyacetylenic compounds is found distributed across all of the five families of Haplosclerida included so far and their patterns of occurrence should trigger careful examination of the studied species. For example, four *Callyspongia* species (Callyspongiidae) produce compounds very similar to two *Haliclona* species (Chalinidae), two *Cribochalina* species, and one *Niphates* species (Niphatidae), but most of the polyacetylenic compounds are really representative of *Petrosia* and *Xestospongia* genera (Petrosiidae). Recent molecular phylogenies suggest that *Callyspongia* and some *Haliclona* fall into one major clade (Clade A) at the base of which appears *Cribochalina* [8, 22]. *Niphates* species fall into a distantly related clade, also with some *Haliclona* (Clade C), while species from *Petrosia* and *Xestospongia* are distributed across multiple clades (but distinct from the aforementioned genera) and are poorly represented on phylogenetic trees drawn from molecular data as yet [8, 22].

The value of a chemosystematics approach for haplosclerid sponges has been questioned in the past due to disagreement between patterns of chemical diversity and morphological classification [176]. With molecular data throwing light on possible evolutionary pathways in sponges, it is evident that the morphological classification is flawed and is likely to change, and this is especially true for the Haplosclerida. An additional concern could be the varying influence of microbes residing in sponge tissue, which may be responsible for some of the compounds isolated. A diversity of microbial sequences has been reported from some *Haliclona* species and *Xestospongia*, amongst others [177–180], even though these studies do not show the presence of microbes in the sponge tissue nor do they confirm the source of compounds of interest as being from sponge or microbial cells. Sponges filter feed and concentrate microbes and microbial constituents from their environment, and these elements will be present in approaches that only isolate DNA sequences from sponge cells (even though they may only represent food for the sponge). Therefore, the presence of many bacterial DNA sequences in sponges does not equate directly to a sponge-bacterial association or to the production of detectable levels of compounds of interest by said bacteria. Ultrastructural studies on Irish and Mediterranean *Haliclona* species (Marra, unpublished data) show extremely low numbers of bacterial cells in four species, while an association between one or very few bacteria is evident in three species and many bacterial species are evident in one. Consideration of such patterns in these and other species will also be required to truly understand how patterns of microbial diversity impact patterns of chemical diversity.

## Conclusion

Our main conclusion is that chemical diversity does not fit with the current classification of this major group of marine sponges, but supports a clear revision of all the species of the Order Haplosclerida included in chemical publications taking into account the presence or absence of key enzymes leading to 3-alkylpyridine derivatives or polyacetylenes. Despite apparent discrepancies there are tantalizing insights that patterns of chemical diversity may well agree in good part with an updated classification that incorporates molecular and other data including morphology. Assessment of the possible bacterial origin of sponge-derived compounds should also be considered. Three other limitations to this approach are the existence of mistakes made in the identification of the species under investigation, a discrepancy between a classical taxonomic approach utilizing shared sponge morphological characters and the approach that uses molecular phylogenetic methods, and, thirdly, poor sampling of the huge diversity of sponges and the chemistry they contain. However, we believe that this review should help in the further development of an integrative approach between biology and chemistry for this group of sponges. The issues highlighted here are not restricted to the group of marine sponges included, but are relevant to all organisms that are of interest in the search for novel chemistry for whatever reason and where taxonomy is still unresolved. New advances in 'omics' sciences, and especially metabolomic approaches using mass spectrometry and nuclear magnetic resonance with some characteristic signals, could quickly give important clues for integrative systematics.

## Conflict of Interest

The authors declare no conflict of interest.

## References

- Zhang F, Blasiak LC, Karolin JO, Powell RJ, Geddes CD, Hill RT. Phosphorus sequestration in the form of polyphosphate by microbial symbionts in marine sponges. *Proc Natl Acad Sci USA* 2015; 112: 4381–4386
- De Goeij JM, van Oevelen D, Vermeij MJA, Osinga R, Middelburg JJ, de Goeij AF, Admiraal W. Surviving in a marine desert: the sponge loop retains resources within coral reefs. *Science* 2013; 342: 108–110
- Genta-Jouve G, Thomas OP. Sponge chemical diversity: from biosynthetic pathways to ecological roles. *Adv Mar Biol* 2012; 62: 183–230
- Cardenas P, Perez T, Boury-Esnault N. Sponge systematics facing new challenges. *Adv Mar Biol* 2012; 61: 79–209
- Van Soest RM, Hooper JA. Order Haplosclerida Topsent, 1928. In: Hooper JA, Van Soest RM, Willenz P, editors. *Systema Porifera – A guide to the classification of sponges*. New York: Springer US; 2002: 831–832
- Fromont J, Kerr S, Kerr R, Riddle M, Murphy P. Chemotaxonomic relationships within, and comparisons between, the orders Haplosclerida and Petrosida (Porifera: Demospongiae) using sterol complements. *Biochem Syst Ecol* 1994; 22: 735–752
- Andersen RJ, van Soest RWM, Kong F. 3-Alkylpiperidine alkaloids isolated from marine sponges in the order Haplosclerida. In: Pelletier SW, editor. *Alkaloids: chemical and biological perspectives*. Oxford: Pergamon Press; 1996: 301–355
- Redmond NE, Raleigh J, van Soest RWM, Kelly M, Travers SAA, Bradshaw B, Vartia S, Stephens KM, McCormack GP. Phylogenetic relationships of the marine Haplosclerida (Phylum Porifera) employing ribosomal (28S rRNA) and mitochondrial *cox1*, *nad1* gene sequence data. *PLoS One* 2011; 6: e24344
- Van Soest RWM, Boury-Esnault N, Hooper JNA, Rützler K, de Voogd NJ, Alvarez de Glasby B, Hajdu E, Pisera AB, Manconi R, Schoenberg C, Janussen D, Tabachnick KR, Klautau M, Picton B, Kelly M, Vacelet J, Dohrmann M, Díaz MC, Cárdenas P. World Porifera database. Available

- at: <http://www.marinespecies.org/porifera>. Accessed September 1, 2015
- 10 Tsukamoto S, Takahashi M, Matsunaga S, Fusetani N, Van Soest RW. Hachijodines A–G: seven new cytotoxic 3-alkylpyridine alkaloids from two marine sponges of the genera *Xestospongia* and *Amphimedon*. *J Nat Prod* 2000; 63: 682–684
  - 11 Takekawa Y, Matsunaga S, van Soest RW, Fusetani N. Amphimedosides, 3-alkylpyridine glycosides from a marine sponge *Amphimedon* sp. *J Nat Prod* 2006; 69: 1503–1505
  - 12 Hirano K, Kubota T, Tsuda M, Mikami Y, Kobayashi J. Pyrinodemins B–D, potent cytotoxic bis-pyridine alkaloids from marine sponge *Amphimedon* sp. *Chem Pharm Bull (Tokyo)* 2000; 48: 974–977
  - 13 Tsuda M, Hirano K, Kubota T, Kobayashi J. Pyrinodemin A, a cytotoxic pyridine alkaloid with an isoxazolidine moiety from sponge *Amphimedon* sp. *Tetrahedron Lett* 1999; 40: 4819–4820
  - 14 Kubota T, Kura K, Fromont J, Kobayashi J. Pyrinodemins G–I, new bis-3-alkylpyridine alkaloids from a marine sponge *Amphimedon* sp. *Tetrahedron* 2013; 69: 96–100
  - 15 Kura K, Kubota T, Fromont J, Kobayashi J. Pyrinodemins E and F, new 3-alkylpyridine alkaloids from sponge *Amphimedon* sp. *Bioorg Med Chem Lett* 2011; 21: 267–270
  - 16 Nishi T, Kubota T, Fromont J, Sasaki T, Kobayashi J. Nakinadines B–F: new pyridine alkaloids with a  $\beta$ -amino acid moiety from sponge *Amphimedon* sp. *Tetrahedron* 2008; 64: 3127–3132
  - 17 Quiñoà E, Crews P. Niphatynes, methoxylamine pyridines from the marine sponge, *Niphates* sp. *Tetrahedron Lett* 1987; 28: 2467–2468
  - 18 Kobayashi J, Zeng CM, Ishibashi M, Shigemori H, Sasaki T, Mikami Y. Niphatesines E–H, new pyridine alkaloids from the Okinawan marine sponge *Niphates* sp. *J Chem Soc Perkin Trans 1* 1992; 11: 1291–1294
  - 19 Kobayashi J, Murayama T, Kosuge S, Kanda F, Ishibashi M, Kobayashi H, Ohizumi Y, Ohta T, Nozoe S, Sasaki T. Niphatesines A–D, new antineoplastic pyridine alkaloids from the Okinawan marine sponge *Niphates* sp. *J Chem Soc Perkin Trans 1* 1990; 12: 3301–3303
  - 20 Kariya Y, Kubota T, Fromont J, Kobayashi J. Pyrinadines B–G, new bispyridine alkaloids with an azoxy moiety from sponge *Cribrochalina* sp. *Bioorg Med Chem* 2006; 14: 8415–8419
  - 21 Kariya Y, Kubota T, Fromont J, Kobayashi J. Pyrinadine A, a novel pyridine alkaloid with an azoxy moiety from sponge *Cribrochalina* sp. *Tetrahedron Lett* 2006; 47: 997–998
  - 22 Redmond NE, Morrow CC, Thacker RW, Diaz MC, Boury-Esnault N, Cárdenas P, Hajdu E, Lôbo-Hajdu G, Picton BE, Pomponi SA, Kayal E, Collins AG. Phylogeny and systematics of Demospongiae in light of new small-subunit ribosomal DNA (18 S) sequences. *Integr Comp Biol* 2013; 53: 388–415
  - 23 Xu NJ, Sun X, Yan XJ. A new cyclostelletamine from sponge *Amphimedon compressa*. *Chin Chem Lett* 2007; 18: 947–950
  - 24 Schmitz FJ, Hollenbeak KH, Campbell DC. Marine natural products: halitoxin, toxic complex of several marine sponges of the genus *Haliclona*. *J Org Chem* 1978; 43: 3916–3922
  - 25 Thompson MN, Gallimore WA, Townsend MM, Chambers NA, Williams LA. Bioactivity of amphitoxin, the major constituent of the Jamaican sponge *Amphimedon compressa*. *Chem Biodivers* 2010; 7: 1904–1910
  - 26 Albrizio S, Ciminiello P, Fattorusso E, Magno S, Pawlik JR. Amphitoxin, a new high molecular weight antifeedant pyridinium salt from the caribbean sponge *Amphimedon compressa*. *J Nat Prod* 1995; 58: 647–652
  - 27 Talpir R, Rudi A, Ilan M, Kashman Y. Niphatoxin A and B; two new ichthyo- and cytotoxic tripyridine alkaloids from a marine sponge. *Tetrahedron Lett* 1992; 33: 3033–3034
  - 28 De Oliveira JH, Grube A, Köck M, Berlinck RG, Macedo ML, Ferreira AG, Hajdu E. Ingenamine G and cyclostelletamines GI, K, and L from the new Brazilian species of marine sponge *Pachychalina* sp. *J Nat Prod* 2004; 67: 1685–1689
  - 29 De Oliveira JH, Selegim MH, Timm C, Grube A, Köck M, Nascimento GG, Martins ACT, Silva EG, de Souza AO, Minarini PR. Antimicrobial and antimycobacterial activity of cyclostelletamine alkaloids from sponge *Pachychalina* sp. *Mar Drugs* 2006; 4: 1–8
  - 30 Laville R, Thomas OP, Berrue F, Reyes F, Amade P. Pachychalines A–C: novel 3-alkylpyridinium salts from the marine sponge *Pachychalina* sp. *European J Org Chem* 2008; 2008: 121–125
  - 31 Buchanan MS, Carroll AR, Addepalli R, Avery VM, Hooper JNA, Quinn RJ. Niphatoxin C, a cytotoxic tripyridine alkaloid from *Callyspongia* sp. *J Nat Prod* 2007; 70: 2040–2041
  - 32 Torres YR, Berlinck RG, Magalhães A, Schefer AB, Ferreira AG, Hajdu E, Muricy G. Arenosclerins A–C and haliclonacyclamine E, new tetracyclic alkaloids from a Brazilian endemic Haplosclerid sponge *Arenosclera brasiliensis*. *J Nat Prod* 2000; 63: 1098–1105
  - 33 Torres YR, Berlinck RG, Nascimento GG, Fortier SC, Pessoa C, de Moraes MO. Antibacterial activity against resistant bacteria and cytotoxicity of four alkaloid toxins isolated from the marine sponge *Arenosclera brasiliensis*. *Toxicol* 2002; 40: 885–891
  - 34 Thacker RW, Hill AL, Hill MS, Redmond NE, Collins AG, Morrow CC, Spicer L, Carmack CA, Zappe ME, Pohlmann D, Hall C, Diaz MC, Bangalore PV. Nearly complete 28 S rRNA gene sequences confirm new hypotheses of sponge evolution. *Integr Comp Biol* 2013; 53: 373–387
  - 35 Schmidt G, Timm C, Koeck M. Haliclocyclin C, a new monomeric 3-alkylpyridinium alkaloid from the arctic marine sponge *Haliclona viscosa*. *Z Naturforsch B* 2011; 66: 745–748
  - 36 Timm C, Volk C, Sasse F, Köck M. The first cyclic monomeric 3-alkylpyridinium alkaloid from natural sources: identification, synthesis, and biological activity. *Org Biomol Chem* 2008; 6: 4036–4040
  - 37 Schmidt G, Timm C, Grube A, Volk CA, Köck M. Viscosamines B(1,2) and E(1,2): challenging new 3-alkylpyridinium alkaloids from the marine sponge *Haliclona viscosa*. *Chemistry* 2012; 18: 8180–8189
  - 38 Volk CA, Köck M. Viscosamine: the first naturally occurring trimeric 3-alkylpyridinium alkaloid. *Org Lett* 2003; 5: 3567–3569
  - 39 Teruya T, Kobayashi K, Suenaga K, Kigoshi H. Cyclohaliclonamines A–E: dimeric, trimeric, tetrameric, pentameric, and hexameric 3-alkylpyridinium alkaloids from a marine sponge *Haliclona* sp. *J Nat Prod* 2006; 69: 135–137
  - 40 Casapullo A, Pinto OC, Marzocco S, Autore G, Riccio R. 3-Alkylpyridinium alkaloids from the pacific sponge *Haliclona* sp. *J Nat Prod* 2009; 72: 301–303
  - 41 Lee Y, Jang KH, Jeon JE, Yang WY, Sim CJ, Oh KB, Shin J. Cyclic Bis-1, 3-dialkylpyridiniums from the sponge *Haliclona* sp. *Mar Drugs* 2012; 10: 2126–2137
  - 42 Damodaran V, Ryan JL, Keyzers RA. Cyclic 3-alkylpyridinium alkaloid monomers from a New Zealand *Haliclona* sp. marine sponge. *J Nat Prod* 2013; 76: 1997–2001
  - 43 Laville R, Genta-Jouve G, Urda C, Fernandez R, Thomas OP, Reyes F, Amade P. Njaoaminiums A, B, and C: cyclic 3-alkylpyridinium salts from the marine sponge *Reniera* sp. *Molecules* 2009; 14: 4716–4724
  - 44 Laville R, Amade P, Thomas OP. 3-Alkylpyridinium salts from Haplosclerida marine sponges: isolation, structure elucidations, and biosynthetic considerations. *Pure Appl Chem* 2009; 81: 1033–1040
  - 45 Baker BJ, Scheuer PJ, Shooley JN. Papuamine, an antifungal pentacyclic alkaloid from a marine sponge, *Haliclona* sp. *J Am Chem Soc* 1988; 110: 965–966
  - 46 Fahy E, Molinski TF, Harper MK, Sullivan BW, Faulkner DJ, Parkanyi L, Clardy J. Haliclonadiamine, an antimicrobial alkaloid from the sponge *Haliclona* sp. *Tetrahedron Lett* 1988; 29: 3427–3428
  - 47 Kashman Y, Koren-Goldshlager G, Gravalos MDG, Schleyer M. Halitulins, a new cytotoxic alkaloid from the marine sponge *Haliclona tulearensis*. *Tetrahedron Lett* 1999; 40: 997–1000
  - 48 Sorek H, Rudi A, Akinin M, Gaydou EM, Kashman Y. Isohalitulins and halicloreinsins B and C, three marine alkaloids from *Haliclona tulearensis*. *J Nat Prod* 2010; 73: 456–458
  - 49 Heinrich MR, Kashman Y, Spittler P, Steglich W. Revision of the structure of halicloreinsin to (S)-7-methyl-1,5-diazacyclotetradecane and confirmation of the new structure by synthesis. *Tetrahedron* 2001; 57: 9973–9978
  - 50 Koren-Goldshlager G, Kashman Y, Schleyer M. Halicloreinsin, a novel diamino alkaloid from the marine sponge *Haliclona tulearensis*. *J Nat Prod* 1998; 61: 282–284
  - 51 Fusetani N, Yasumuro K, Matsunaga S, Hirota H. Haliclamines A and B, cytotoxic macrocyclic alkaloids from a sponge of the genus *Haliclona*. *Tetrahedron Lett* 1989; 30: 6891–6894
  - 52 Schmidt G, Timm C, Kock M. New haliclamines E and F from the Arctic sponge *Haliclona viscosa*. *Org Biomol Chem* 2009; 7: 3061–3064
  - 53 Volk CA, Lippert H, Lichte E, Köck M. Two new haliclamines from the Arctic sponge *Haliclona viscosa*. *Eur J Org Chem* 2004; 2004: 3154–3158
  - 54 Cychon C, Schmidt G, Mordhorst T, Koeck M. Structure elucidation of submilligram quantities of natural products – application to haliclamines G and H from the Arctic marine sponge *Haliclona viscosa*. *Z Naturforsch B* 2012; 67: 944–950

- 55 Jaspars M, Pasupathy V, Crews P. A tetracyclic diamine alkaloid, halicyclamine A, from the marine sponge *Haliclona* sp. *J Org Chem* 1994; 59: 3253–3255
- 56 Arai M, Liu L, Fujimoto T, Setiawan A, Kobayashi M. DedA protein relates to action-mechanism of halicyclamine A, a marine spongean macrocyclic alkaloid, as an anti-dormant mycobacterial substance. *Mar Drugs* 2011; 9: 984–993
- 57 Arai M, Sobou M, Vilchéze C, Baughn A, Hashizume H, Pruksakorn P, Ishida S, Matsumoto M, Jacobs jr. WR, Kobayashi M. Halicyclamine A, a marine spongean alkaloid as a lead for anti-tuberculosis agent. *Bioorg Med Chem* 2008; 16: 6732–6736
- 58 Charan RD, Garson MJ, Brereton IM, Willis AC, Hooper JNA. Haliclona-cyclamines A and B, cytotoxic alkaloids from the tropical marine sponge *Haliclona* sp. *Tetrahedron* 1996; 52: 9111–9120
- 59 Clark RJ, Field KL, Charan RD, Garson MJ, Brereton M, Willis AC. The haliclona-cyclamines, cytotoxic tertiary alkaloids from the tropical marine sponge *Haliclona* sp. *Tetrahedron* 1998; 54: 8811–8826
- 60 Matsunaga S, Miyata Y, van Soest RW, Fusetani N. Tetradehydrohalicyclamine A and 22-Hydroxyhalicyclamine A, new cytotoxic bis-piperidine alkaloids from a marine sponge *Amphimedon* sp. *J Nat Prod* 2004; 67: 1758–1760
- 61 De Oliveira JH, Nascimento AM, Kossuga MH, Cavalcanti BC, Pessoa CO, Moraes MO, Macedo ML, Ferreira AG, Hajdu E, Pinheiro US. Cytotoxic alkyloxy-piperidine alkaloids from the Brazilian marine sponge *Pachychalina alcaloidifera*. *J Nat Prod* 2007; 70: 538–543
- 62 Dewi AS, Hadi TA, Fajarningsih ND, Blanchfield JT, Bernhardt PV, Garson MJ. Acanthocyclamine A from the Indonesian marine sponge *Acanthostrongylophora ingens*. *Aust J Chem* 2014; 67: 1205–1210
- 63 Nakagawa M, Endo M, Tanaka N, Gen-Pei L. Structures of xestospongins A, B, C and D, novel vasodilative compounds from marine sponge, *Xestospongia exigua*. *Tetrahedron Lett* 1984; 25: 3227–3230
- 64 Kobayashi M, Kawazoe K, Kitagawa I. Araguspungines B, C, D, E, F, G, H, and J, new vasodilative bis-1-oxaquinolizidine alkaloids from an Okinawan marine sponge, *Xestospongia* sp. *Chem Pharm Bull (Tokyo)* 1989; 37: 1676–1678
- 65 Quirion JC, Sevenet T, Husson HP, Weniger B, Debitus C. Two new alkaloids from *Xestospongia* sp., a New Caledonian sponge. *J Nat Prod* 1992; 55: 1505–1508
- 66 Venkateswarlu Y, Reddy MVR, Rao JV. Bis-1-oxaquinolizidines from the sponge *Haliclona exigua*. *J Nat Prod* 1994; 57: 1283–1285
- 67 Liu H, Mishima Y, Fujiwara T, Nagai H, Kitazawa A, Mine Y, Kobayashi H, Yao X, Yamada J, Oda T. Isolation of araguspungine M, a new stereoisomer of an araguspungine/xestospongine alkaloid, and dopamine from the marine sponge *Neopetrosia exigua* collected in Palau. *Mar Drugs* 2004; 2: 154–163
- 68 Li Y, Qin S, Guo YW, Gu YC, van Soest RW. 9'-Epi-3 $\beta$ , 3' $\beta$ -dimethylxestospongins C, a new macrocyclic diamine alkaloid from the Hainan sponge *Neopetrosia exigua*. *Planta Med* 2011; 77: 179–181
- 69 Wei X, Nieves K, Rodriguez AD. Neopetrosiamine A, biologically active bis-piperidine alkaloid from the Caribbean sea sponge *Neopetrosia proxima*. *Bioorg Med Chem Lett* 2010; 20: 5905–5908
- 70 Morinaka BI, Molinski TF. Xestoproxamines A–C from *Neopetrosia proxima*. Assignment of absolute stereostructure of bis-piperidine alkaloids by integrated degradation-CD analysis. *J Nat Prod* 2011; 74: 430–440
- 71 Braekman JC, Daloz D, Cimino G, Trivellone E. 2D-NMR study of petrosins: Revised structure for petrosin-A. *Bull Soc Chim Belg* 1988; 97: 519–524
- 72 Braekman JC, Daloz D, Defay N, Zimmermann D. Petrosin-A and -B, two new bis-quinolizidine alkaloids from the sponge *Petrosia seriata*. *Bull Soc Chim Belg* 1984; 93: 941–944
- 73 Defant A, Mancini I, Raspor L, Guella G, Turk T, Sepčić K. New structural insights into saraines A, B, and C, macrocyclic alkaloids from the mediterranean sponge *Reniera (Haliclona) sarai*. *Eur J Org Chem* 2011; 2011: 3761–3767
- 74 Cimino G, Stefano SD, Scognamiglio G, Sodano G, Trivellone E. Sarains: A new class of alkaloids from the marine sponge *Reniera sarai*. *Bull Soc Chim Belg* 1986; 95: 783–800
- 75 Caprioli V, Cimino G, de Giulio A, Madaio A, Scognamiglio G, Trivellone E. Selected biological activities of saraines. *Comp Biochem Physiol B* 1992; 103: 293–296
- 76 Jang KH, Kang GW, Jeon JE, Lim C, Lee HS, Sim CJ, Oh KB, Shin J. Haliclolinin A, a new macrocyclic diamide from the sponge *Haliclona* sp. *Org Lett* 2009; 11: 1713–1716
- 77 Sorek H, Rudi A, Benayahu Y, Kashman Y. Njaoamines G and H, two new cytotoxic polycyclic alkaloids and a tetrahydroquinolone from the marine sponge *Neopetrosia* sp. *Tetrahedron Lett* 2007; 48: 7691–7694
- 78 Reyes F, Fernández R, Urda C, Francesch A, Bueno S, de Eguilior C, Cuevas C. Njaoamines A–F, new cytotoxic polycyclic alkaloids from the haplosclerid sponge *Reniera* sp. *Tetrahedron* 2007; 63: 2432–2438
- 79 Hwang BS, Oh JS, Jeong EJ, Sim CJ, Rho JR. Densanins A and B, new macrocyclic pyrrole alkaloids isolated from the marine sponge *Haliclona densaspicula*. *Org Lett* 2012; 14: 6154–6157
- 80 Cavalcanti BC, Sombra CML, de Oliveira JH, Berlink RG, de Moraes MO, Pessoa C. Cytotoxicity and genotoxicity of ingenamine G isolated from the Brazilian marine sponge *Pachychalina alcaloidifera*. *Comp Biochem Physiol C Toxicol Pharmacol* 2008; 147: 409–415
- 81 Sakai R, Kohmoto S, Higa T, Jefford CW, Bernardinelli G. Manzamine B and C, two novel alkaloids from the sponge *Haliclona* sp. *Tetrahedron Lett* 1987; 28: 5493–5496
- 82 Sakai R, Higa T, Jefford CW, Bernardinelli G. Manzamine A, a novel antitumor alkaloid from a sponge. *J Am Chem Soc* 1986; 108: 6404–6405
- 83 Ashok P, Ganguly S, Murugesan S. Manzamine alkaloids: isolation, cytotoxicity, antimalarial activity and SAR studies. *Drug Discov Today* 2014; 19: 1781–1791
- 84 Kobayashi M, Chen YJ, Aoki S, In Y, Ishida T, Kitagawa I. Four new  $\beta$ -carboline alkaloids isolated from two Okinawan marine sponges of *Xestospongia* sp. and *Haliclona* sp. *Tetrahedron* 1995; 51: 3727–3736
- 85 Nakamura H, Deng S, Kobayashi J, Ohizumi Y, Tomotake Y, Matsuzaki T, Hirata Y. Keramamine-A and -B, novel antimicrobial alkaloids from the Okinawan marine sponge *Pellina* sp. *Tetrahedron Lett* 1987; 28: 621–624
- 86 Ichiba T, Sakai R, Kohmoto S, Saucy G, Higa T. New manzamine alkaloids from a sponge of the genus *Xestospongia*. *Tetrahedron Lett* 1988; 29: 3083–3086
- 87 Kobayashi J, Tsuda M, Kawasaki N, Sasaki T, Mikami Y. 6-Hydroxymanzamine A and 3, 4-dihydrmanzamine A, new alkaloids from the Okinawan marine sponge *Amphimedon* sp. *J Nat Prod* 1994; 57: 1737–1740
- 88 Kobayashi J, Tsuda M, Kawasaki N, Matsumoto K, Adachi T. Keramaphidin B, a novel pentacyclic alkaloid from a marine sponge *Amphimedon* sp.: A plausible biogenetic precursor of manzamine alkaloids. *Tetrahedron Lett* 1994; 35: 4383–4386
- 89 Tsuda M, Kawasaki N, Kobayashi J. Keramaphidin C and keramamine C plausible biogenetic precursors of manzamine C from an Okinawan marine sponge. *Tetrahedron Lett* 1994; 35: 4387–4388
- 90 Tsuda M, Kawasaki N, Kobayashi J. Ircinols A and B, first antipodes of manzamine-related alkaloids from an Okinawan marine sponge. *Tetrahedron* 1994; 50: 7957–7960
- 91 Tsuda M, Inaba K, Kawasaki N, Honma K, Kobayashi J. Chiral resolution of ( $\pm$ )-keramaphidin B and isolation of manzamine L, a new  $\beta$ -carboline alkaloid from a sponge *Amphimedon* sp. *Tetrahedron* 1996; 52: 2319–2324
- 92 Watanabe D, Tsuda M, Kobayashi J. Three new manzamine congeners from *Amphimedon* sponge. *J Nat Prod* 1998; 61: 689–692
- 93 Takahashi Y, Kubota T, Fromont J, Kobayashi J. Zamamidines A and B, new manzamine alkaloids from the sponge *Amphimedon* species. *Org Lett* 2008; 11: 21–24
- 94 Kubota T, Kamijyo Y, Takahashi-Nakaguchi A, Fromont J, Gonoï T, Kobayashi J. Zamamiphidin A, a new manzamine related alkaloid from an Okinawan marine sponge *Amphimedon* sp. *Org Lett* 2013; 15: 610–612
- 95 Rao KV, Kananah N, Wahyuono S, Tekwani BL, Schinazi RF, Hamann MT. Three new manzamine alkaloids from a common Indonesian sponge and their activity against infectious and tropical parasitic diseases. *J Nat Prod* 2004; 67: 1314–1318
- 96 Rao KV, Donia MS, Peng J, Garcia-Palmero E, Alonso D, Martinez A, Medina M, Franzblau SG, Tekwani BL, Khan SI. Manzamine B and E and ircinal A related alkaloids from an Indonesian *Acanthostrongylophora* sponge and their activity against infectious, tropical parasitic, and Alzheimer's diseases. *J Nat Prod* 2006; 69: 1034–1040
- 97 Wahba AE, Fromentin Y, Zou Y, Hamann MT. Acantholactone, a new manzamine related alkaloid with an unprecedented  $\delta$ -lactone and  $\epsilon$ -lactam ring system. *Tetrahedron Lett* 2012; 53: 6329–6331
- 98 El Sayed KA, Kelly M, Kara UAK, Ang KKH, Katsuyama I, Dunbar DC, Khan AA, Hamann MT. New manzamine alkaloids with potent activity against infectious diseases. *J Am Chem Soc* 2001; 123: 1804–1808

- 99 Samoylenko V, Khan SI, Jacob MR, Tekwani BL, Walker LA, Hufford CD, Muhammad I. Bioactive (+)-manzamine A and (+)-8-hydroxymanzamine A tertiary bases and salts from *Acanthostrongylophora ingens* and their preparations. *Nat Prod Commun* 2009; 4: 185–192
- 100 Furusato A, Kato H, Nehira T, Eguchi K, Kawabata T, Fujiwara Y, Losung F, Mangindaan RE, de Voogd NJ, Takeya M. Acanthoanzamines A–E with new manzamine frameworks from the marine sponge *Acanthostrongylophora ingens*. *Org Lett* 2014; 16: 3888–3891
- 101 El-Desoky AH, Kato H, Eguchi K, Kawabata T, Fujiwara Y, Losung F, Mangindaan RE, de Voogd NJ, Takeya M, Yokosawa H. Acantholactam and pre-neo-kaulamine, manzamine-related alkaloids from the Indonesian marine sponge *Acanthostrongylophora ingens*. *J Nat Prod* 2014; 77: 1536–1540
- 102 Rodríguez J, Crews P. Revised structure of xestocyclamine A and description of a new analogue. *Tetrahedron Lett* 1994; 35: 4719–4722
- 103 Rodríguez J, Peters BM, Kurz L, Schatzman RC, McCarley D, Lou L, Crews P. An alkaloid protein kinase C inhibitor, xestocyclamine A, from the marine sponge *Xestospongia* sp. *J Am Chem Soc* 1993; 115: 10436–10437
- 104 Kong F, Andersen RJ, Allen TM. Ingenamine, a novel pentacyclic alkaloid from the marine sponge *Xestospongia ingens*. *Tetrahedron Lett* 1994; 35: 1643–1646
- 105 Kong F, Andersen RJ, Allen TM. Ingamines A and B, new cytotoxic alkaloids from the marine sponge *Xestospongia ingens*. *Tetrahedron* 1994; 50: 6137–6144
- 106 Kong F, Andersen RJ, Allen TM. Madangamine A, a novel cytotoxic alkaloid from the marine sponge *Xestospongia ingens*. *J Am Chem Soc* 1994; 116: 6007–6008
- 107 Kong F, Andersen RJ. Ingenamine alkaloids isolated from the sponge *Xestospongia ingens*: Structures and absolute configurations. *Tetrahedron* 1995; 51: 2895–2906
- 108 Ichiba T, Corgiat JM, Scheuer PJ, Kelly-Borges M. 8-Hydroxymanzamine A, a  $\beta$ -carboline alkaloid from a sponge, *Pachypellina* sp. *J Nat Prod* 1994; 57: 168–170
- 109 Umeyama A, Nagano C, Arihara S. Three novel C21 polyacetylenes from the marine sponge *Callyspongia* sp. *J Nat Prod* 1997; 60: 131–133
- 110 Youssef DTA, Yoshida WY, Kelly M, Scheuer PJ. Polyacetylenes from a Red Sea sponge *Callyspongia* species. *J Nat Prod* 2000; 63: 1406–1410
- 111 Nakao Y, Uehara T, Matunaga S, Fusetani N, van Soest RWM. Callyspongynic acid, a polyacetylenic acid which inhibits  $\alpha$ -glucosidase, from the marine sponge *Callyspongia truncata*. *J Nat Prod* 2002; 65: 922–924
- 112 Braekman JC, Daloz D, Devijver C, Dubut D, van Soest RWM. A new C-20 polyacetylene from the sponge *Callyspongia pseudoreticulata*. *J Nat Prod* 2003; 66: 871–872
- 113 Youssef DTA, van Soest RWM, Fusetani N. Callyspongamide A, a new cytotoxic polyacetylenic amide from the Red Sea sponge *Callyspongia fistularis*. *J Nat Prod* 2003; 66: 861–862
- 114 Youssef DT, van Soest RWM, Fusetani N. Callyspongensols A–C, new cytotoxic C22-polyacetylenic alcohols from a Red Sea sponge, *Callyspongia* species. *J Nat Prod* 2003; 66: 679–681
- 115 Umeyama A, Matsuoka N, Mine R, Nakata A, Arimoto E, Matsui M, Shoji N, Arihara S, Takei M, Hashimoto T. Polyacetylene diols with anti-proliferative and driving Th1 polarization effects from the marine sponge *Callyspongia* sp. *J Nat Med* 2010; 64: 93–97
- 116 Rotem M, Kashman Y. New polyacetylenes from the sponge *Siphonochalina* sp. *Tetrahedron Lett* 1979; 20: 3193–3196
- 117 Tada H, Yasuda F. Siphonodiols, a new polyacetylenic metabolite from the sponge *Siphonochalina truncate*. *Chem Lett* 1984; 13: 779–780
- 118 Ayyad SE, Angawy R, Saqer EA, Badria FA. Cytotoxic polyacetylenes from the Red Sea sponge *Siphonochalina siphonella*. *Z Naturforsch C* 2014; 69: 117–123
- 119 Cimino G, de Stefano S. New acetylenic compounds from the sponge *Reniera fulva*. *Tetrahedron Lett* 1977; 18: 1325–1328
- 120 Ortega MJ, Zubía E, Carballo JL, Salvá J. Fulvinol, a new long-chain diacetylenic metabolite from the sponge *Reniera fulva*. *J Nat Prod* 1996; 59: 1069–1071
- 121 Nuzzo G, Ciavatta ML, Villani G, Manzo E, Zarfardino A, Varcamonti M, Gavagnin M. Fulvynes, antimicrobial polyoxygenated acetylenes from the Mediterranean sponge *Haliclona fulva*. *Tetrahedron* 2012; 68: 754–760
- 122 Williams DH, Faulkner DJ. Three new acetylenes from the Palauan sponge *Haliclona* sp. *J Nat Prod* 1996; 59: 1099–1101
- 123 Shin J, Seo Y, Cho KW, Rho JR, Paul VJ. Osirisynes A–F, highly oxygenated polyacetylenes from the sponge *Haliclona osiris*. *Tetrahedron* 1998; 54: 8711–8720
- 124 Chill L, Miroz A, Kashman Y. Haliclonyne, a new highly oxygenated polyacetylene from the marine sponge *Haliclona* species. *J Nat Prod* 2000; 63: 523–526
- 125 Aoki S, Matsui K, Tanaka K, Satari R, Kobayashi M. Lembehynes A, a novel neuritogenic polyacetylene, from a marine sponge of *Haliclona* sp. *Tetrahedron* 2000; 56: 9945–9948
- 126 Aoki S, Matsui K, Wei H, Murakami N, Kobayashi M. Structure-activity relationship of neuritogenic spongean acetylene alcohols, lembehynes. *Tetrahedron* 2002; 58: 5417–5422
- 127 Zhou GX, Molinski T. Long-chain acetylenic ketones from the Micronesian sponge *Haliclona* sp. Importance of the 1-yn-3-ol group for antitumor activity. *Mar Drugs* 2003; 1: 46–53
- 128 Aratake S, Trianto A, Hanif N, de Voogd N, Tanaka J. A new polyunsaturated brominated fatty acid from a *Haliclona* sponge. *Mar Drugs* 2009; 7: 523–527
- 129 Kobayashi M, Mahmud T, Tajima H, Wang W, Aoki S, Nakagawa S, Mayumi T, Kitagawa I. Marine natural products. XXXVI. Biologically active polyacetylenes, adociacetylenes A, B, C, and D, from an Okinawan marine sponge of *Adocia* sp. *Chem Pharm Bull* 1996; 44: 720–724
- 130 Wright AE, McConnell OJ, Kohmoto S, Lui MS, Thompson W, Snader KM. Duryne, a new cytotoxic agent from the marine sponge *Cribrachalina dura*. *Tetrahedron Lett* 1987; 28: 1377–1379
- 131 Gunasekera SP, Faircloth GT. New acetylenic alcohols from the sponge *Cribrachalina vasculum*. *J Org Chem* 1990; 55: 6223–6225
- 132 Aiello A, Fattorusso E, Menna M, Pansini M. Further bioactive acetylenic compounds from the caribbean sponge *Cribrachalina vasculum*. *J Nat Prod* 1992; 55: 1275–1280
- 133 Hallock YF, Cardellina JH, Balaschak MS, Alexander MR, Prather TR, Shoemaker RH, Boyd MR. Antitumor activity and stereochemistry of acetylenic alcohols from the sponge *Cribrachalina vasculum*. *J Nat Prod* 1995; 58: 1801–1807
- 134 Dai JR, Hallock YF, Cardellina JH, Boyd MR. Vasculyne, a new cytotoxic acetylenic alcohol from the marine sponge *Cribrachalina vasculum*. *J Nat Prod* 1996; 59: 88–89
- 135 Legrave N, Hamrouni-Buonomo S, Dufies M, Guérineau V, Vacelet J, Auberger P, Amade P, Mehiri M. Nepheliosyne B, a new polyacetylenic acid from the new caledonian marine sponge *Niphates* sp. *Mar Drugs* 2013; 11: 2282–2292
- 136 De Jesus RP, Faulkner DJ. Chlorinated acetylenes from the San Diego sponge *Haliclona lunisimilis*. *J Nat Prod* 2003; 66: 671–674
- 137 Castiello D, Cimino G, de Rosa S, de Stefano S, Sodano G. High molecular weight polyacetylenes from the nudibranch *Peltodoris actromaculata* and the sponge *Petrosia ficiformis*. *Tetrahedron Lett* 1980; 21: 5047–5050
- 138 Cimino G, Crispino A, de Rosa S, de Stefano S, Sodano G. Polyacetylenes from the sponge *Petrosia ficiformis* found in dark caves. *Experientia* 1981; 37: 924–926
- 139 Cimino G, de Giulio A, de Rosa S, de Stefano S, Sodano G. Further high molecular weight polyacetylenes from the sponge *Petrosia ficiformis*. *J Nat Prod* 1985; 48: 22–27
- 140 Cimino G, de Giulio A, de Rosa S, di Marzo V. High molecular weight polyacetylenes from *Petrosia ficiformis*: Further structural analysis and biological activity. *Tetrahedron Lett* 1989; 30: 3563–3566
- 141 Cimino G, de Giulio A, de Rosa S, di Marzo V. Minor bioactive polyacetylenes from *Petrosia ficiformis*. *J Nat Prod* 1990; 53: 345–353
- 142 Guo Y, Gavagnin M, Trivellone E, Cimino G. Absolute stereochemistry of petroformynes, high molecular polyacetylenes from the marine sponge *Petrosia ficiformis*. *Tetrahedron* 1994; 50: 13261–13268
- 143 Guo Y, Gavagnin M, Trivellone E, Cimino G. Further structural studies on the petroformynes. *J Nat Prod* 1995; 58: 712–722
- 144 Guo Y, Gavagnin M, Salierno C, Cimino G. Further petroformynes from both Atlantic and Mediterranean populations of the sponge *Petrosia ficiformis*. *J Nat Prod* 1998; 61: 333–337
- 145 Isaacs S, Kashman Y, Loya S, Hizi A, Loya Y. Petrosynol and petrosolic acid, two novel natural inhibitors of the reverse transcriptase of human immunodeficiency virus from *Petrosia* sp. *Tetrahedron* 1993; 49: 10435–10438
- 146 Guerriero A, Debitus C, Laurent D, d'Ambrosio M, Pietra F. Aztèquynol A, the first clearly defined, C-branched polyacetylene and the analogue Aztèquynol B. Isolation from the tropical marine sponge *Petrosia* sp. *Tetrahedron Lett* 1998; 39: 6395–6398

- 147 Fusetani N, Shiragaki T, Matsunaga S, Hashimoto K. Bioactive marine metabolites XX. Petrosynol and petrosynone, antimicrobial C30 polyacetylenes from the marine sponge *Petrosia* sp.: Determination of the absolute configuration. *Tetrahedron Lett* 1987; 28: 4313–4314
- 148 Fusetani N, Li HY, Tamura K, Matsunaga S. Antifungal brominated C18 acetylenic acids from the marine sponge, *Petrosia volcano* Hoshino. *Tetrahedron* 1993; 49: 1203–1210
- 149 Li H-Y, Matsunaga S, Fusetani N. Corticatic acids A–C, antifungal acetylenic acids from the marine sponge, *Petrosia corticata*. *J Nat Prod* 1994; 57: 1464–1467
- 150 Nishimura S, Matsunaga S, Shibasaki M, Suzuki K, Harada N, Naoki H, Fusetani N. Corticatic acids D and E, polyacetylenic geranylgeranyltransferase type I inhibitors, from the marine sponge *Petrosia corticata*. *J Nat Prod* 2002; 65: 1353–1356
- 151 Shin J, Seo Y, Cho KW. Five new polyacetylenes from a sponge of the genus *Petrosia*. *J Nat Prod* 1998; 61: 1268–1273
- 152 Seo Y, Cho KW, Rho J-R, Shin J, Sim CJ. Petrocortynes and petrosiacetylenes, novel polyacetylenes from a sponge of the genus *Petrosia*. *Tetrahedron* 1998; 54: 447–462
- 153 Kim JS, Lim YJ, Im KS, Jung JH, Shim CJ, Lee CO, Hong J, Lee H. Cytotoxic polyacetylenes from the marine sponge *Petrosia* sp. *J Nat Prod* 1999; 62: 554–559
- 154 Lim YJ, Lee CO, Hong J, Kim DK, Im KS, Jung JH. Cytotoxic polyacetylenic alcohols from the marine sponge *Petrosia* species. *J Nat Prod* 2001; 64: 1565–1567
- 155 Lim YJ, Park HS, Im KS, Lee CO, Hong J, Lee MY, Kim DK, Jung JH. Additional cytotoxic polyacetylenes from the marine sponge *Petrosia* species. *J Nat Prod* 2001; 64: 46–53
- 156 Lee YJ, Yoo SJ, Kang JS, Yun J, Shin HJ, Lee JS, Lee HS. Cytotoxic petrosiacetylenes from the marine sponge *Petrosia* sp. *Lipids* 2013; 48: 87–91
- 157 Juan YS, Lee CC, Tsao CW, Lu MC, El-Shazly M, Shih HC, Chen YC, Wu YC, Su JH. Structure elucidation and cytotoxic evaluation of new polyacetylenes from a marine sponge *Petrosia* sp. *Int J Mol Sci* 2014; 15: 16511–16521
- 158 Watanabe K, Tsuda Y, Hamada M, Omori M, Mori G, Iguchi K, Naoki H, Fujita T, van Soest RW. Acetylenic strongylodiols from a *Petrosia* (Strongylophora) Okinawan marine sponge. *J Nat Prod* 2005; 68: 1001–1005
- 159 Watanabe K, Mori G, Iguchi K, Suzuki M, van Soest RW. Nine acetylenic alcohols isolated from the Okinawan marine sponge of the genus *Petrosia* (Strongylophora). *Nat Prod Res* 2007; 21: 710–720
- 160 Okamoto C, Nakao Y, Fujita T, Iwashita T, van Soest RWM, Fusetani N, Matsunaga S. Cytotoxic C47-polyacetylene carboxylic acids from a marine sponge *Petrosia* sp. *J Nat Prod* 2007; 70: 1816–1819
- 161 Ueoka R, Ise Y, Matsunaga S. Cytotoxic polyacetylenes related to petroformyne-1 from the marine sponge *Petrosia* sp. *Tetrahedron* 2009; 65: 5204–5208
- 162 Hitora Y, Takada K, Okada S, Matsunaga S. Miyakosynes A–F, cytotoxic methyl branched acetylenes from a marine sponge *Petrosia* sp. *Tetrahedron* 2011; 67: 4530–4534
- 163 Ohta S, Ogawa T, Ohta E, Ikeuchi T, Kamemura K, Ikegami S. Petroacetylene, a new polyacetylene from the marine sponge *Petrosia solida* that inhibits blastulation of starfish embryos. *Nat Prod Res* 2013; 27: 1842–1847
- 164 Horikawa K, Yagyu T, Yoshioka Y, Fujiwara T, Kanamoto A, Okamoto T, Ojika M. Petrosiols A–E, neurotrophic diyne tetraols isolated from the Okinawan sponge *Petrosia strongylata*. *Tetrahedron* 2013; 69: 101–106
- 165 Schmitz FJ, Gopichand Y. (7E,13Z,15Z)-14,16-dibromo-7,13,15-hexadecatrien-5-ynoic acid. A novel dibromo acetylenic acid from the marine sponge *Xestospongia muta*. *Tetrahedron Lett* 1978; 19: 3637–3640
- 166 Quinn RJ, Tucker DJ. A brominated bisacetylenic acid from the marine sponge *Xestospongia testudinaria*. *Tetrahedron Lett* 1985; 26: 1671–1672
- 167 Quinn RJ, Tucker DJ. Further acetylenic acids from the marine sponge *Xestospongia testudinaria*. *J Nat Prod* 1991; 54: 290–294
- 168 Bourguet-Kondracki ML, Rakotoarisoa MT, Martin MT, Guyot M. Bioactive bromopolyacetylenes from the marine sponge *Xestospongia testudinaria*. *Tetrahedron Lett* 1992; 33: 225–226
- 169 Hirsh S, Carmely S, Kashman Y. Brominated unsaturated acids from the marine sponge *Xestospongia* sp. *Tetrahedron* 1987; 43: 3257–3261
- 170 Quiñoà E, Crews P. Melynes, polyacetylene constituents from a Vanuatu marine sponge. *Tetrahedron Lett* 1988; 29: 2037–2040
- 171 Kobayashi J, Naitoh K, Ishida K, Shigemori H, Ishibashi M. Nepheliosyne A, new C47 acetylenic acid from the okinawan marine sponge *Xestospongia* sp. *J Nat Prod* 1994; 57: 1300–1303
- 172 Brantley SE, Molinski TF, Preston CM, DeLong EF. Brominated acetylenic fatty acids from *Xestospongia* sp., a marine spongebacteria association. *Tetrahedron* 1995; 51: 7667–7672
- 173 Dai JR, Hallock YF, Cardellina JH, Gray GN, Boyd MR. Triangulynes A–H and triangulynic acid, new cytotoxic polyacetylenes from the marine sponge *Pellina triangulata*. *J Nat Prod* 1996; 59: 860–865
- 174 Fu X, Abbas SA, Schmitz FJ, Vidavsky I, Gross ML, Laney M, Schatzman RC, Cabuslay RD. New acetylenic metabolites from the marine sponge *Pellina triangulata*. *Tetrahedron* 1997; 53: 799–814
- 175 Matsunaga S, Okada Y, Fusetani N, van Soest RWM. An antimicrobial C14 acetylenic acid from a marine sponge *Oceanapia* species. *J Nat Prod* 2000; 63: 690–691
- 176 Erpenbeck D, van Soest RWM. Status and perspective of sponge chemosystematics. *Mar Biotechnol* 2007; 9: 2–19
- 177 Montalvo NF, Hill RT. Sponge-associated bacteria are strictly maintained in two closely related but geographically distant sponge hosts. *Appl Environ Microbiol* 2011; 77: 7207–7216
- 178 Trindade-Silva AE, Rua C, Silva GGZ, Dutilh BE, Moreira APB, Edwards RA, Hajdu E, Lobo-Hajdu G, Vasconcelos AT, Berlinck RGS, Thompson FL. Taxonomic and functional microbial signatures of the endemic marine sponge *Arenosclera brasiliensis*. *PLoS One* 2012; 7: e39905
- 179 Khan ST, Musarrat J, Alkhedhairi AA, Kazuo S. Diversity of bacteria and polyketide synthase associated with marine sponge *Haliclona* sp. *Ann Microbiol* 2014; 64: 199–207
- 180 Jasmin C, Anas A, Nair S. Bacterial diversity associated with *Cinachyra cavernosa* and *Haliclona pigmentifera*, cohabiting sponges in the Coral Reef Ecosystem of Gulf of Mannar, Southeast Coast of India. *PLoS One* 2015; 10: e0123222