Difficult enteral stenting made simple

Fig. 1 Fluoroscopic image showing injection of contrast using a ball-tip catheter to locate the duodenal stricture.

Fig. 2 Pressure injection of contrast under fluoroscopy to assess the length of the duodenal stricture.

Video 1

Radiographic imaging showing abdominal metastases and features of gastric outlet obstruction in a man with malignant duodenal stricture; endoscopic view of the stricture; fluoroscopic and endoscopic views showing a ball-tip catheter being carefully advanced across the stricture, an obstruction duodenogram to determine the length of the stricture, and a self-expandable metal stent being positioned across the stricture.

Enteral stenting is an accepted palliative option for malignant strictures. Long narrow duodenal strictures can be technically challenging [1,2]. A 55-year-old man with a history of Crohn's disease and a newly diagnosed malignant duodenal stricture presented with poor oral intake. Abdominal imaging revealed extensive metastatic disease with proximal duodenal obstruction without biliary obstruction. An upper gastrointestinal endoscopy with stent placement was performed. A gastroscope was advanced to the duodenal bulb, where the stricture was encountered but could not be traversed. Under endoscopic and fluoroscopic guidance, a ball-tip catheter with a preloaded guidewire was advanced into the stricture. Resistance to initial passage of the wire was encountered because of the stricture's tortuosity and length. A series of maneuvers, under fluoroscopic guidance with contrast injection, with slow advancement of the ball-tip catheter over the wire allowed the stricture to be crossed. The ball-tip catheter allowed for advancement of the wire with greater precision (● Fig. 1).

Once beyond the stricture, the wire was coiled in the distal small-bowel lumen. The ball-tip catheter was exchanged for a 15-mm extraction balloon, which was inflated in the small-bowel lumen just beyond the stricture. An occlusion duodenogram revealed a narrow stricture of 10 cm in length (● Fig. 2). Because of stent shortening during deployment, it is preferable to deploy a stent at least 2 cm longer than the stricture [3]. Therefore, a 22-mm × 12-cm uncovered self-expandable metal stent (SEMS) was deployed under endoscopic and fluoroscopic guidance. Gentle traction was applied to combat radial and propulsion forces during deployment. Contrast injection confirmed the patency and positioning of the stent (● Video 1). The patient was monitored post-procedure and was discharged home the same day, tolerating an oral diet. Endoscopic stent placement across long narrow duodenal strictures can be challenging. Use of a ball-tip catheter allows for precise advancement of the wire, leading to successful stenting of difficult enteral strictures.

Endoscopy_UCTN_Code_TTT_1AO_2AZ

Competing interests: Michel Kahaleh has received grant support from Boston Scientific, Fujinon, EMcison, Xlumen Inc., W.L. Gore, MaunaKea, Apollo Endosurgery, Cook Endoscopy, ASPIRE Bariatrics, GI Dynamics, NinePoint Medical, Merit Medical, Olympus and MI Tech. He is a consultant for Boston Scientific, Xlumen Inc., Concordia Laboratories Inc. and Mauna-Kea Tech.

Amit Desai, Amy Tyberg, Steve Zerbo, Reem Sharaiha, Michel Kahaleh

Division of Gastroenterology and Hepatology, Weill Cornell Medical College, New York, New York, USA

References

- 1 Kim JH, Song HY, Shin JH et al. Metallic stent placement in the palliative treatment of malignant gastroduodenal obstructions: prospective evaluation of results and factors influencing outcome in 213 patients. *Gastrointest Endosc* 2007; 66: 256–264

- 2 Chiu KW, Razack A, Maraveyas A. Self-expandable metal stent placement for malignant duodenal obstruction distal to the bulb Eur J Gastroenterol Hepatol 2015; 27: 1466–1472
- 3 Adler DG, Baron TH. Endoscopic palliation of malignant gastric outlet obstruction using self-expanding metal stents: experience in 36 patients. Am J Gastroenterol 2002; 97: 72–78

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-106963>
Endoscopy 2016; 48: E194–E195
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Michel Kahaleh, MD

Division of Gastroenterology and Hepatology
Weill Cornell Medical College
1305 York Avenue, 4th floor
New York
NY 10021
USA
mkahaleh@gmail.com