Subacute food bolus obstruction secondary to a migrated Overstitch suture from a previous esophageal perforation repair

The Overstitch device has been used for closure of esophageal perforations [1]. A 40-year-old woman with type I achalasia underwent fluoroscopic pneumatic dilation in September 2015. Her mean lower

esophageal sphincter (LES) pressure was 40 mmHg. She had 100% esophageal aperistalsis and absence of LES relaxation. Endoscopy showed a tight but traversable esophagogastric junction (EGJ). A 30-mm

achalasia balloon was used to perform dilation for 1 minute at 5 PSI followed by 1 minute at 8 PSI.

She became pyrexial on the third day after dilation. Computed tomography (CT) showed a distal esophageal perforation with a small paraesophageal collection and left basal pleural effusion (● Fig. 1). Intravenous antibiotics were commenced and ultrasound-guided aspiration of the pleural fluid yielded 8 mL of hemerosous fluid. An endoscopy on day 6 after dilation showed a 5-mm linear laceration above the EGJ (● Fig. 2a). It was repaired with two Overstitch polydioxanone (PDS) sutures using an Olympus 2T160 gastroscope (● Fig. 2b). Endoscopy and fluoroscopy 6 days after this repair showed that the repair was intact and the patient was discharged 13 days after the initial dilation. Gastroscopy at 4 weeks after repair showed good healing of the laceration with residual sutures at the repair site (● Fig. 2c). The patient's LES pressure was 20.7 mmHg; however, she was asymptomatic.

At follow-up at 5 months, the patient complained of spasmodic epigastric pain without vomiting or dysphagia. A subse-

Fig. 1 Computed tomography (CT) scan showing a distal esophageal perforation with extraluminal gas locules abutting the posterior surface of the descending thoracic aorta at the level of the hiatus, along with a small paraesophageal collection and left basal pleural effusion.

Fig. 2 a–e Endoscopic views showing: a a 5-mm linear deep laceration just above the esophagogastric junction; b repair of the laceration with Overstitch polydioxanone (PDS) sutures; c good healing of the laceration with a residual suture visible at the repair site 4 weeks after the Overstitch repair; d complete healing of the laceration and no sutures visible at the repair site 6 months after the Overstitch repair; e the suture that was found lodged at the D2/3 junction during the same endoscopic examination. f The Overstitch suture along with the food bolus around its T tag after it had been extracted endoscopically.

quent endoscopy showed good healing of the perforation site (● Fig. 2d); however, a suture with a 3-cm food bolus around its T tag was lodged at the D2/3 junction (● Fig. 2e). The suture and the food bolus was completely removed endoscopically using rat-tooth forceps (● Fig. 2f). The patient's symptoms resolved after this procedure.

Endoscopy_UCTN_Code_CPL_1AH_2AJ

Competing interests: None

Baldwin P. M. Yeung, Shannon M. Chan, Philip W. Y. Chiu

Department of Surgery, Prince of Wales Hospital, The Chinese University of Hong Kong, Hong Kong, China

Reference

- 1 Henderson JB, Sorser SA, Atia AN et al. Repair of esophageal perforations using a novel endoscopic suturing system. *Gastrointest Endosc* 2014; 80: 535–537

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-106965>
Endoscopy 2016; 48: E177–E178
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Baldwin P. M. Yeung, MBChB, PhD
Chinese University of Hong Kong – Surgery
30-32 Ngan Shing Street
Shatin
NT Hong Kong
Hong Kong
China
Fax: +852-26377974
byeung@doctors.net.uk