Accuracy of Point-of-Care B-Line Lung Ultrasound in Comparison to NT-ProBNP for Screening Acute Heart Failure

Authors

E. Glöckner¹, M. Christ², F. Geier¹, P. Otte³, U. Thiem⁴, S. Neubauer¹, V. Kohfeldt¹, K. Singler^{5,6}

Affiliations

Affiliation addresses are listed at the end of the article

Key words

- thorax
- chest ultrasound
- thoracic ultrasound
- emergency care

Abstract

Aim: The objective of this pilot study was to determine the accuracy of point-of-care B-line lung ultrasound in comparison to NT Pro-BNP for screening acute heart failure.

Materials and Methods: An 8-zone lung ultrasound was performed by experienced sonographers in patients presenting with acute dyspnea in the ED. AHF was determined as the final diagnosis by 2 independent reviewers.

Results: Contrary to prior studies, B-line ultrasound in our study was highly specific, but moderately sensitive for identifying patients with AHF. There was a strong association between elevated NT-proBNP levels and an increased number of B-lines.

Conclusion: In conclusion, point-of-care lung ultrasound is a helpful tool for ruling in or ruling out important differential diagnoses in ED patients with acute dyspnea.

received 21.10.2015
revised 04.03.2016
accepted 21.04.2016

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-108343>
Published online:
July 26, 2016
Ultrasound International Open
2016; 2: E90–E92
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 2199-7152

Correspondence

K. Singler

Department of Geriatrics
Klinikum Nuernberg
Prof. Ernst Nathan Str. 1
90419, Nürnberg
Germany
Tel.: +49/911 398 2434
katrin.singler@klinikum-nuernberg.de

Introduction

Acute dyspnoea is a common chief complaint of patients presenting to the emergency department (ED) and is associated with high morbidity and mortality. About 50% of dyspnoeic patients display acute heart failure (AHF) [1]. Rapid accurate diagnosis of AHF is hindered due to lacking sensitivity and specificity of clinical signs and symptoms. The primary aim of this pilot study was to determine the accuracy of point of care ultrasound for evaluation of acute dyspnea in the ED in comparison to circulating N-terminal pro-hormone of brain natriuretic peptide (NT-proBNP) levels.

Methods

We enrolled a convenience sample of 25 patients (≥ 18 years) presenting with undifferentiated acute dyspnoea to the ED of a German urban academic hospital. The study was approved by the local ethical committee. After obtaining written informed consent, 2 experienced sonographers performed an 8-zone lung ultrasound (LUS; 2–5 MHz phased array transducer, General Electric Vivid S6). Scans were evaluated offline by 2 medical experts blinded to clinical data (Cohens kappa=0.9). Positive ultrasound confirmation of

AHF was defined, as the bilateral existence of 2 or more positive regions with 3 or more B-lines. Clinical and demographic data, comorbidities, laboratory test results, transthoracic echocardiographic (TTE) data and 12-lead electrocardiography (ECG) data were obtained by reviewing all medical records available.

Final adjudicated diagnosis of acute heart failure was done by 2 experience physicians (cardiologist, emergency physician). If a disagreement about AHF diagnosis was present, a third experienced cardiologist settled disagreement [2, 3]. Primary endpoint was the accuracy of LUS for screening of AHF and compare it to circulating NT-proBNP levels. Continuous variables are presented as means ($\pm$ SD) or medians (interquartile range [IQR]), categorical variables as numbers and percentages. Comparisons of different diagnostic tools or in different subgroups of patients were performed using the Mann-Whitney U test for independent variables. For categorical data the Pearson chi-square, respectively Fisher's exact test was used. Proportions are described with 95% confidence intervals (CI). P-values < 0.05 were considered statistically significant. Data were analyzed using SPSS IBM Statistik 20 version for Windows, Munich, Germany.

License terms

Results

25 patients were included for analysis (Table 1). Median age was 72 years (IQR 60.5–80.5), 68% (n=17) were male and 76% (n=19) had a previous history of chronic heart failure (CHF). 60% (n=15) of patients had a final adjudicated diagnosis of AHF. Evaluation of laboratory tests, ECG, transthoracic echocardiography, chest X-ray and LUS are presented in Table 1. The sensitivity for LUS to detect AHF was 40% and the specificity 100% (Positive predictive value: 100%; negative predictive value: 52.6%). Analysis displays an association among the total number of B-lines and NT-proBNP levels in AHF patients ($p=0.005$, Table 1, Fig. 1).

Decline in amount of total B-lines after treatment accompanied decrease in symptoms as assessed by a VAS-dyspnoea scale.

Discussion

B-line LUS is proposed as a tool used at the point-of-care to support clinical decision making in patients with acute dyspnoea. [4–6]. In this pilot study, B-line ultrasound in the ED was highly specific, but moderately sensitive to identify patients with AHF. The major finding of our study corroborates and extends previous findings. 2 or more positive regions with 3 or more B-lines as dem-

variables	whole cohort, n = 25	AHF, n = 15	no AHF, n = 10	p =
demography				
age, median (Q1–Q3)	72 (60.5–80.5)	80 (65–81)	63.5 (40.25–72.25)	0.011
male, n (%)	17 (68.0)	11 (73.3)	6 (60.0)	0.667
vitals at first contact				
heart rate, bpm	90	88	92	0.698
systolic blood pressure, mmHg	138	136	140	0.651
diastolic blood pressure, mmHg	78	76	82	0.438
oxygen saturation, %	96.0	96.0	96.0	0.922
respiratory rate, bpm	17	18	16	0.64
temperature, C	36.7	36.7	36.7	0.897
history				
chronic heart failure, n (%)	19 (76.0)	15 (100)	4 (40.0)	0.001
previous myocardial infarction, n (%)	5 (20.0)	4 (26.7)	1 (10.0)	0.615
coronary artery disease, n (%)	14 (56.0)	11 (73.3)	3 (30.0)	0.032
hypertension, n (%)	22 (88.0)	15 (100)	7 (70.0)	0.052
chronic obstructive pulmonary disease, n (%)	6 (24.0)	3 (20.0)	3 (30.0)	0.653
malignancy, n (%)	5 (20.0)	4 (26.7)	1 (10.0)	0.615
final diagnosis				
cardiac, n (%)	15 (60.0)	14 (93.3)	1 (10.0)	<0.001
pulmonary, n (%)	3 (12.0)	0 (0.0)	3 (30.0)	0.052
mixed cardiac pulmonary, n (%)	1 (4.0)	1 (6.7)	0 (0.0)	1
other, n (%)	6 (24.0)	0 (0.0)	6 (60.0)	0.001
AHF, n (%)	15 (60.0)	15 (100)	0 (0.0)	
no AHF, n (%)	10 (40.0)	0 (0.0)	10 (100)	
laboratory test				
sodium, mmol/L	139.6	139	140.5	0.503
potassium, mmol/L	4.192	4.28	4.06	0.253
creatinine, mg/dL	1.23	1.45	0.9	0.007
BUN, mmol/L	23.75	30.72	13.31	0.001
NT-proBNP, (pg/ml)	5244	7214	320	0.005
cTnT-hs, ng/L	0.0391	0.0443	0.0312	0.005
12-lead ECG				
sinus rhythm, n (%)	15 (60.0)	6 (40.0)	9 (90.0)	0.012
atrial fibrillation, n (%)	9 (36.0)	8 (53.3)	1 (10.0)	0.027
other rhythm, n (%)	1 (4.0)	1 (6.7)	0 (0.0)	1
left bundle branch block, n (%)	3 (12.0)	2 (13.3)	1 (10.0)	1
right bundle branch block, n (%)	1 (4.0)	0 (0.0)	1 (10.0)	0.4
pacemaker rhythm, n (%)	1 (4.0)	1 (6.7)	0 (0.0)	1
ST elevation, n (%)	0 (0.0)	0 (0.0)	0 (0.0)	–
myocardial ischemia, n (%)	5 (20.0)	3 (20.0)	2 (20.0)	1
lung ultrasound				
LUS positive for AHF, n (%)	6 (24.0)	6 (40.0)	0 (0.0)	0.051
echocardiography				
LVEDD, mm	56.1	56.7	54.5	0.777
EF, %	44.6	40.5	54.8	0.062
LA, ml	90.8	93.9	83	1
chest X-ray				
Overall impression, (mean)	1.9	2.1	1.6	0.059
1, n (%)	6 (24.0)	2 (13.3)	4 (40.0)	0.175
2, n (%)	16 (64.0)	10 (66.7)	6 (60.0)	0.734
3, n (%)	3 (12.0)	3 (20.0)	0 (0.0)	0.25
cardiothoracic ratio	0.54	0.58	0.48	<0.001

Table 1 Patient characteristics and diagnostic parameters.

Fig. 1 Boxplot of B-lines by different NT-proBNP groups in patients with acute dyspnea (n = 14 patients in total with available NT-proBNP values). The graph shows a strong association among elevated NT-proBNP levels and the increased number of B-lines.

onstrated by point-of-care LUS display a very high specificity to support the diagnosis of AHF in ED patients with acute dyspnea (specificity 75–91%) [7,8]. In contrast to previous examinations, diagnostic sensitivity in our trial (40%) is moderate as compared to previous reports (70–92%) [7,8]. It is tempting to speculate that this may be due to the method of final adjudication of AHF diagnosis in respective trials: Final adjudicated diagnosis of AHF in our trial was done by 2 experienced physicians (cardiologist, emergency physician) using all available patient data for adjudication [1,10], while other reports used scores including descriptive methods of pulmonary congestions and/or cardiac decompensation for final diagnosis questioning accuracy of final adjudicated diagnosis [6,8]. Of note, some of the enrolled patients were pre-treated by EMS based emergency physicians before being admitted to ED. Use of pre-hospital diuretics may have influenced the presence of B-lines due to their rapid resolution after treatment. Nevertheless, emergency LUS of study subjects was performed within 1 h of presentation to the ED confirming the validity of our findings. Miglironza found a sensitivity for LUS of 85% in outpatients with a mean age of 53 years [8]. Our cohort was older with a median age of 72 years (IQR 60.5–80.5) and contrary to prior ED studies a history of CHF was present in nearly all patients (vs. 75% described by Anderson, 2013) [9].

As demonstrated by our data, radiographic signs show a moderate diagnostic accuracy to detect AHF [11], while NT-proBNP levels display a high diagnostic accuracy for identifying AHF as the cause of acute dyspnea [12]. In addition, it has to be recognized that LUS technique to correctly identify B-lines is not fully standardized. In particular, heterogeneity of equipment (type of probe used, frequency), and individual technique used (how long are the pleural line segments visualized at each spot, type of the probe, pressure applied, scanning sector angle) may influence the diagnostic accuracy of LUS. The conclusions of our findings may also be limited due to the low sample size and bias may have occurred due to the convenience sample of this trial. The strong association among elevated NT-proBNP levels and the increased number of B-Lines (≥ 12) confirms the validity and strength of our data, which has also been shown by previous reports [13]. Adequately powered diagnostic studies are required to confirm the utility of this evaluation strategy.

Conclusion

Lung ultrasound appears to be a helpful screening method for patients with acute dyspnea regarding AHF. LUS is a rapid, timely available and very specific method to support the diagnosis of AHF in patients presenting with acute dyspnea to the ED. In contrast to circulating natriuretic peptide levels, sensitivity of LUS is intermediate. Therefore a combined strategy of NT-proBNP testing and LUS will lead to fast and reliable results to support the diagnosis of AHF.

Affiliations

- ¹ Geriatrics, Klinikum Nuernberg, Paracelsus Medical University, Nürnberg, Germany
- ² Department of Emergency and Critical Care Medicine, Klinikum Nürnberg, Paracelsus Medical University, Nürnberg, Germany
- ³ Radiology, Klinikum Nuernberg, Paracelsus Medical University, Nürnberg, Germany
- ⁴ Department of Geriatrics, Marienhospital Herne, University Bochum, Nürnberg, Germany
- ⁵ Department of Geriatrics, Klinikum Nuernberg, Paracelsus Medical University, Nürnberg, Germany
- ⁶ Institute for Biomedicine of Aging, Friedrich-Alexander University Erlangen-Nuremberg, Nürnberg, Germany

References

- 1 Christ M, Laule-Kilian K, Hochholzer W et al. Gender-specific risk stratification with B-type natriuretic peptide levels in patients with acute dyspnea: insights from the B-type natriuretic peptide for acute shortness of breath evaluation study. *J Am Coll Cardiol* 2006; 48: 1808–1812
- 2 McMurray JJ, Adamopoulos S, Anker SD et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J* 2012; 33: 1787–1847 Erratum in: *Eur Heart J* 2013; 34: 158
- 3 Nieminen MS, Böhm M, Cowie MR et al. Executive summary of the guidelines on the diagnosis and treatment of acute heart failure: the Task Force on Acute Heart Failure of the European Society of Cardiology. *Eur Heart J* 2005; 26: 384–416
- 4 Volpicelli G, Elbarbary M, Blaivas M et al. International evidence-based recommendations for point-of-care lung ultrasound. *Intensive Care Med* 2012; 38: 577–591
- 5 Lichtenstein DA, Mezière GA. Relevance of lung ultrasound in the diagnosis of acute respiratory failure: the BLUE protocol. *Chest* 2008; 134: 117–125
- 6 Neskovic AN, Hagendorff A, Lancellotti P et al. Emergency echocardiography: the European Association of Cardiovascular Imaging recommendations. *European Association of Cardiovascular Imaging. Eur Heart J Cardiovasc Imaging* 2013; 14: 1–11
- 7 Bitar Z, Maadarani O, Almerri K. Sonographic chest B-lines anticipate elevated B-type natriuretic peptide level, irrespective of ejection fraction. *Ann Intensive Care* 2015; 5: 56
- 8 Miglironza MH, Gargani L, Sant'Anna RT et al. Lung ultrasound for the evaluation of pulmonary congestion in outpatients: a comparison with clinical assessment, natriuretic peptides, and echocardiography. *JACC Cardiovasc Imaging* 2013; 6: 1141–1151
- 9 Anderson KL, Jenq KY, Fields JM et al. Point-of-care ultrasound diagnoses acute decompensated heart failure in the ED regardless of examination findings. *Am J Emerg Med* 2014; 32: 385–388
- 10 Bahrmann P, Christ M, Hofner B, Bahrmann A, Achenbach S, Sieber CC, Bertsch T. Prognostic value of different biomarkers for cardiovascular death in unselected older patients in the emergency department. *Eur Heart J Acute Cardiovasc Care* 2015; 19:
- 11 Studler U, Kretzschmar M, Christ M, Breidthardt T, Noveanu M, Schoetzau A, Perruchoud AP, Steinbrich W, Mueller C. Accuracy of chest radiographs in the emergency diagnosis of heart failure. *Eur Radiol* 2008; 18: 1644–1652
- 12 Mueller C, Scholer A, Laule-Kilian K et al. Use of B-type natriuretic peptide in the evaluation and management of acute dyspnea. *N Engl J Med* 2004; 350: 647–654
- 13 Gargani L, Frassi F, Soldati G et al. Ultrasound lung comets for the differential diagnosis of acute cardiogenic dyspnea: a comparison with natriuretic peptides. *Eur J Heart Fail* 2008; 10: 70–77