Submucosal tunneling endoscopic resection for multiple esophageal leiomyomas

Fig. 1 Endoscopic view showing two protruding lesions in the mid-esophagus.

Fig. 2 Endoscopic ultrasonography (EUS) view showing that the tumors were originating from the muscularis propria layer.

Fig. 3 Computed tomography (CT) scan of the lesion, which was suggestive of benign tumors.

A 61-year-old man presented for treatment of esophageal submucosal tumors (SMTs). The SMTs had been found 1 month previously at his local hospital during a health examination. Upon presentation, his physical examination and laboratory tests were unremarkable. Esophago-gastroduodenoscopy (EGD) revealed two protruded lesions in the middle of the esophagus (• **Fig. 1**; • **Video 1**). Endoscopic ultrasonography (EUS) revealed that the tumors were originating from the muscularis propria layer (• **Fig. 2**). Computed tomography (CT) also showed the lesions in the mid-esophagus, which were suspected to be benign tumors (• **Fig. 3**; • **Video 2**).

A submucosal tunneling endoscopic resection (STER) was performed. After a longitudinal mucosal incision had been made, a submucosal tunnel was created, which allowed us to see the first SMT. Submucosal injection of methylene blue was performed to help locate the second tumor, and two separate tumors about 2 cm apart from each other could then be seen (• **Fig. 4a**; • **Video 3**). The tumors were carefully dissected off the muscularis propria layer (• **Fig. 4b**). The tunnel entry had to be enlarged to allow successful extraction of the larger SMT (• **Video 4**). The mucosal entry was then closed. The STER procedure was completed uneventfully within 110 minutes. The resected SMTs measured 3.8×3.2 cm and 1.5×0.9 cm (• **Fig. 5**) and, histopathologically, they were both leiomyomas.

Endoscopy showing two protruding lesions in the esophagus.

Computed tomography (CT) scan of the lesion.

STER has been demonstrated to be safe and effective for treating upper gastrointestinal SMTs; most of the reported cases were with solitary and small SMTs (≤ 3.5 cm) [1]. Although several cases/studies have been reported regarding STER for multiple SMTs and large SMTs,

the procedure is technically difficult and has a higher rate of complications [2–5]. In the present case, two SMTs were found, which were not strictly in a straight plane and one of them was larger than 3.5 cm. We successfully removed them both using the STER technique uneventfully within a

Fig. 4 Endoscopic views showing: **a** the two tumors found in the submucosal tunnel; **b** the wound surface after removal of the tumors.

Fig. 5 Macroscopic appearance of the resected tumors.

Video 3

After the first tumor had been successfully exposed, the second one could not be seen so methylene blue was injected submucosally to help locate it.

Video 4

En bloc extraction of the larger tumor was difficult, so the mucosotomy was enlarged, the tumor then dropped into the stomach allowing it to be extracted en bloc with a snare.

single submucosal tunnel after locating the second tumor with methylene blue and enlarging the tunnel entry.

Endoscopy_UCTN_Code_TTT_1AO_2AG

Competing interests: None

Yuyong Tan, Changmei Hu, Yuqian Zhou, Yi Chu, Jirong Huo, Deliang Liu

Department of Gastroenterology,
The Second Xiangya Hospital of Central
South University, Hunan, China

References

- 1 Chen T, Zhou PH, Chu Y et al. Long-term outcomes of submucosal tunneling endoscopic resection for upper gastrointestinal submucosal tumors. *Ann Surg* 2016; doi: 10.1097/SLA.0000000000001650
- 2 Wang H, Tan Y, Zhou Y et al. Submucosal tunneling endoscopic resection for upper gastrointestinal submucosal tumors originating from the muscularis propria layer. *Eur J Gastroenterol Hepatol* 2015; 27: 776–780
- 3 Chen T, Zhang C, Yao LQ et al. Management of the complications of submucosal tunneling endoscopic resection for upper gastrointestinal submucosal tumors. *Endoscopy* 2016; 48: 149–55

- 4 Chen H, Xu Z, Huo J et al. Submucosal tunneling endoscopic resection for simultaneous esophageal and cardia submucosal tumors originating from the muscularis propria layer (with video). *Dig Endosc* 2015; 27: 155–158
- 5 Zhang C, Hu JW, Chen T et al. Submucosal tunneling endoscopic resection for upper gastrointestinal multiple submucosal tumors originating from the muscularis propria layer: a feasibility study. *Indian J Cancer* 2015; 51 (Suppl. 02): e52–e55

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-109602>
Endoscopy 2016; 48: E234–E235
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Delian Liu, MD
Department of Gastroenterology
The Second Xiangya Hospital
of Central South University
NO. 139 Middle Renmin Road
Changsha
Hunan, 410011
China
Fax: 86-0731-85533525
liudeliang@medmail.com.cn