Novel technique to relax the lower esophageal sphincter during challenging peroral endoscopic myotomy (POEM)

Fig. 1 Illustration of the effect of amyl nitrite on lower esophageal sphincter relaxation during peroral endoscopic myotomy (POEM).

The use of amyl nitrite inhalation to relax the lower esophageal sphincter and facilitate submucosal tunneling in two patients with achalasia.

Amyl nitrite is easily administered and is a potent short-acting smooth muscle relaxant. Therefore, it can be used to relax the lower esophageal sphincter (LES) during submucosal tunneling in peroral endoscopic myotomy (POEM) (Fig. 1) [1,2]. One study demonstrated that the diameter of the LES increased by >3 mm in patients with idiopathic achalasia in response to amyl nitrite [3]. POEM involves the creation of a submucosal tunnel, for which dissection of fibers should be car-

ried out close to the muscle layer to avoid mucosal injury. When the LES is tight, this dissection of the fibers can be challenging to perform safely without risking mucosal injury. Video 1 demonstrates the use of amyl nitrite inhalation to relax the LES and facilitate submucosal tunneling in two patients.

The first patient was a 59-year-old man who underwent POEM for type III achalasia. During tunneling, the distal esophagus and LES were spastic, which inhibited the advancement of the endoscope. The decision was made to use amyl nitrite. During its administration, the patient's heart rate increased by 10 beats per minute (bpm) from a baseline of 70 bpm and his blood pressure (BP) decreased to 95/68 mmHg from 117/70 mmHg. The second patient was a 38-year-old woman who underwent POEM for type II achalasia. During POEM her LES was found to be tight, which prevented effective and safe tunneling. When amyl nitrite was given, her heart rate increased by 10 bpm from baseline and her systolic BP dropped to 80 mmHg from a baseline of 116 mmHg.

Both patients were admitted to the hospital for observation following their procedures. They were discharged home on

soft diet and returned to clinic 4 weeks later with complete resolution of their symptoms.

We conclude that amyl nitrite is a safe and effective method to relax the LES. This technique may be utilized in patients where the safe creation of a submucosal tunnel through the LES might otherwise not have been possible.

Endoscopy_UCTN_Code_TTT_1AO_2AN

Competing interests: Dr. Khashab is a consultant for Boston Scientific. All other authors have no relevant disclosures.

Majidah Bukhari¹, Vivek Kumbhari¹, Yamile Haito-Chavez¹, Yen-I Chen¹, Saowanee Ngamruengphong¹, Payal Saxena², Mouen A. Khashab¹

¹ Department of Medicine and Division of Gastroenterology and Hepatology, The Johns Hopkins Medical Institutions, Baltimore, Maryland, USA

² Department of Medicine and Division of Gastroenterology and Hepatology, Royal Prince Alfred Hospital, Camperdown, New South Wales, Australia

References

- 1 Kahrilas PJ, Kishk SM, Helm JF et al. Comparison of pseudoachalasia and achalasia. *Am J Med* 1987; 82: 439–446
- 2 Gonzalez M, Mearin F, Vasconez C et al. Oesophageal tone in patients with achalasia. *Gut* 1997; 41: 291–296
- 3 Dodds WJ, Stewart ET, Kishk SM et al. Radiologic amyl nitrite test for distinguishing pseudoachalasia from idiopathic achalasia. *AJR Am J Roentgenol* 1986; 146: 21–23

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-109776>
Endoscopy 2016; 48: E252
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Mouen A. Khashab, MD
Therapeutic Endoscopy
Johns Hopkins Hospital
Sheikh Zayed Bldg
Division of Gastroenterology and Hepatology
1800 Orleans Street, Suite 7125B
Baltimore
MD 21287
USA
Fax: +1-443-683-8335
mkhasha1@jhmi.edu