Submucosal tunneling endoscopic resection for leiomyomas located on opposite esophageal walls: the role of a tortuous submucosal tunnel

Fig. 1 Endoscopic views showing: **a** two protruding lesions in the esophagus; **b** the first tumor in the submucosal tunnel; **c** the second tumor within the same submucosal tunnel; **d** the wound surface after removal of the two tumors; **e** the mucosal entry point closed with several clips.

A 50-year-old man presented to our hospital with a 1-month history of epigastric discomfort. Esophagogastroduodenoscopy (EGD) revealed two protruding lesions on the anterior and posterior esophageal walls, about 27 cm and 30 cm from the incisors, respectively (● **Fig. 1 a**; ● **Video 1**). Endoscopic ultrasonography (EUS) revealed that the tumors were originating from the muscularis propria layer and had no risk features.

Submucosal tunneling endoscopic resection (STER) was performed. After a longitudinal mucosal incision had been made, a submucosal tunnel was created and the first submucosal tumor (SMT) could be seen (● **Fig. 1 b**). As the two tumors were located on opposite walls of the esophagus, it was impossible to locate the second tumor using purely a straight submucosal tunnel, so a tortuous tunnel was created. Submucosal injection of methylene blue

was performed to preset the tunnel route, and the two separate tumors were identified, with a distance of about 3 cm between them (● **Fig. 1 c**). The tumors were carefully dissected off the muscularis propria layer (● **Fig. 1 d**), and the mucosal entry was closed (● **Fig. 1 e**; ● **Video 2**). The STER procedure was completed uneventfully within 80 minutes. The diameters of the resected SMTs were 2 cm and 1.5 cm (● **Fig. 2**) and they were both leiomyomas histopathologically.

STER has been demonstrated to be safe and effective for treating upper gastrointestinal SMTs [1, 2]. There have been only a few reports regarding STER for multiple SMTs, and the tumors in each of the reported cases were in the same or adjacent parts of the esophageal/gastric wall, so a single submucosal tunnel was sufficient to remove the tumors [3–5]. In the present case, the two SMTs were found on opposite

Fig. 2 Macroscopic appearance of the two resected tumors.

esophageal walls, and at first two submucosal tunnels were recommended. This however would mean excessive submucosal dissection and two tunnel entries, which, as well as being time-consuming,

Video 1

Endoscopy showing two protruding lesions on opposite walls of the esophagus.

Video 2

The submucosal tunneling endoscopic resection (STER) procedure being performed for two tumors on opposite esophageal walls.

would increase the complication rate. We successfully removed both of the tumors by STER within a single tunnel by creating a tortuous submucosal tunnel.

Endoscopy_UCTN_Code_TTT_1AO_2AG

Competing interests: None

Fuxi Zhou, Yuyong Tan, Yi Chu, Changmei Hu, Jirong Huo, Deliang Liu

Department of Gastroenterology,
The Second Xiangya Hospital of Central
South University, Changsha, Hunan, China

References

- 1 Chen T, Zhou PH, Chu Y et al. Long-term outcomes of submucosal tunneling endoscopic resection for upper gastrointestinal submucosal tumors. *Ann Surg* 2016. Epub ahead of print 2016 Jan 19
- 2 Wang H, Tan Y, Zhou Y et al. Submucosal tunneling endoscopic resection for upper gastrointestinal submucosal tumors originating from the muscularis propria layer. *Eur J Gastroenterol Hepatol* 2015; 27: 776–780
- 3 Chen H, Xu Z, Huo J et al. Submucosal tunneling endoscopic resection for simultaneous esophageal and cardia submucosal tumors originating from the muscularis propria layer (with video). *Dig Endosc* 2015; 27: 155–158
- 4 Zhang C, Hu JW, Chen T et al. Submucosal tunneling endoscopic resection for upper gastrointestinal multiple submucosal tumors originating from the muscularis propria layer: a feasibility study. *Indian J Cancer* 2015; 51 (Suppl. 02): e52–e55
- 5 Tan Y, Hu C, Zhou Y et al. Submucosal tunneling endoscopic resection for multiple esophageal leiomyomas. *Endoscopy* 2016; 48 (Suppl. 01): E234–E235

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-111323>
Endoscopy 2016; 48: E255–E256
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Delian Liu, MD

Department of Gastroenterology
 The Second Xiangya Hospital of Central South
 University
 NO. 139 Middle Renmin Road
 Changsha
 Hunan, 410011
 China
 Fax: 86-0731-85533525
liudeliant@medmail.com.cn