Endoscopic ultrasound-guided gastrojejunostomy with a Nagi stent for relief of jejunal loop obstruction following hepaticojejunostomy

Fig. 1 Contrast-enhanced computed tomography (CT) scan of the abdomen showing a dilated loop of jejunum with an intact hepaticojejunostomy anastomosis.

Fig. 2 Endoscopic ultrasound (EUS) images showing: **a** the puncture of the dilated jejunal loop with a 10-Fr cystotome; **b** the Niti-S Nagi stent being deployed over the wire.

Fig. 3 Endoscopic views showing: **a** the Niti-S Nagi stent in position allowing free flow of bile across the gastrojejunostomy; **b** a 7-Fr double-pigtail plastic stent (7 cm in length) that has been inserted through the stent to prevent migration.

Video 1

Radiographic and endoscopic views of the dilated jejunal loop being punctured, a Nagi stent being positioned across the gastrojejunostomy, with a double-pigtail stent positioned through it to prevent migration.

A 27-year-old man with unresectable periampullary neoplasm underwent a Roux-en-Y hepaticojejunostomy with gastrojejunostomy 1 year previously. He presented with a 1-month history of progressive yellowish discoloration of his eyes, abdominal pain and distension, and intermittent high fever with chills and rigors. On admission his results showed a bilirubin of 9 mg/dL, alkaline phosphatase of 1467 IU/mL, and total white blood cell (WBC) count of 28 700/mm³. A blood culture grew *Escherichia coli*. A contrast-enhanced computed tomography (CT) scan of the abdomen showed a grossly dilated loop of jejunum with distal obstruction, in communication with a patent hepaticojejunostomy anastomosis (Fig. 1).

We planned to decompress the jejunal loop to relieve the cholangitis; however, the presence of a distal obstruction precluded enteroscope-assisted drainage. The distended jejunal loop was punctured with a 10-Fr cystotome (Cook Medical, Winston-Salem, North Carolina, USA) using endoscopic ultrasound (EUS) guidance (Fig. 2a). The needle was removed and a 0.035-inch guidewire was placed through the inner catheter into the jejunal loop. The over-the-wire 10-Fr outer catheter of the cystotome with a diathermic ring was advanced into the jejunal loop using pure cut (Video 1). The cystotome was then removed. A 30-mm Niti-S Nagi stent (Taewoong Medical, Seoul, South Korea)

was placed across the tract between the stomach and jejunal limb (► Fig. 2b), and bile was seen to drain through the gastro-jejunal anastomosis (► Fig. 3a). To prevent migration of the stent, a 7-Fr double-pig-tail plastic stent (7 cm in length) was placed across the Nagi stent (► Fig. 3b).

After the procedure the patient's fever subsided, his bilirubin decreased to 3.1 mg/dL, his total WBC count decreased to 15 200/mm³ and he was moved from the intensive care unit. He was discharged from hospital after 15 days.

Obstruction of the jejunal loop can occur as a late complication of hepaticojejunostomy due to either adhesions or tumor recurrence. EUS-guided gastrojejunostomy for afferent-loop syndrome using a Hot Axios stent was first described by Ikeuchi et al. in 2015 [1]. Mutignani et al. recently used the Nagi stent to gain access to the jejunal loop in a patient with an hepaticojejunostomy [2]. In our patient, we used the Nagi stent to decompress the obstructed jejunal loop. The procedure appears to be a safe alternative for the management of patients with hepaticojejunostomy presenting with cholangitis secondary to jejunal loop obstruction.

Endoscopy_UCTN_Code_TTT_1AS_2AD

Competing interests: None

**Sudipta Dhar Chowdhury,
Reuben Thomas Kurien, A. Kumar
Bharath, Amit Kumar Dutta,
Deepu David, C. Kumar Bharath,
Anjilivelil Joseph Joseph**

Department of Gastroenterology, Christian Medical College, Vellore, Tamil Nadu, India

References

- 1 Ikeuchi N, Itoi T, Tsuchiya T et al. One-step EUS-guided gastrojejunostomy with use of lumen-apposing metal stent for afferent loop syndrome treatment. *Gastrointest Endosc* 2015; 82: 166
- 2 Mutignani M, Manta R, Pugliese F et al. Endoscopic ultrasound-guided duodenojejunal anastomosis to treat postsurgical Roux-en-Y hepaticojejunostomy stricture: a dream or a reality? *Endoscopy* 2015; 47 (Suppl. 01): E350–E351

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-112971>
Endoscopy 2016; 48: E263–E264
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Sudipta Dhar Chowdhury, MD
Department of Gastroenterology
Christian Medical College
Vellore
Tamil Nadu
India
Fax: +91-416-2282054
sudipto.d.c@gmail.com
sudipta@cmcvellore.ac.in