The use of stand-alone fully covered self-expandable metallic stents for drainage of walled-off necrosis or pseudocyst may lead to poor results

In current practice, endoscopic transmural drainage is the preferred method for the treatment of symptomatic walled-off necrosis and pseudocysts in patients with acute pancreatitis [1]. In order to maintain a fistulous tract between these lesions and the gut lumen, fully covered self-expandable metallic stents (FCSEMS) and/or plastic pigtail stents can be used. The migration of FCSEMS to the gut lumen can lead to the disconnection of the cavity from the gut lumen, abscess development in the cavity, and life-threatening septic complications.

We present the case of a 20-year-old man with acute biliary pancreatitis who was admitted to our hospital with abdominal pain, early satiety, nausea, and vomiting. Endoscopic ultrasound and computed tomography scans demonstrated a pseudocyst (diameter 71 × 126 mm) in the tail of the pancreas. The gastric corpus was compressed by the cyst (● Fig. 1). Endoscopic drainage was planned.

Cystogastrostomy was performed and an FCSEMS (Micro-Tech, 16 × 30 mm; Nanjing, Jiangsu Province, China) was placed into the fistulous tract (● Fig. 2). After the procedure, the patient's symptoms improved. However, he developed fever, leukocytosis, and progressive abdominal pain on the third postoperative day. On plain abdominal radiography, the metal stent was seen to be located in the rectum (● Fig. 3). Endoscopy was performed and large amounts of pus was observed coming from the orifice of the cystogastrostomy into the gastric lumen. A nasocystic drainage catheter was placed into the cyst cavity via the same orifice. The patient expelled the stent on defecation the same day.

Fever and leukocytosis resolved following nasocystic drainage and broad-spectrum antibiotic therapy. The nasocystic drainage catheter was then exchanged for two 10-Fr double-pigtail plastic stents, and the patient was discharged.

Fig. 1 Computed tomography image of pseudocyst compressing the stomach (arrow).

Fig. 2 Computed tomography image of the metal stent (arrow) in the cystogastrostomy.

There are several well-known complications related to the metal stents used in cystogastrostomy. Migration is one of them and can occur either into the cystic cavity or into the gut lumen [2]. It is important to fix FCSEMS in place using a double-pigtail stent (stent-in-stent arrangement) or by the addition of a nasocystic drainage catheter in order to prevent migration and related complications.

Endoscopy_UCTN_Code_CPL_1AL_2AD

Competing interests: None

**Bulent Odemis¹, Erkin Oztas¹,
Muhammet Yener Akpınar¹,
Sabite Kacar¹, Ufuk Baris Kuzu¹,
Mustafa Ozdemir²**

¹ Department of Gastroenterology, Türkiye Yüksek İhtisas Training and Research Hospital, Ankara, Turkey

² Department of Radiology, Türkiye Yüksek İhtisas Training and Research Hospital, Ankara, Turkey

Fig. 3 Metal stent (arrow) in the rectum on plain abdominal radiograph.

References

- 1 Schmidt PN, Novovic S, Roug S et al. Endoscopic, transmural drainage and necrosectomy for walled-off pancreatic and peripancreatic necrosis is associated with low mortality – a single-center experience. *Scand J Gastroenterol* 2015; 50: 611–618
- 2 Walter D, Will U, Sanchez-Yague A et al. A novel lumen-apposing metal stent for endoscopic ultrasound-guided drainage of pancreatic fluid collections: a prospective cohort study. *Endoscopy* 2015; 47: 63–67

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-112978>
Endoscopy 2016; 48: E271–E272
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Muhammet Yener Akpınar, MD

Türkiye Yüksek İhtisas Training
 and Research Hospital
 Kızılay Street
 Sıhhiye
 Ankara 06100
 Turkey
 Fax: +90-312-3124120
muhammet.yener@gmail.com