Salvage endoscopic ultrasound–radiologic rendezvous to re-establish biliary flow following duodenectomy with ablation of papilla

Fig. 1 a, b Staple line on the common bile duct (arrow) with dilation of the biliary tree and a normal pancreatic duct in a 54-year-old patient after curative duodenectomy with duodenojejunal anastomosis for adenocarcinoma.

Fig. 2 Looping of the guidewire in the cystic stump.

Video 1
Endoscopic ultrasound (EUS)-guided cholangiography showing dilatation of the common bile duct with no passage into the duodenum because of complete transection of the papilla after duodenectomy with duodenojejunal anastomosis.

Video 2
Fistulotomy on guidewire allowing a communication between the duodenum and the common bile duct (CBD) to be created. The wall of the lower CBD can be seen.

Fig. 3 Passage of the guidewire by percutaneous transhepatic cholangiography, through the common bile duct and into the duodenum via the bilioduodenal access previously created under endoscopic ultrasound (EUS).

Fig. 4 Fully covered metal and plastic stents in place, restoring the biliary flow.

A 54-year-old woman underwent curative duodenectomy with duodenojejunal anastomosis for adenocarcinoma. The day after the procedure she developed jaundice. The computed tomography (CT) scan showed a staple line on the lower part of the common bile duct (CBD) with biliary tree dilatation (Fig. 1) and a normal pancreatic duct. Under endoscopic ultrasound (EUS) control a transbulbar cholangiography was performed on the lower part of the CBD, confirming biliary dilatation without any leakage of contrast medium into the duodenum (Video 1). A fistulotomy (6Fr) was then performed (Video 2). Although different guidewires and catheters were used, selective cannulation of the cystic stump was achieved without reaching the intrahe-

patic ducts because of the anatomically low cystic duct insertion (Fig. 2). A transhepatic cholangiography was performed. Passage of the guidewire into the duodenum was easily done through the bilioduodenal access previously created by EUS fistulotomy (Fig. 3). A fully covered self-expanding metal stent (FCSEMS) and a plastic stent were delivered to maintain biliary flow (Fig. 4). Follow-up was uneventful with normalization of liver function test results, and endoscopic retrograde cholangiopancreatography (ERCP) was scheduled for stent replacement.

Iatrogenic injury of the biliary tree following surgery on organs of the epigastrium other than the biliary tract is well known [1]. Successful EUS-guided choledoch-

duodenostomy has been described using a lumen-apposing FCSEMS after failed ERCP [2]. Perez-Miranda et al. reported successful repair, by EUS-guided choledochoduodenostomy and magnets, of a disconnected CBD following orthotopic liver transplantation [3].

In this particular case we did not use a lumen-apposing SEMS; this was to avoid sump syndrome and malfunction once the CBD was empty. We decided to perform a EUS–radiologic rendezvous [4,5] in order to deliver the FCSEMS in a position that more nearly replicated the normal anatomy to prevent possible later adverse events. In conclusion, interventional EUS is the key technique with which to access the biliary tree in cases of altered anatomy and EUS–radiologic rendezvous helps in difficult situations, sparing the need for major surgery with its related morbidity and mortality.

Endoscopy_UCTN_Code_TTT_1AR_2A]

Competing interests: None

Gianfranco Donatelli¹, Serge Derhy², David Fuks³, Hugues Levard³, Bruno Meduri¹, Brice Gayet³

¹ Unité d'Endoscopie Interventionnelle, Ramsay Générale de Santé, Hôpital Privé des Peupliers, Paris, France

² Unité de Radiologie Interventionnelle, Ramsay Générale de Santé, Hôpital Privé des Peupliers, Paris, France

³ Department of Digestive Diseases, Institut Mutualiste Montsouris, Paris, France

References

- 1 Jablonska B, Lampe P. Iatrogenic bile duct injuries: etiology, diagnosis and management. *World J Gastroenterol* 2009; 15: 4097–4104
- 2 Kunda R, Pérez-Miranda M, Will U et al. EUS-guided choledochoduodenostomy for malignant distal biliary obstruction using a lumen-apposing fully covered metal stent after failed ERCP. *Surg Endosc* 2016. DOI: <http://dx.doi.org/10.1007/s00464-016-4845-6>
- 3 Perez-Miranda M, Aleman N, de la Serna Higuera C et al. Magnetic compression anastomosis through EUS-guided choledochoduodenostomy to repair a disconnected bile duct in orthotopic liver transplantation. *Gastrointest Endosc* 2014; 80: 520–521

4 Donatelli G, Vergeau BM, Derhy S et al. Combined endoscopic and radiologic approach for complex bile duct injuries (with video). *Gastrointest Endosc* 2014; 79: 855–864

5 Meduri B, Derhy S, Dhumane P et al. Extra-anatomical intraduodenal endoscopic-radiologic biliary rendezvous for treatment of iatrogenic complete stenosis of the common bile duct. *Endoscopy* 2015; 47 (Suppl. 01): E565–566

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-116818>
Endoscopy 2016; 48: E293–E294
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Gianfranco Donatelli, MD

Unité d'Endoscopie Interventionnelle
Ramsay Générale de Santé
Hôpital Privé des Peupliers
8 Place de l'Abbé G. Hénocque
75013 Paris
France
Fax: +33-01-44165615
donatelligianfranco@gmail.com