Perigastric fluid collection after endoscopic sleeve gastroplasty

Fig. 1 Endoscopic views showing: **a** the technique used to perform endoscopic sleeve gastroplasty (ESG); **b** the appearance after completion of the ESG procedure.

Fig. 2 Computed tomography (CT) scan of the abdomen with intravenous contrast 12 days post-endoscopic sleeve gastroplasty: **a** on coronal view, showing a rim-enhancing 3.3×3.7-cm collection; **b** on axial view, showing a rim-enhancing 3.3×2.9-cm collection.

A 39-year-old woman with a body mass index (BMI) of 32 kg/m² was evaluated for suitability for endoscopic management of her obesity. Because of her interest in a durable weight loss intervention, she elected to undergo an endoscopic sleeve gastroplasty (ESG). The procedural details have been described previously by Sha-

raihia et al. and Lopez Nava et al. [1–3]. In brief, the volume of the stomach is reduced by approximately 70% using a full-thickness endoscopic suturing system (OverStitch; Apollo Endosurgery, Austin, Texas, USA) (Fig. 1).

The patient recovered well in the days immediately after the procedure but pre-

sented with gradual onset vomiting and abdominal pain 12 days after her ESG. An erect abdominal radiograph showed a non-obstructive bowel gas pattern without any evidence of free gas. An abdominal contrast-enhanced computed tomography (CT) scan revealed a rim-enhancing 3.3×2.9×3.7-cm fluid collection in the fundus along the greater curvature of the stomach, suggestive of a perigastric collection, with surrounding stranding of the omental fat (Fig. 2).

She did not have fever or chills, and her vital signs were stable, with a normal temperature and a white blood cell (WBC) count of 11.1×10³/μL. The patient was given a stat dose of intravenous etrapenem and was discharged with oral ciprofloxacin 500 mg twice daily and metronidazole 400 mg three times daily for 7 days. At follow-up 3 weeks later, her symptoms had completely resolved and follow-up imaging revealed no residual collection.

There has only been one documented case of perigastric inflammatory serous fluid collection after ESG (adjacent to the fundus), which resolved with percutaneous drainage and antibiotics [4]. One possible cause of the perigastric fluid collection may be intractable vomiting causing a small leak along the thin-walled fundus. When such a collection is identified, we recommend initial conservative therapy with oral antibiotics if there are no signs or symptoms of sepsis. Depending on the size and location of the collection, percutaneous drainage could be performed.

Endoscopy_UCTN_Code_CPL_1AH_2AJ

Competing interests: Mouen A. Khashab is a consultant for Boston Scientific and Olympus America and has received research support from Cook Medical. Vivek Kumbhari is a consultant for Boston Scientific and Apollo Endosurgery. All other authors have no disclosures

Sindhu Barola¹, Abhishek Agnihotri², Mouen A. Khashab¹, Vivek Kumbhari¹

¹ Department of Medicine and Division of Gastroenterology and Hepatology, The Johns Hopkins Medical Institutions, Baltimore, Maryland, USA

² Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

References

- 1 Sharaiha RZ, Kedia P, Kumta N et al. Initial experience with endoscopic sleeve gastropasty: technical success and reproducibility in the bariatric population. *Endoscopy* 2015; 47: 164–166
- 2 Lopez-Nava G, Galvao M, Bautista-Castaño I et al. Endoscopic sleeve gastropasty with 1-year follow-up: factors predictive of success. *Endosc Int Open* 2016; 4: E222–E227
- 3 Lopez-Nava G, Galvão MP, da Bautista-Castaño I et al. Endoscopic sleeve gastropasty for the treatment of obesity. *Endoscopy* 2015; 47: 449–452
- 4 Abu Dayyeh BK, Acosta A, Camilleri M et al. Endoscopic sleeve gastropasty alters gastric physiology and induces loss of body weight in obese individuals. *Clin Gastroenterol Hepatol*. Epub ahead of print 2015 Dec 31

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-117844>
Endoscopy 2016; 48: E340–E341
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Vivek Kumbhari, MD

Division of Gastroenterology and Hepatology
 Johns Hopkins Medical Institutions
 4940 Eastern Avenue
 AA Building, 3rd floor
 Baltimore
 Maryland 21224
 USA
 Fax: +1-410-550-7861
vkumbhari@gmail.com