Does carbon dioxide insufflation impact adenoma detection rate? A single-center retrospective analysis

Authors

Yaseen B. Perbtani¹, Michael Rivero¹, Jonathan J. Shuster², Joydeep Chakraborty¹, Tony S. Brar¹, Mitali Agarwal¹, Han Zhang¹, Anand Gupte³, Shailendra S. Chauhan³, Christopher E. Forsmark³, Peter V. Draganov³, Dennis Yang³

Institutions

¹ Department of Medicine, University of Florida, Gainesville, Florida, United States

² Department of Health Outcomes and Policy University of Florida, Gainesville, Florida, United States

³ Division of Gastroenterology, University of Florida, Gainesville, Florida, United States

submitted 30. April 2016
accepted after revision
12. September 2016

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-118177>
Published online: 24.11.2016
Endoscopy International Open 2016; 04: E1275–E1279
© Georg Thieme Verlag KG
Stuttgart · New York
E-ISSN 2196-9736

Corresponding author

Dennis Yang, MD
Division of Gastroenterology,
University of Florida
1329 SW 16th Street,
Suite 5251
Gainesville, FL 32608
United States
Phone: +1-352-273-9474
Fax: +1-352-627-9002
Dennis.Yang@medicine.ufl.edu

Background and study aims: Carbon dioxide (CO₂) has been associated with reduced post-procedural pain and improved patient satisfaction when compared to air insufflation (AI). The effect of CO₂ insufflation (CO₂I) on the adenoma detection rate (ADR) remains unclear. The aims of this study are to compare ADR in patients undergoing screening colonoscopy with AI vs. CO₂I and identify predictors of ADR.

Patients and methods: Single-center retrospective cohort study of 2,107 patients undergoing screening colonoscopy at the University of Florida Hospital between November 2011 and June 2015. Patient demographics, procedural parameters, and histology results were retrospectively obtained from a prospectively maintained colonoscopy database. Univariate and multivariate analysis were performed to identify predictors of ADR.

Results: A total of 2107 colonoscopies (644 with AI and 1463 with CO₂I) were analyzed. Overall ADR was 27.8%. There was no significant difference in ADR between AI (27.6%) vs. CO₂I (27.8%) ($P=0.93$). Method of insufflation (AI vs. CO₂I) was not significantly associated with ADR (OR 0.9; 95% CI: 0.7–1.2). Older age (OR: 1.02; 95% CI: 1.001–1.03 per year increase), male gender (OR 1.48; 95% CI: 1.17–1.87), and longer scope withdraw time (OR 1.13; 95% CI: 1.1–1.16 per minute) were associated with a higher ADR. Fellow involvement was negatively associated with ADR (OR 0.60; 95% CI: 0.47–0.77).

Conclusion: ADR was similar between patients who underwent screening colonoscopy with AI vs. CO₂I. While CO₂I has been associated with improved patient comfort and post-procedural recovery time, there is no definitive evidence to suggest that this method of luminal distention enhances ADR.

Introduction

Colorectal cancer (CRC) remains the second leading cause of cancer death in the United States. Colonoscopy has been shown to reduce risk of death from CRC through the prompt identification and removal of premalignant adenomas or early stage cancerous lesions [1]. Indeed, the increased utilization of screening colonoscopy has been associated with a 30% reduction in cancer deaths due to CRC in the past decade [2]. Yet the reported impact of colonoscopy on CRC incidence has been widely variable, presumably due to discrepancies in the quality of the procedure [3].

The adenoma detection rate (ADR), defined as the proportion of screening colonoscopies performed by a provider with the detection of at least 1 histologically confirmed adenoma or adenocarcinoma, has been emphasized as an important quality indicator for colonoscopy, given the supporting evidence on ADR and its impact on CRC [4]. There-

fore, the American Society for Gastrointestinal Endoscopy/American College of Gastroenterology Task Force on Quality in Endoscopy has recently proposed ADR targets of $\geq 30\%$ in men and $\geq 20\%$ in women [5]. Furthermore, in the setting of recent data demonstrating a 3.0% decrease in the risk of interval CRC with each 1.0% increase in ADR [6], it is not surprising that there has been an increased emphasis on identifying and regulating factors that contribute to variability in adenoma detection and thereby improve quality of colonoscopies among practitioners.

Luminal distention during colonoscopy is necessary to allow adequate inspection of the colonic mucosa. Room air insufflation (AI) traditionally has been the most commonly used method for bowel insufflation during colonoscopy. More recently, there has been increasing data supporting the use carbon dioxide (CO₂) as an alternate method for luminal distention. CO₂ is more rapidly absorbed across the intestines when compared

License terms

Table 1 Study population.

Variable	AI (n=644)	CO ₂ I (n=1463)	P value
Age, mean ± SD (years)	58.7 ± 8.9	59.1 ± 8.7	0.32
Gender	289 (44.9)	680 (46.5)	0.50
Male, n (%)			
BMI, mean ± SD	32.2 ± 9.4	31.5 ± 8.8	0.14
ASA score			0.18
I	25 (3.9)	50 (3.4)	
II	286 (44.4)	625 (42.7)	
III	316 (49.1)	719 (49.2)	
IV	17 (2.6)	69 (4.7)	
Fellow involvement, n (%)	320 (49.7)	553 (37.8)	<.001
Bowel preparation grading			0.0016
Excellent	104 (16.2)	187 (12.8)	
Good	386 (59.9)	836 (57.1)	
Fair	111 (17.2)	320 (21.9)	
Poor	43 (6.7)	120 (8.2)	
Cecal intubation rate, n (%)	638 (99.1)	1437 (98.2)	0.14
Total procedure time, mean ± SD (minutes)	20.9 ± 9.9	19.6 ± 9.6	0.0022
Scope withdrawal time, mean ± SD (minutes)	10.2 ± 6.9	9.4 ± 6.4	0.023

AI, air insufflation; CO₂I, carbon dioxide insufflation; SD, standard deviation.

to air and it has been shown in a large meta-analysis to be associated with a reduction in procedure-related pain [7–10]. However, data are limited on the impact of CO₂ insufflation (CO₂I) on ADR. The aim of this study was to (1) compare ADR in patients undergoing screening colonoscopy with CO₂I versus AI and to (2) identify factors influencing ADR in patients undergoing screening colonoscopy in a tertiary-care endoscopy unit.

Patients and methods

Study design and patients

The study was approved by the institutional review board (IRB) of the University of Florida in which a waiver for informed consent was obtained. To be eligible for inclusion in this report, the endoscopic database at the University of Florida Health (UF Health) was retrospectively searched for all patients who had undergone a screening colonoscopy between November 2011 and June 2015. Eligibility was restricted to those with average CRC risk. Patients were excluded if they had a personal or first-degree relative family history of CRC, history of colon polyps, inflammatory bowel disease, gastrointestinal bleeding, prior history of partial colon resection, and/or an aborted procedure (i.e. due to unsatisfactory bowel preparation, patient intolerance, procedure-related factors) based on information from the endoscopic database and/or electronic chart record. Informed procedural consents were obtained from all patients.

Colonoscopy procedure

All patients underwent screening colonoscopies by 1 of the 24 experienced board-certified gastroenterologists or by gastroenterology fellows (first- to third-year of fellowship training) under direct supervision of a staff attending. The bowel preparation agent used was predominantly 4L of polyethylene glycol solution. Bowel preparation quality was rated as excellent, good, fair, or poor based on the Aronchick scale [11]. All colonoscopies were performed under provider-anesthesiologist-administered conscious sedation (fentanyl and midazolam) or propofol sedation. Cecal intubation was documented by the endoscopist using landmark descriptions (i.e. identification of the ileocecal valve and/or

appendiceal orifice). Total procedure time (defined as amount of time from scope insertion to scope removal from the patient) and withdrawal time (defined as the amount of time spent examining the mucosa as the colonoscope is withdrawn) were recorded prospectively and documented in the electronic report by the nurses. Air insufflation was used for luminal distention in all screening colonoscopies prior to January 21st, 2013. Following that date, routine use of CO₂ for insufflation was adopted universally for all endoscopic procedures in our institution.

Data collection

Demographic information and histopathology reports were obtained from chart review. Patient demographics included age, sex, American Society of Anesthesiology (ASA) physical status grade, and body mass index (BMI). Adenomas were classified as tubular, tubulovillous, villous or carcinoma based on the Vienna criteria [12]. Dysplasia was defined as either low-grade or high-grade. Colonoscopy-related data obtained from the prospectively maintained report generating database included: type of sedation, quality of bowel preparation, fellow involvement, cecal intubation, total procedure time, withdrawal time, and number of polyps removed. Endoscopic adverse events (AEs) were defined based on previously established criteria by the American Society of Gastrointestinal Endoscopy (ASGE) [13]. AEs were determined by reviewing the colonoscopy report and the immediate post-procedural note.

Study outcomes

The primary aim of this study was to compare the ADR in patients undergoing screening colonoscopy with AI versus CO₂I. ADR was defined as the proportion of colonoscopies in which at least 1 histologically confirmed adenoma was detected. A secondary aim was to identify factors associated with ADR in our cohort.

Statistical methods

Baseline characteristics for Table 1 between the 2 cohorts AI and CO₂I were compared by (a) the Student's *t*-test with the Satterthwaite correction for unequal variances for quantitative variables (age, BMI, total procedure time, and scope withdrawal time); (b) Pearson's chi-square for binary variables (gender, fel-

Pathology	AI (n=644)	CO ₂ I (n=1463)	OR (95% CI)	P value
Tubular adenoma, n (%)	156 (24.22)	379 (25.91)	0.91 (0.74 – 1.13)	0.41
Tubulovillous adenoma, n (%)	23 (3.57)	50 (3.42)	1.05 (0.63 – 1.73)	0.86
Villous adenoma, n (%)	5 (0.78)	4 (0.27)	2.85 (0.76 – 10.66)	0.10
Adenocarcinoma, n (%)	0	5 (0.34)	–	0.14

AI, air insufflation; CO₂I, carbon dioxide insufflation.

Table 2 Histologic classification of adenomatous polyps.

Clinical variable	Univariate analysis		Multivariate Analysis	
	OR (95% CI)	P value	OR (95% CI)	P value
Age (years)	1.02 (1.004 – 1.026)	.0067	1.02 (1.001 – 1.028)	0.034
Gender (male vs. female)	1.56 (1.29 – 1.89)	<.001	1.48 (1.17 – 1.87)	0.001
BMI (kg/M ²)	1.00 (0.99 – 1.01)	0.87	1.01 (0.99 – 1.02)	0.51
ASA score 1 vs. 2,3 and/or 4	1.56 (0.88 – 2.77)	0.13	1.61 (0.84 – 3.12)	0.15
Type of sedation (conscious sedation vs. propofol)	0.99 (0.81 – 1.22)	0.93	1.02 (0.71 – 1.45)	0.93
Fellow involvement (yes vs. no)	1.00 (0.82 – 1.21)	0.98	0.60 (0.47 – 0.77)	<.001
Quality of bowel preparation (excellent vs good/fair/poor)	1.43 (1.06 – 1.92)	0.018	1.37 (0.96 – 1.96)	0.087
Cecal intubation (yes vs. no)	2.10 (0.80 – 5.47)	0.13	1.18 (0.23 – 5.94)	0.84
Total procedure time (Min)	1.06 (1.05 – 1.07)	<.001	1.01 (0.99 – 1.02)	0.55
Scope withdrawal time (Min)	1.13 (1.11 – 1.15)	<.001	1.13 (1.10 – 1.16)	<.001
Type of insufflation (AI vs. CO ₂ I)	0.99 (0.81 – 1.22)	0.93	0.90 (0.69 – 1.17)	0.41

AI, air insufflation; CO₂I, carbon dioxide insufflation.

Table 3 Factors associated with ADR.

low-involvement, cecal intubation); and (c) the Wilcoxon test for ordinal variables (ASA score and Bowel preparation grading).

Univariate analysis for **Table 2** and **Table 3** was conducted by univariate and multiple logistic regression. The odds ratios for the quantitative independent variables in **Table 3** reflect the ratio of odds, for 2 subjects with 1 with a value 1 unit higher than the other, but otherwise equivalent on other covariates in the model, if any, higher value to lower value. The multivariate model estimates the odds ratio (and compares it to the null value of 1.00) adjusting for all other variables in the model. Significance in the multiple regression model means that the variable has independent significant prognostic value that cannot be accounted for by the other variables in the model.

All P values are 2-sided. SAS (Statistical Analysis Systems) version 9.4 was used in all of the analyses.

Results

Study population

A total of 2107 screening colonoscopies were performed among the eligible patients during the study period. Of these colonoscopies, 644 (30.6%) were performed with AI compared to 1463 (69.4%) with CO₂I (**Table 1**). There were no significant differences in age, gender, BMI or ASA score between patients undergoing colonoscopy with AI vs. CO₂I. The cecal intubation rate was also similar between the 2 groups (99.1% in the AI group vs. 98.2% in the CO₂I group; *P*=0.14). The total procedure time and scope withdrawal time were slightly longer in patients undergoing colonoscopy with AI vs. CO₂I. Fellows were more commonly involved in colonoscopies with AI (49.7%) vs. CO₂I (37.8%) (*P*<0.001). The average scope withdrawal time was longer when a fellow was involved/present during the procedure (11.5±7.5 min vs. 8.3±5.5 min; *P*<0.001). Cumulatively,

the quality of the bowel preparation was rated better in patients undergoing colonoscopies with AI vs. CO₂I (*P*=0.0016).

Adenoma detection rate

Overall, a total of 622 adenomatous lesions were detected in this study. The cumulative ADR in this study was 27.8%. There was no statistically significant difference in the ADR in patients undergoing colonoscopy with AI (178/644; 27.6%) vs. CO₂I (407/1463; 27.8%) (*P*=0.93). The histologic classification of the adenomatous polyps detected was also similar in both groups (**Table 2**). Overall, tubular adenoma was the most common histopathology reported for both groups (*P*=0.41). There were a total of 5 adenocarcinomas diagnosed on histopathology, all from patients who underwent CO₂I (*P*=0.14). Only a small number of adenomas detected during colonoscopies with AI (4.3%) and CO₂I (3.9%) revealed high-grade dysplasia (OR 0.89; 95% CI: 0.38 – 2.09).

Factors associated with ADR

Univariate and multiple logistic regression analyses were performed to identify factors associated with ADR (**Table 3**). The variables included were age, gender, BMI, fellow involvement (yes vs. no), cecal intubation (yes vs. no), scope withdrawal time, total procedure time, type of sedation (conscious sedation vs. propofol), quality of bowel preparation and method of bowel insufflation (AI vs CO₂I). Patient characteristics, including older age and male gender, were associated with a higher ADR in both univariate and multivariate analysis. While both scope withdrawal time and total procedure time correlated positively with ADR on univariate analysis, only scope withdrawal time was found to positively impact ADR on multivariate analysis. Neither type of sedation nor method of bowel insufflation (AI vs. CO₂I) had a significant association with ADR. Quality of bowel preparation was also not significantly associated with ADR (OR 1.37; 95% CI: 0.96 – 1.96). On the other hand, fellow involvement during the

colonoscopy was the only covariate that was negatively associated with ADR on multivariate analysis.

Adverse events

There were no procedural or sedation-related AEs reported in the prospective colonoscopy database or in the immediate post-operative note on chart review. Overall, 28 procedures (1.3%) were aborted due to inadequate bowel preparation and/or tortuous colon as indicated on the colonoscopy report. There were a total of 6 cases terminated prematurely due to patient discomfort. Out of these, 4 cases were done with AI vs 2 with CO₂I ($P=0.07$).

Discussion

Colonoscopy is considered the gold standard for CRC screening. The effectiveness of this strategy at reducing the morbidity and mortality associated with CRC is dependent on optimal detection and resection of premalignant or early stage cancerous lesions. Hence, ADR is currently regarded as the most important measure of quality in colonoscopy. In this study, there was no difference in ADR in patients undergoing screening colonoscopy with AI versus those with CO₂I after adjusting for patient and procedural variables.

AI is the most commonly utilized method for colonic insufflation during colonoscopy. More recently, alternate methods of luminal distention, such as water-assisted colonoscopy and CO₂I, have been sought as to reduce post-procedural patient pain and bloating associated with AI. Yet the current literature on methods of bowel distention and its effect on ADR is scarce. In a recent Cochrane review, Hafner et al. compared technical quality and screening efficacy between patients undergoing AI versus water infusion during colonoscopy [14]. Their analysis, which included 16 randomized controlled trials consisting of 2933 colonoscopies, showed a slight improvement in ADR with water infusion vs. AI (risk ratio 1.16; 95% CI: 1.04–1.30, $P=0.007$). It has been speculated that this increase in ADR may be secondary to additional bowel lavage with water infusion, with the drawback of significantly longer insertion time [15]. Conversely, the previous literature on the impact of CO₂I on ADR has been limited to a single abstract submission by Mills and colleagues [16]. The preliminary results from their retrospective review suggested a higher ADR in patients with CO₂I vs. AI for colonoscopy (OR 1.36; 95% CI: 1.01–1.85). However, information on whether and how these findings were adjusted for other potential confounding factors was unavailable, which makes the interpretation of their results difficult. In our study of 2107 screening colonoscopies, there was no significant difference in ADR between patients undergoing the procedure with AI vs. CO₂I. Furthermore, method of bowel insufflation (AI vs. CO₂I) was not found to be significantly correlated with ADR (OR 1.12; 95% CI: 0.87–1.45) after adjusting for patient characteristics (age, gender, BMI) and procedural variations (fellow involvement, cecal intubation rate, type of sedation, quality of bowel preparation, total procedure and scope withdrawal time). Nonetheless, while our current data did not demonstrate a potential advantage of CO₂I over AI on ADR, several other studies have alluded to the benefits of CO₂I in terms of patient comfort and satisfaction [10,17,18]. Congruent with prior findings, in this study we demonstrated that a slightly higher number of cases cancelled due to patient discomfort when the procedure was performed with AI vs. CO₂I; albeit this difference

did not quite reach statistical significance ($P=0.07$). In general, patient acceptability and tolerability would intuitively seem to improve the efficacy of the procedure. Whether the overall widely reported positive perception and attitude towards CO₂I during colonoscopy actually translates to higher compliance and thereby effective screening remains to be determined.

Differences in patient demographics and procedure-related factors have varying effects on the ADR. Both older age and male gender have been shown to be independently associated with a higher ADR on previous studies [19,20]. Our results are in line with these prior reports as both increasing age and male gender positively correlated with ADR. Similarly, procedural process measures, particularly scope withdrawal time, have been linked with ADR and hence the quality of the examination. In a landmark study by Barclay and colleagues, mean withdrawal times of 6 minutes or more were found to have higher rates of detection of neoplasia (28.3% vs. 11.8%, $P<0.001$) [21]. Several other studies have further confirmed a linear correlation between withdrawal time and ADR [22,23]. In our study, the mean withdrawal time was more than 6 minutes in patients undergoing colonoscopy irrespective of method of insufflation. Congruent with prior studies, this study demonstrated that longer scope withdrawal time was associated with a higher ADR (OR 1.13; 95% CI: 1.1–1.16). Our findings further corroborate the importance of withdrawal time as a surrogate marker for the thoroughness and quality of the examination for the detection of neoplastic lesions during colonoscopy.

In this study, fellow involvement was associated with a decrease in ADR in multivariate analysis (Table 3). The current available data on the effect of fellow involvement in colonoscopy on ADR is conflicting, with some studies supporting a positive correlation [24], whereas others suggesting no effect or a lower ADR [25]. This discrepancy among studies may be in part associated with differences in the level of training of fellows included as well as with the degree of supervision provided by the staff endoscopist at the time of the procedure. In this study, fellow involvement was negatively associated with ADR even though the average withdrawal time was significantly longer when a fellow was present. Future prospective studies are needed to clarify the association between fellow involvement and ADR. This is not only imperative in order to maintain quality of care but also to identify benchmarks during colonoscopy training.

Our findings should be interpreted in light of the strengths and limitations of the study. The main strengths of our study are the comprehensive and detailed assessment of ADR in 2107 colonoscopies performed in our institution. Multiple established quality indicators were all collected prospectively over a 4-year period and included in our analysis. Our findings on the effect of patient demographics (age, gender) and procedure parameters (scope withdrawal time) on ADR are in line with those previously reported and allude to the validity of the study. Furthermore, this is the first study evaluating the impact of CO₂I vs. AI on ADR. Our findings suggest that CO₂I did not significantly correlate with ADR after adjusting for patient and procedural covariates. With the increasing number of reports supporting the use of CO₂I for endoscopic procedures in regards to patient comfort and post-procedural recovery, further prospective studies evaluating its effect on ADR and thereby the effectiveness of the colonoscopy are needed. We also recognize the limitations of our study. First, the study was performed in a single tertiary care center and results may not be generalizable to all ambulatory endoscopic units. Furthermore, this was a retrospective study with its inher-

ent limitations, including baseline differences in rate of fellow involvement, quality of bowel preparation, and procedural time between patients undergoing colonoscopy with AI vs. CO₂I. Nonetheless, the effect of these variables was adjusted by performing a multivariate analysis to specifically determine whether any of the confounding variables contributed specifically to the ADR. Furthermore, data on the type of colonoscopes (i.e. standard definition vs. high-definition) used in this study was not readily available in the final analysis. Our ongoing data extraction suggests that close to 90% of all cases in both groups were performed with high-definition colonoscopes (data not shown). While we recognize that differences in the type of colonoscopes between the 2 groups may affect the interpretability of our findings, its impact on polyp or adenoma detection rate remains debatable based on the conflicting literature [26–28]. Lastly, while the adequacy of bowel cleansing has been clearly linked to ADR, the quality of bowel preparation was not significantly associated with ADR in our study (OR 1.4; 95% CI: 0.96–1.96). This difference could potentially be explained by the bowel cleansing grading used in this cohort (Aronchick scale), which was specifically designed and validated to compare the efficacy of purgatives rather than outcomes such as ADR.

Conclusion

In conclusion, this study demonstrates that the detection of adenomatous polyps in preventive colonoscopy was not significantly increased by the use of CO₂I compared to AI. While the implementation of CO₂I has been clearly associated with improved patient comfort and post-procedural recovery time, there is no definitive evidence to suggest that this method of luminal distention enhances ADR. Future prospective trials are needed to compare the effect of different methods of luminal distention on ADR and on the overall effectiveness of screening colonoscopy.

Competing interests: None

Acknowledgements

Research reported in this publication was partly supported by the National Center For Advancing Translational Sciences of the National Institutes of Health under Award Number UL1TR001427. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

- 1 Zauberg AG, Winawer SJ, O'Brien MJ et al. Colonoscopic polypectomy and long-term prevention of colorectal-cancer deaths. *N Engl J Med* 2012; 366: 687–696
- 2 Siegel R, Desantis C, Jemal A. Colorectal cancer statistics, 2014. *CA Cancer J Clin* 2014; 64: 104–117
- 3 Pohl H, Robertson DJ. Colorectal cancers detected after colonoscopy frequently result from missed lesions. *Clin Gastroenterol Hepatol* 2010; 8: 858–864
- 4 Kaminski MF, Regula J, Kraszewska E et al. Quality indicators for colonoscopy and the risk of interval cancer. *N Engl J Med* 2010; 362: 1795–1803
- 5 Rex DK, Schoenfeld PS, Cohen J et al. Quality indicators for colonoscopy. *Gastrointest Endosc* 2015; 81: 31–53
- 6 Corley DA, Jensen CD, Marks AR et al. Adenoma detection rate and risk of colorectal cancer and death. *N Engl J Med* 2014; 370: 1298–1306
- 7 Wu J, Hu B. The role of carbon dioxide insufflation in colonoscopy: a systematic review and meta-analysis. *Endoscopy* 2012; 44: 128–136
- 8 Brandt LJ, Boley SJ, Sammartano R. Carbon dioxide and room air insufflation of the colon. Effects on colonic blood flow and intraluminal pressure in the dog. *Gastrointest Endosc* 1986; 32: 324–329
- 9 Yasumasa K, Nakajima K, Endo S et al. Carbon dioxide insufflation attenuates parietal blood flow obstruction in distended colon: potential advantages of carbon dioxide insufflated colonoscopy. *Surg Endosc* 2006; 20: 587–594
- 10 Sumanac K, Zealley I, Fox BM et al. Minimizing postcolonoscopy abdominal pain by using CO(2) insufflation: a prospective, randomized, double blind, controlled trial evaluating a new commercially available CO(2) delivery system. *Gastrointest Endosc* 2002; 56: 190–194
- 11 Aronchick CA, Lipshutz WH, Wright SH et al. A novel tableted purgative for colonoscopic preparation: efficacy and safety comparisons with Colyte and Fleet Phospho-Soda. *Gastrointest Endosc* 2000; 52: 346–352
- 12 Schlemper RJ, Riddell RH, Kato Y et al. The Vienna classification of gastrointestinal epithelial neoplasia. *Gut* 2000; 47: 251–255
- 13 Cotton PB, Eisen GM, Aabakken L et al. A lexicon for endoscopic adverse events: report of an ASGE workshop. *Gastrointest Endosc* 2010; 71: 446–454
- 14 Hafner S, Zolk K, Radaelli F et al. Water infusion versus air insufflation for colonoscopy. *Cochrane Database Syst Rev* 2015; 5: Cd009863
- 15 Leung FW, Amato A, Ell C et al. Water-aided colonoscopy: a systematic review. *Gastrointest Endosc* 2012; 76: 657–666
- 16 Mills CD, Swaine A, Mccamley C et al. Su1686 The impact of carbon dioxide insufflation on colonic polyp and adenoma detection rate. *Gastrointestinal Endoscopy* 2015: 81
- 17 Riss S, Akan B, Mikola B et al. CO2 insufflation during colonoscopy decreases post-interventional pain in deeply sedated patients: a randomized controlled trial. *Wien Klin Wochenschr* 2009; 121: 464–468
- 18 Wang WL, Wu ZH, Sun Q et al. Meta-analysis: the use of carbon dioxide insufflation vs. room air insufflation for gastrointestinal endoscopy. *Aliment Pharmacol Ther* 2012; 35: 1145–1154
- 19 Schramm C, Mbaya N, Franklin J et al. Patient- and procedure-related factors affecting proximal and distal detection rates for polyps and adenomas: results from 1603 screening colonoscopies. *Int J Colorectal Dis* 2015; 30: 1715–1722
- 20 Cai B, Liu Z, Xu Y et al. Adenoma detection rate in 41,010 patients from Southwest China. *Oncol Lett* 2015; 9: 2073–2077
- 21 Barclay RL, Vicari JJ, Doughty AS et al. Colonoscopic withdrawal times and adenoma detection during screening colonoscopy. *N Engl J Med* 2006; 355: 2533–2541
- 22 Jover R, Zapater P, Polania E et al. Modifiable endoscopic factors that influence the adenoma detection rate in colorectal cancer screening colonoscopies. *Gastrointest Endosc* 2013; 77: 381–389.e381
- 23 Butterly L, Robinson CM, Anderson JC et al. Serrated and adenomatous polyp detection increases with longer withdrawal time: results from the New Hampshire Colonoscopy Registry. *Am J Gastroenterol* 2014; 109: 417–426
- 24 Buchner AM, Shahid MW, Heckman MG et al. Trainee participation is associated with increased small adenoma detection. *Gastrointest Endosc* 2011; 73: 1223–1231
- 25 Leffler DA, Kheraj R, Bhansali A et al. Adenoma detection rates vary minimally with time of day and case rank: a prospective study of 2139 first screening colonoscopies. *Gastrointest Endosc* 2012; 75: 554–560
- 26 Burke CA, Choure AG, Sanaka MR et al. A comparison of high-definition versus conventional colonoscopes for polyp detection. *Dig Dis Sci* 2010; 55: 1716–1720
- 27 Tribonias G, Theodoropoulou A, Konstantinidis K et al. Comparison of standard vs high-definition, wide-angle colonoscopy for polyp detection: a randomized controlled trial. *Colorectal Dis* 2010; Oct 12: e260–266
- 28 East JE, Stavrindis M, Thomas-Gibson S et al. A comparative study of standard vs. high definition colonoscopy for adenoma and hyperplastic polyp detection with optimized withdrawal technique. *Aliment Pharmacol Ther* 2008; 28: 768–776