Balloon enteroscopy-assisted biliary drainage using a diathermic dilator followed by placement of a novel ultra-slim metallic stent

Endoscopic management of malignant biliary stricture after hepaticojejunostomy is challenging. Although balloon enteroscopy-assisted endoscopic retrograde cholangiopancreatography (ERCP) has been developed, dedicated devices are still insufficient. We present a case of successful enteroscopic dilation using a wire-guided diathermic dilator [1–5], and placement of a novel, ultra-slim, uncovered, self-expandable, metallic stent (USEMS) for the treatment of a severe malignant stricture of the perihilar bile duct after hepaticojejunostomy.

A 52-year-old man with obstructive jaundice was referred to our department. He had undergone pancreaticoduodenectomy with hepaticojejunostomy for pancreatic head cancer 27 months earlier. He had received 6 months of chemotherapy for multiple liver metastases, but this treatment was suspended because of multiple liver abscesses, which developed 1 month before we saw him.

Computed tomography revealed intrahepatic bile duct dilation, perihilar biliary mass, and ascites. Double-balloon enteroscopy (EI-530B; Fujifilm Corp., Tokyo, Japan) was selected for biliary drainage. Double-balloon enteroscopy showed jejunal invasion of recurrent tumor involving the hepaticojejunal anastomosis (► Fig. 1 a, b). A tapered ERCP catheter and a 0.025-inch guidewire were passed through the stricture of the hepaticojejunal anastomosis (► Fig. 1 b). ERCP showed perihilar biliary stricture (► Fig. 2 a, ► Video 1), which could not be passed by the tapered ERCP catheter (► Video 1).

The severe stricture was dilated successfully with a wire-guided diathermic dilator (Cysto-Gastro-Set, 6Fr, length 180cm; Endo-Flex, Voerde, Germany) (► Fig. 2 b, ► Video 1). Total dilation time was 3 seconds for the hepaticojejunal anastomosis and 13 seconds for the

► Fig. 1 Endoscopic images. a Jejunal invasion. b Edematous mucosa around the hepaticojejunal anastomosis.

perihilar biliary stricture, respectively. A slim USEMS (Zilver 635, 6.0Fr delivery system, 10mm diameter, 200cm long; Cook Japan, Tokyo, Japan) failed to pass the stricture. However, a novel ultra-slim USEMS (BileRush Selective, 5.7Fr delivery system, 10mm diameter, 185cm long; Piolax Medical Devices, Kanagawa, Japan) was successfully deployed, followed by placement of an additional USEMS, in a stent-in-stent fashion, with-

out any complications (► Fig. 3, ► Fig. 4, ► Video 1).

In summary, using balloon enteroscopy, a wire-guided diathermic dilator and novel ultra-slim USEMS were useful in facilitating dilation and stenting for the treatment of severe malignant anastomotic and biliary strictures in a patient with altered anatomy.

Endoscopy_UCTN_Code_TTT_1AS_2AD

► VIDEO 1

► Video 1: Wire-guided diathermic dilation after failed dilation using a tapered endoscopic retrograde cholangiopancreatography catheter, and deployment of novel, ultra-slim, uncovered, self-expandable, metallic stents.

► **Fig. 2** Radiographic images. **a** Perihilar biliary stricture. **b** Wire-guided diathermic dilator passing the perihilar bile duct stricture (inset: 6 Fr wire-guided diathermic dilator insertion under endoscopic guidance).

► **Fig. 4** Radiograph showing deployed novel ultra-slim uncovered self-expandable metallic stent (inset: enteroscopic view).

► **Fig. 3** Comparison of a novel, ultra-slim, uncovered, self-expandable, metallic stent (USEMS; BileRush Selective, 5.7 Fr delivery system, 10 mm diameter, 185 cm long; PIOLAX Medical Devices, Kanagawa, Japan) (above), and a slim USEMS (Zilver 635, 6.0 Fr delivery system, 10 mm diameter, 200 cm long; Cook Japan, Tokyo, Japan) (below).

Competing interests

Dr. Kawakami has collaborated with Piolax Medical Devices on the development of the BileRush Selective, and is a consultant and gives lectures for the company.

The Authors

Hiroshi Kawakami^{1,2}, Yoshimasa Kubota^{1,2}, Shinya Ashizuka^{2,3}, Masaru Haraguchi^{2,3}, Ichiro Sannomiya^{2,3}

- 1 Department of Gastroenterology and Hepatology, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan
- 2 Center for Digestive Disease, University of Miyazaki Hospital, Miyazaki, Japan
- 3 Circulatory and Body Fluid Regulation, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan

Corresponding author

Hiroshi Kawakami, MD, PhD

Department of Gastroenterology and Hepatology, Faculty of Medicine, University of Miyazaki, Center for Digestive Disease, University of Miyazaki Hospital, 5200, Kihara, Kiyotake Miyazaki 889-1692, Japan
Fax: +81-985-859802
hiropon@med.miyazaki-u.ac.jp

References

- [1] Kawakami H, Kuwatani M, Eto K et al. Resolution of a refractory severe biliary stricture using a diathermic sheath. *Endoscopy* 2012; 44 (Suppl. 02): E119 – 120
- [2] Kawakami H, Kuwatani M, Kawakubo K et al. Transpapillary dilation of refractory severe biliary stricture or main pancreatic duct by using a wire-guided diathermic dilator (with video). *Gastrointest Endosc* 2014; 79: 338 – 343
- [3] Kawakami H, Kuwatani M, Sakamoto N. Endoscopic ultrasound-guided antegrade diathermic dilation followed by self-expandable metallic stent placement for anastomotic stricture after hepaticojejunostomy (with video). *Dig Endosc* 2014; 26: 121 – 122
- [4] Kawakami H, Kuwatani M, Kawakubo K et al. Endoscopic ultrasound-guided antegrade diathermic dilation followed by self-expandable metal stent placement for malignant distal biliary stricture. *Endoscopy* 2014; 46 (Suppl. 01): E328 – 329
- [5] Kawakami H, Abo D, Kawakubo K et al. Rendezvous biliary recanalization combining percutaneous and endoscopic techniques using a diathermic dilator for bile duct obstruction. *Endoscopy* 2014; 46 (Suppl. 01): E460 – 461

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-124499>
Endoscopy 2017; 49: E78–E80
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X