Submucosal tunneling endoscopic resection (STER) with full-thickness muscle excision for a recurrent para-aortic esophageal leiomyoma after surgery

► **Fig. 1** Endoscopic view showing a recurrent leiomyoma in the mid-esophagus.

► **Fig. 2** Endoscopic ultrasound (EUS) showing a huge para-aortic leiomyoma developing from the muscularis propria layer.

We report the case of a 49-year-old woman with a recurrent esophageal leiomyoma following two surgical resections. The surgical procedures had been performed 23 and 21 years previously. Follow-up showed lesion recurrence in the mid esophagus (► **Fig. 1** and ► **Fig. 2**). Submucosal endoscopic tunneling resection (STER) was preferred to surgery because of the previous interventions. A single shot of 2g ceftriaxone was administered intravenously prior to the procedure. Submucosal injection, mucosal incision, and tunnel creation was started 5cm above the lesion. Enucleation was performed using a DualKnife J (Olympus, Tokyo, Japan). The lesion was close to the aorta, so the final dissection was carried out by synchronizing with aortic movements. Full-thickness muscle resection was required to achieve en bloc resection being careful to preserve the esophageal adventitia (► **Video 1**). The leiomyoma was grasped with a 30-mm snare and was easily removed (► **Fig. 3**). Six standard clips (QuickClip Pro; Olympus) were deployed to close the mucosal incision.

A computed tomography (CT) scan with swallow study was performed on post-operative day 1; oral diet was restarted on day 2. Histological examination did not show any malignancy.

Tumors originating from the muscularis propria require surgery in most cases [1]. STER is a novel approach for the treatment of subepithelial tumors of the gastrointestinal tract. The risk of perforation may reach up to 15% [2]. If the tumor develops from the muscularis propria, preservation of the serosal layer is difficult; circumferential incision of the serosa is therefore often required to

complete en bloc resection [3]. Even though a large muscular defect exists, mediastinitis does not occur if the mucosal continuity is maintained [4]. Recurrent esophageal leiomyoma has been anecdotally reported and is usually related to incomplete resection or enlargement of a previously undetected nodule [5]. Here, we report the first STER treatment for recurrent esophageal

► **VIDEO 1**

 A screenshot from a video showing a peroral endoscopic view. A large, dark, lobulated mass is visible through a mucosal incision. Surgical instruments are used to manipulate the mass. A QR code is located to the right of the video frame.

► **Video 1:** Peroral endoscopic tunneling dissection of a huge recurrent para-aortic esophageal leiomyoma. Muscularis propria dissection and full-thickness resection was necessary to mobilize the lesion with preservation of the esophageal adventitia.

► **Fig. 3** Endoscopic view showing the 15×30-mm leiomyoma being grasped by a 30-mm snare.

leiomyoma with full-thickness muscle resection, which was required because of involvement of the deep muscularis propria and fibrosis deriving from previous surgery.

Endoscopy_UCTN_Code_TTT_1AO_2AG

Competing interests

None

The Authors

Gianfranco Donatelli¹, David Fuks², Guillaume Pourcher², Isabelle Dumontier¹, Fabrizio Cereatti¹, Thierry Perniceni², Brice Gayet²

- 1 Unité d'Endoscopie Interventionnelle, Ramsay Générale de Santé, Hôpital Privé des Peupliers, Paris, France
- 2 Department of Digestive Surgery, Institut Mutualiste Monsouris, Université Paris Descartes, Paris, France

Corresponding author

Gianfranco Donatelli, MD

Unité d'Endoscopie Interventionnelle, Ramsay Générale de Santé, Hôpital Privé des Peupliers, 8 Place de l'Abbé G. Hénocque, 75013 Paris, France
Fax: + 33-1-44165615
donatelligianfranco@gmail.com

References

- [1] Eleftheriadis N, Inoue H, Ikeda H et al. Submucosal tunnel endoscopy: Peroral endoscopic myotomy and peroral endoscopic tumor resection. *World J Gastrointest Endosc* 2016; 8: 86 – 103
- [2] Chiu PW, Inoue H, Rösch T. From Poem to Poet: Applications and perspectives for submucosal tunnel endoscopy. *Endoscopy* 2016; 48: 1134 – 1142
- [3] Xu MD, Cai MY, Zhou PH et al. Submucosal tunneling endoscopic resection: a new technique for treating upper GI submucosal tumors originating from the muscularis propria layer (with videos). *Gastrointest Endosc* 2012; 75: 195 – 199
- [4] Inoue H, Ikeda H, Hosoya T et al. Submucosal endoscopic tumor resection for subepithelial tumors in the esophagus and cardia. *Endoscopy* 2012; 44: 225 – 230
- [5] Standerfer RJ, Paneth M. Recurrent leiomyoma of the oesophagus. *Thorax* 1982; 37: 478 – 479

Bibliography

DOI <http://dx.doi.org/10.1055/s-0043-100212>
Endoscopy 2017; 49: E86–E87
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X