SpyGlass DS-guided conversion of transmural pancreaticogastrostomy drainage to transpapillary drainage by rendezvous via a lumen-apposing metal stent

Recently, we reported a successful endoscopic ultrasound-guided pancreatic duct (PD) drainage using a lumen-apposing metal stent (LAMS) plus a pigtail stent in a 44-year-old man with chronic pancreatitis and PD stricture [1]. The patient's course was satisfactory and 5 months later conversion of the transmural drainage to transpapillary drainage was planned.

The mature fistula, derived from the pancreaticogastrostomy using the LAMS, was used as a port to facilitate access of a SpyScope (Boston Scientific, Natick, Massachusetts, USA) through a therapeutic upper scope. First, a "buried stent effect" was found in the stomach (► Fig. 1). The LAMS was unearthed using grasping forceps. Then the cholangiopancreatoscope was positioned at the inner end of the stent, and a 0.035-inch guidewire was advanced in an antegrade direction towards the papilla (► Fig. 2 and ► Fig. 3). Contrast filling using an endoscopic retrograde cholangiopancreatography cannula revealed a severe PD stricture, and allowed advancement of the guidewire to inside the duodenal lumen (► Fig. 4). Curiously, no contrast filling of the small-bowel lumen was observed, and the antegrade approach was aborted and changed to a rendezvous method. The scope was exchanged for a duodenoscope, and good positioning of the guidewire in the duodenal lumen was confirmed (► Fig. 5). This allowed retrograde pancreatic cannulation, and dilation of the stricture in the pancreatic head using a 6-mm biliary balloon. Finally, a double-pigtail plastic stent (7 Fr × 10 cm; Boston Scientific) was inserted into the major papilla and inside the PD, and its inner end was located in the gastric body, through the LAMS (► Fig. 6; ► Video 1). No adverse events were reported.

There is a lack of knowledge about the behavior of transmural drainage in long-term follow-up [2], but it is known that transpapillary drainage more closely re-

► **Fig. 1** SpyGlass DS-guided conversion of transmural drainage to transpapillary drainage in a 44-year-old man with chronic pancreatitis and pancreatic duct stricture. The fistula tract orifice in the gastric body can be seen but the lumen-apposing metal stent (AXIOS, 6 × 8 mm) is "buried" by hyperplasia of the gastric mucosa.

► **Fig. 2** Image from a peroral single-operator cholangiopancreatoscope through a therapeutic upper endoscope. The inner end of the lumen-apposing stent is impacted on the pancreatic duct wall.

plicates normal physiology because pancreatic juice drains directly into the duodenum, an alkaline environment, and enterokinase converts trypsinogen into its active form, trypsin, resulting in the activation of pancreatic digestive enzymes [3].

Endoscopy_UCTN_Code_TTT_1AR_2AI

Competing interests

None

VIDEO 1

► Video 1: Pancreaticogastrostomy transmural drainage converted to transpapillary drainage despite severe pancreatic duct stricture: a rendezvous procedure using intraductal cholangioscopy (SpyGlass DS) and direct visualization via a lumen-apposing metal stent.

► **Fig. 3** SpyGlass DS-guided conversion of transmural pancreatic drainage to transpapillary drainage: a 0.035-inch guidewire is advanced, through the cholangiopancreatoscope, in an antegrade direction towards the papilla.

► **Fig. 4** Fluoroscopic view: an antero-grade ductogram. An endoscopic retrograde cholangiopancreatography catheter is inserted through the internal ostomy (pancreaticogastrostomy), and a guidewire is advanced until it reaches the duodenal lumen.

► **Fig. 5** After scope exchange: endoscopic view using a duodenoscope. Rendezvous image of the guidewire passing out of the papilla and being grasped by forceps.

► **Fig. 6** Transpapillary double-pigtail plastic stent inserted retrogradely through the pancreatic head stricture.

The Authors

Joan B. Gornals¹, Claudia Consiglieri¹, Susana Redondo²

- 1 Endoscopy Unit, Department of Digestive Diseases, Hospital Universitari de Bellvitge-IDIBELL, Barcelona, Catalonia, Spain
- 2 Department of Anaesthesiology, Hospital Universitari de Bellvitge-IDIBELL Catalonia, Spain

Corresponding author

Joan B. Gornals, MD, PhD

Endoscopy Unit, Department of Digestive Diseases, Hospital Universitari de Bellvitge-IDIBELL (Bellvitge Biomedical Research Institute), Feixa Llarga s/n, 08907 L'Hospitalet de Llobregat, Barcelona, Catalonia, Spain
Fax: +34-93-2607681
jgornals@bellvitgehospital.cat

References

- [1] Gornals JB, Consiglieri C, Vida F et al. Endoscopic ultrasound-guided pancreaticogastrostomy using a lumen-apposing metal stent plus a double-pigtail plastic stent. *Endoscopy* 2016; 48: 276–277
- [2] Chapman CG, Waxman I, Siddiqui UD. Endoscopic ultrasound (EUS)-guided pancreatic duct drainage: the basics of when and how to perform EUS-guided pancreatic duct interventions. *Clin Endosc* 2016; 49: 161–167
- [3] Feldman M, Friedman LS, Sleisenger MH. *Gastrointestinal and liver disease. Volume 1. Pathophysiology, diagnosis, management.* 10th edn. Philadelphia: Saunders-Elsevier; 2016

Bibliography

DOI <http://dx.doi.org/10.1055/s-0043-100216>
Endoscopy 2017; 49: E97–E98
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X