Tunnel-and-bridge strategy for rectal endoscopic submucosal dissection: tips to allow strong countertraction without clip and line

Endoscopic submucosal dissection (ESD) is the reference method for curative resection of rectal tumors, reducing the rate of recurrence and allowing complete specimen assessment [1]. Nevertheless, it is technically challenging and new strategies to reduce the procedure time and difficulty, and to improve learning are needed [2–4].

Countertraction, as has been proposed with the clip-and-line technique, allows the submucosal space to be enlarged, making dissection easier [5]. In the rectum, the line creates tangential traction without triangulation.

We present here the tunnel-and-bridge strategy (► Fig. 1 and ► Fig. 2; ► Video 1). First, the two edges (oral and anal) of the lesion are cut, with incision followed by dissection. A tunnel is then created from the anal to the oral edge. After this step, the two lateral incisions are made without dissection. Finally, the scope is passed through the channel and retroflexion is performed above the oral side, making permanent countertraction for the procedure to be completed by cutting the two lateral submucosal residues with the scope in the retroflexed position. Because of the weight of the scope, a real triangulation is obtained.

This strategy needs to be compared prospectively but seems to offer a good method of countertraction without any additional cost, such as for a clip.

Endoscopy_UCTN_Code_TTT_1AQ_2AD

► **Fig. 1** Schematic of the tunnel-and-bridge procedure for rectal endoscopic submucosal dissection: (1) an incision is made at the anal side; (2) an incision is made at the oral side with the scope in a retroflexed position; (3) a submucosal tunnel is created under the lesion; (4) the lateral incisions are made; (5) a bridge is created with the scope being retroflexed after being passed through the tunnel.

► Video 1

► **Video 1:** Example of rectal endoscopic submucosal dissection being performed using the tunnel-and-bridge strategy.

► **Fig. 2** Further views of the tunnel and bridge created by retroflexion of the scope.

► **Fig. 3** Endoscopic views showing: **a** the bridge effect viewed from the retroflexed position; **b** dissection of the lateral edge being performed with the scope in the retroflexed position and stretching of the tunnel.

Corresponding author

Mathieu Pioche, MD

Endoscopy unit – Digestive Disease
Department, H Pavillon – Edouard Herriot
Hospital, 69437 Lyon, France
Fax: +33-4-72110147
mathieu.pioche@chu-lyon.fr

References

- [1] Saito Y, Uraoka T, Yamaguchi Y et al. A prospective, multicenter study of 1111 colorectal endoscopic submucosal dissections (with video). *Gastrointest Endosc* 2010; 72: 1217 – 1225
- [2] Pioche M, Mais L, Guillaud O et al. Endoscopic submucosal tunnel dissection for large esophageal neoplastic lesions. *Endoscopy* 2013; 45: 1032 – 1034
- [3] Pioche M, Ciocirlan M, Lépilliez V et al. High-pressure jet injection of viscous solutions for endoscopic submucosal dissection: a study on ex vivo pig stomachs. *Surg Endosc* 2014; 28: 1742 – 1747
- [4] Pioche M, Rivory J, Nishizawa T et al. Randomized comparative evaluation of an ESD self-learning software in France and Japan. *Endoscopy* 2016; 48: 1076 – 1083
- [5] Jacques J, Geyl S, Carrier P et al. A combination of the clip-with-line method and the tunnel technique during esophageal endoscopic submucosal dissection: a technical solution? *Endoscopy* 2015; 47 (Suppl. 01): E307 – E308

Competing interests

Thierry Ponchon: Olympus: advisory board member, lecture and clinical research, Boston Scientific: advisory board member, lecture and clinical research, Cook Medical: advisory board member and clinical research, Fujifilm: lecture and clinical research, Medtronic: advisory board member, lecture and clinical research, Ipsen pharma: advisory board member, lecture and clinical research, Ferring: lecture, Nestis: shareholder.

The Authors

Mathieu Pioche^{1,2}, Jérôme Rivory¹, Vincent Lépilliez³, Jean-Christophe Saurin¹, Thierry Ponchon^{1,2}, Jérémie Jacques⁴

- 1 Gastroenterology and Endoscopy Unit, Edouard Herriot Hospital, Lyon, France
- 2 Inserm U1032, Labtau, Lyon, France
- 3 Gastroenterology and Endoscopy Unit, Private Hospital Jean Mermoz, Lyon, France
- 4 Gastroenterology and Endoscopy Unit, Dupuytren University Hospital, Limoges, France

Bibliography

DOI <http://dx.doi.org/10.1055/s-0043-100757>
Endoscopy 2017; 49: E123–E124
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X