Giant duodenal hematoma after duodenal biopsy in a patient with suspected acute gastrointestinal graft-versus-host disease

A 21-year-old man developed nausea and weight loss 2 months after undergoing allogeneic hematopoietic stem cell transplantation for treatment of acute myeloid leukemia. He was referred to gastrointestinal endoscopy for a duodenal biopsy, which is a procedure commonly used to confirm the diagnosis of acute graft-versus-host disease (GvHD) [1]. His coagulation test results were within the normal ranges and his platelet count was $52 \times 10^9/L$. The examination was uneventful and a duodenal biopsy was performed without significant bleeding (► Fig. 1).

Shortly after the intervention, the patient complained of upper abdominal pain, which initially responded to symptomatic therapy. However, after 24 hours, he presented with vomiting and progressive abdominal tenderness, suspicious of an ileus. Laboratory tests showed his hemoglobin had decreased from 4.8 mmol/L to 3.3 mmol/L and he had an elevated C-reactive protein (CRP) level at 211 mg/L. Transabdominal ultrasound revealed a large inhomogeneous hypoechoic structure in the horizontal part of the duodenum (► Fig. 2; ► Video 1). A computed tomography (CT) scan confirmed a large hematoma of the duodenal wall that was causing intestinal obstruction (► Fig. 3).

The patient was treated with a nasogastric tube, parenteral nutrition, and platelet transfusion to prevent further bleeding. Despite these attempts, he developed progressive jaundice during the following days and ultrasound confirmed common bile duct dilatation due to compression of the region of the duodenal papilla by the hematoma (► Fig. 4). The conservative therapeutic approach was continued, with frequent ultrasound and laboratory monitoring, which showed that the hematoma diminished within 3 weeks, and the patient's symptoms resolved.

► Fig. 1 Endoscopic view showing the biopsy site (arrow) in the descending duodenum.

► Fig. 2 Transabdominal ultrasound showing a large hypoechoic structure (orange line) with a mean diameter of 5 cm in the horizontal part of the duodenum. Ao, aorta.

Endoscopy-guided duodenal biopsy is the gold standard for the diagnosis of gastrointestinal GvHD and has a low incidence of severe complications [2]. However, duodenal hematoma following duodenal biopsy may occur in patients with coagulopathy and platelet dysfunction, especially after stem cell therapy [3]. The present case clearly illustrates the potentially life-threatening complications of biopsy-induced intestinal

hematoma and highlights the role of bedside ultrasound as the first-line approach for unclear abdominal symptoms after endoscopic interventions.

Endoscopy_UCTN_Code_CPL_1AH_2AB

Competing interests

None

► Video 1: Transabdominal ultrasound showing an inhomogeneous hypoechoic structure in the horizontal part of the duodenum.

► **Fig. 3** Abdominal computed tomography scan (coronal view) showing a large duodenal hematoma (orange line) and the dilated descending duodenum (arrow).

► **Fig. 4** Transabdominal ultrasound showing common bile duct dilatation up to 15 mm (arrow).

The Authors

Robert Henker¹, Sebastian Schwind², Christine Döhning², Mario Heindl¹, Maximilian Beimler³, Albrecht Hoffmeister¹, Thomas Karlas¹

- 1 Division of Gastroenterology and Rheumatology, University Hospital Leipzig, Leipzig, Germany
- 2 Division of Hematology and Oncology, University Hospital Leipzig, Leipzig, Germany
- 3 Department of Diagnostics and Interventional Radiology, University Hospital Leipzig, Leipzig, Germany

Corresponding author

Thomas Karlas, MD

University Hospital Leipzig, Division of Gastroenterology and Rheumatology, Liebigstraße 20, 04103 Leipzig, Germany
Fax: +49-341-9712209
thomas.karlas@medizin.uni-leipzig.de

References

- [1] Zeiser R, Wolff D, Scheid C et al. Graft-versus-Host Erkrankung (akut) für die DAG-KBT, Deutsche Arbeitsgemeinschaft für Knochenmark- und Blutstammzelltransplantation e.V. Available from: <https://www.onkopedia.com/de/onkopedia/guidelines/graft-versus-hosterkrankung-akut/@@view/html/index.html>. Accessed: 29.12. 2016
- [2] Gassas A, Krueger J, Schechter T et al. Safety and role of gastrointestinal endoscopy in the management of gastrointestinal acute GVHD in children after hematopoietic stem cell transplantation. *J Pediatr Hematol Oncol* 2016; 38: 453 – 456
- [3] Sierra A, Ecochard-Dugelay E, Bellaïche M et al. Biopsy-induced duodenal hematoma is not an infrequent complication favored by bone marrow transplantation. *J Pediatr Gastroenterol Nutr* 2016; 63: 627 – 632

Bibliography

DOI <http://dx.doi.org/10.1055/s-0043-103405>
Endoscopy 2017; 49: E132–E133
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X