A new technique for difficult biliary cannulation using endobiliary forceps in a patient with a periampullary diverticulum

An 83-year-old man attended the emergency room complaining of abdominal pain for 1 month. He had a history of having undergone cholecystectomy and had ongoing chronic kidney disease. His appearance was suggestive of chronic illness and there was tenderness on palpation of the right upper quadrant of his abdomen. On admission, the results of his liver function tests and blood chemistry were as follows: aspartate aminotransferase 31 IU/L, alanine aminotransferase 13 IU/L, total bilirubin 1.0 mg/dL, alkaline phosphatase 59 IU/L, γ -glutamyl transferase 12 IU/L, white blood cell count 4600/mm³ (reference 3000–9000/mm³), hemoglobin 8.2 g/dL, creatinine 3.42 mg/dL (reference 0.6–1.5 mg/dL), and albumin 2.5 g/dL. A non-enhanced abdominal computed tomography (CT) scan demonstrated a round

calcified lesion of about 15 mm in size in the distal common bile duct (CBD) (**► Fig. 1**).

Endoscopic retrograde cholangiopancreatography (ERCP) was performed with a side-viewing endoscope (TGF-240; Olympus Optical Corporation, Tokyo, Japan). A huge periampullary diverticulum was visualized; however, the ampulla of Vater was hidden inside the diverticulum (**► Fig. 2a**). A prominent round lesion, presumed to be the ampulla of Vater, was noted at the 5-o'clock position. We failed to identify the ampullary orifice despite multiple attempts to expose it using suction, torque, or papilotome manipulation [1,2].

A small-caliber endobiliary biopsy forceps (Olympus, Aomori Olympus Co., Japan) was inserted through the working channel of the endoscope. By grasping

► Fig. 1 A non-enhanced abdominal computed tomography (CT) scan image showing a round calcified lesion of about 15 mm in size (arrow) in the distal common bile duct.

the mucosa with the forceps, we were able to pull the mucosa back out of the diverticulum and the ampullary orifice was eventually turned outward (**► Fig. 2b**; **► Video 1**). After the orifice had been identified, the mucosa was secured with the forceps and a conventional cannula (Cook, Winston-Salem, North Carolina, USA) was then inserted through the same working channel, before being smoothly and successfully advanced into the bile duct (**► Fig. 2c**). Following deep cannulation, a cholangiogram demonstrated a movable filling defect in the distal CBD (**► Fig. 3**). The ductal stone was left intact and a 7-Fr plastic stent was placed into the bile duct over the guidewire (**► Fig. 4**).

In the present case, we describe a simple but useful technique using a grasp-and-pull method as a rescue technique in a patient with an ampullary orifice that was facing inward from the rim of a diverticulum.

Endoscopy_UCTN_Code_TTT_1AR_2AC

► Video 1 A periampullary diverticulum was seen, but the ampullary orifice was not visible. An endobiliary biopsy forceps was inserted through the endoscope to grasp the mucosa and pull it out of the diverticulum, thereby exposing the ampullary orifice. A conventional cannula was advanced through the same channel and the bile duct was successfully cannulated, with a plastic stent then being inserted over the guidewire.

► **Fig. 2** Endoscopic images showing: **a** a huge perampullary diverticulum with a prominent round lesion, thought to be the ampulla of Vater, at the 5-o'clock position, but the ampullary orifice not visible; **b** the mucosa being grasped with the endobiliary biopsy forceps and pulled out of the diverticulum to expose the ampullary orifice; **c** a conventional cannula smoothly cannulating the bile duct, having been advanced through the same working channel in parallel with the biopsy forceps.

Competing interests

None

The Authors

Kook Hyun Kim, Tae Nyeun Kim

Division of Gastroenterology and Hepatology,
Department of Internal medicine, Yeungnam
University College of Medicine, Daegu, Republic
of Korea

Corresponding author

Kook Hyun Kim, MD

Division of Gastroenterology and
Hepatology, Department of Internal
Medicine, Yeungnam University College of
Medicine, 170 hyunchungro, Nam-gu,
Daegu 42415, Republic of Korea
Fax: +82-53-6548386
kimkh@yu.ac.kr

► **Fig. 3** A cholangiogram showing a movable filling defect in the distal common bile duct and the biopsy forceps in position.

► **Fig. 4** Endoscopic imaging showing a 7-Fr plastic stent that was placed into the bile duct over the guidewire.

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>

References

- [1] Elmunzer BJ, Boetticher NC. Reverse guidewire anchoring of the papilla for difficult cannulation due to a periampullary diverticulum. *Gastrointest Endosc* 2015; 82: 957
- [2] García-Cano J. ERCP cannulation of a hidden papilla within a duodenal diverticulum. *Endoscopy* 2008; 40: (Suppl. 02): E53

Bibliography

DOI <https://doi.org/10.1055/s-0043-111010>
 Published online: 14.6.2017
Endoscopy 2017; 49: 824–826
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X