Successful endoscopic closure of a refractory button-battery tracheoesophageal fistula in a 3-year child using endoscopic submucosal dissection of the surrounding mucosa

► **Fig. 1** Appearance of the tracheoesophageal fistula in a 3-year-old girl after ingestion of a button battery.

Most serious button-battery ingestions are not witnessed and they can cause life threatening complications.

We present here the case of a 3-year-old girl who swallowed a button battery in January 2016 with a delayed diagnosis being made after 10 days. A 5-mm tracheoesophageal fistula was endoscopically diagnosed (► **Fig. 1**). The first attempts at closure involved the deployment of two successive esophageal covered stents between January and May (► **Fig. 2**). The fistula decreased in size but persisted, so we then attempted controlled wound healing with a nasogastric tube, but the fistula still remained. Next, we tried a side fistula abrasion with argon plasma coagulation. Unfortunately, these techniques did not allow full recovery, even though the fistula reduced notably. After 1 year, we tried endoscopic submucosal dissection (ESD) of the mucosa surrounding the fistula, resecting a 1-cm mucosal patch centered on the fistula. After injecting the submucosa and making the mucosal incision, we used a Dual-Knife (Olympus) to dissect the fibrotic area. After the dissection, the fistula was closed with three clips anchored into the submucosa of the resected area (► **Fig. 3**; ► **Video 1**). We arranged a radiologic check with contrast, which

► **Fig. 2** Radiographic images showing: **a** the fistula on a barium swallow; **b** the first attempted closure procedure with a stent positioned in the esophagus.

► **Video 1** Views of the fistula and previous attempts to close it. The endoscopic submucosal dissection procedure is performed to resect the surrounding mucosa, which is subsequently clipped to close the fistula.

► **Fig. 3** Endoscopic images showing the endoscopic submucosal dissection procedure: **a** the submucosal injection being performed; **b** the incision and dissection of the fibrosis; **c** clips placed in the submucosa.

► **Fig. 4** Follow-up imaging. **a** Chest radiograph 2 weeks later. **b** Barium swallow 3 months later.

showed no sign of a fistula in the tracheal tract, and the girl made good and rapid progress without pain. A barium swallow 5 days later showed no signs of the fistula, and she was able to eat again. At 3 months after the procedure, a barium swallow and laryngoscopy confirmed complete healing of the fistula (► **Fig. 4**). The current management of tracheoesophageal fistula after button-battery ingestion is not well defined and, although some studies have demonstrated the role of conservative treatment [1], the place of endoscopy is not known. In

all kind of fistulas, stent placement only permits healing in 53% [2]. ESD can be key in non-surgical treatment to achieve complete closure of such chronic fistulas, as was previously also shown by Rodriguez-Lago et al. [3].

Endoscopy_UCTN_Code_TTT_1AO_2AI

Competing interests

None

The Authors

Melissa Gruner¹, Sophie Heissat², Vincent Pitiot³, Paul Suy⁴, Alain Lachaux², Thierry Ponchon^{1,5}, Mathieu Pioche^{1,5}

- 1 Department of Endoscopy and Gastroenterology, Pavillon L, Edouard Herriot Hospital, Lyon, France
- 2 Department of Pediatric Endoscopy and Gastroenterology, Hôpital Femme Mère Enfant, Lyon, France
- 3 Department of Pediatric Digestive Surgery, Hôpital Femme Mère Enfant, Lyon, France
- 4 Department of Pediatric ENT, Hôpital Femme Mère Enfant, Lyon, France
- 5 Inserm U1032 LabTau, Lyon, France

Corresponding author

Mathieu Pioche, MD

Endoscopy unit – Digestive Disease department, Pavillon L – Edouard Herriot Hospital, 69437 Lyon, France
mathieu.pioche@chu-lyon.fr

References

- [1] Grisel JJ, Richter GT, Casper KA et al. Acquired tracheoesophageal fistula following disc-battery ingestion: can we watch and wait? *Int J Pediatr Otorhinolaryngol* 2008; 72: 699–706
- [2] Silon B, Siddiqui AA, Taylor LJ et al. Endoscopic management of esophagorespiratory fistulas: a multicenter retrospective study of techniques and outcomes. *Dig Dis Sci* 2017; 62: 424–431
- [3] Rodríguez-Lago I, Schroeven M, Rodenstein D et al. Endoscopic closure of tracheoesophageal fistula with submucosal dissection and a biosynthetic mesh. *Endoscopy* 2015; 47: E312–E313

Bibliography

DOI <https://doi.org/10.1055/s-0043-113549>
Published online: 29.6.2017
Endoscopy 2017; 49: E212–E214
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>