Gastric peroral endoscopic myotomy for gastroparesis, after botulinum toxin injection

A 45-year-old woman had refractory Epstein–Barr virus-associated gastroparesis. Despite lifestyle modification and medication therapy, the patient had recurrent hospitalizations for gastroparesis. Management also included four previous sessions of endoscopic botulinum toxin injection. Despite transient symptomatic improvement, the patient's symptoms had recurred. She was not a candidate for surgical implantation of a gastric pacemaker, so she was referred for gastric peroral endoscopic myotomy (G-POEM).

During the procedure (► **Video 1**), a submucosal bleb was created with a methylene blue and saline solution, 6 cm proximal to the gastroesophageal junction. A mucosal entry point was incised with a multipurpose knife and an endoscope was advanced into the submucosa. The submucosal space was dissected using intermittent injection and dissection with spray coagulation current.

In patients who have not undergone previous botulinum toxin injection, injection of the submucosa will facilitate separation between the mucosa and muscle layers (► **Fig. 1**). However, this patient's submucosal tunnel demonstrated significant scarring. The botulinum toxin caused areas of fusion of the mucosa and the muscularis with dense scarring and opaque submucosa leading to a more challenging dissection (► **Video 1**).

Once the submucosal tunnel was dissected down to the pylorus level, pyloromyotomy was performed. The submucosal tunnel was washed with topical liquid gentamicin, and the mucosal entry site was closed using multiple endoscopic sutures.

At 3 months' follow-up, the patient has regained a normal quality of life with weight gain of 10 kg.

G-POEM is a novel endoscopic therapy for refractory gastroparesis, that involves mucosal entry, submucosal tunneling, pyloromyotomy, and closure of the mu-

► **Video 1** Gastric peroral endoscopic myotomy (G-POEM) for refractory gastroparesis in a patient who had previously received botulinum toxin injections; submucosal tunnelling is compared to dissection in a patient without botulinum toxin injection.

► **Fig. 1** Submucosal tunnel dissection for gastric peroral endoscopic myotomy (G-POEM). Right: In the present patient, who had previously received botulinum toxin injections. The botulinum toxin caused areas of fusion of the mucosa and muscularis, forming a dense and opaque submucosa. Left: Submucosal tunnel in a patient who had not had botulinum toxin.

cosal entry site. It is associated with an 86% symptomatic improvement in patients with refractory gastroparesis [1]. Previous endoscopic botulinum toxin injection is associated with a more challenging submucosal tunneling (► **Fig. 1**). Recent studies recommend against endoscopic botulinum toxin injection for

gastroparesis [2–4]. ► **Video 1** highlights the submucosal fibrosis secondary to botulinum toxin injection that leads to a more challenging dissection during G-POEM.

Endoscopy_UCTN_Code_TTT_1AO_2AN

Competing interests

Michel Kahaleh MD: has received grant support from Boston Scientific, Fujinon, EMcision, Xluma Inc., W.L. Gore, MaunaKea, Apollo Endosurgery, Cook Endoscopy, ASPIRE Bariatrics, GI Dynamics, NinePoint Medical, Merit Medical, Olympus and MI Tech. He is a consultant for Boston Scientific, Xluma Inc., Concordia Laboratories Inc, ABBvie, and MaunaKea Tech.

All other authors have no conflicts of interest to report.

The Authors

Monica Saumoy¹, Najib Nassani¹, Joaquin Ortiz², Viviana Parra³, Amy Tyberg¹, Michel Kahaleh¹

- 1 Division of Gastroenterology and Hepatology New York Presbyterian Hospital, Weill Cornell Medical College, New York, United States
- 2 Instituto Nacional de Cancerologia, Delegacion Tlalpan, Mexico City, Mexico
- 3 Clinica Universitaria Columbia, Bogota, Colombia

Corresponding author

Michel Kahaleh, MD

1305 York Ave, 4th Floor, New York, NY 10021, United States
Fax: 01-646-962-0110
mkahaleh@gmail.com

References

- [1] Khashab MA, Ngamruengphong S, Carr-Locke D et al. Gastric per-oral endoscopic myotomy for refractory gastroparesis: results from the first multicenter study on endoscopic pyloromyotomy (with video). *Gastrointest Endosc* 2017; 85: 123 – 128
- [2] Arts J, Holvoet L, Caenepeel P et al. Clinical trial: a randomized-controlled crossover study of intrapyloric injection of botulinum toxin in gastroparesis. *Aliment Pharmacol Ther* 2007; 26: 1251 – 1258
- [3] Friedenberg FK, Palit A, Parkman HP et al. Botulinum toxin A for the treatment of delayed gastric emptying. *Am J Gastroenterol* 2008; 103: 416 – 423
- [4] Camilleri M, Parkman HP, Shafi MA et al. Clinical guideline: management of gastroparesis. *Am J Gastroenterol* 2013; 108: 18 – 37

Bibliography

DOI <https://doi.org/10.1055/s-0043-115888>
Published online: 31.7.2017
Endoscopy 2017; 49: E256–E257
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>