Deep endoscopic submucosal dissection of a refractory tracheoesophageal fistula using clip-and-line traction: a successful closure

► **Fig. 1** Refractory tracheoesophageal fistula: schematic of closure using endoscopic submucosal dissection with clip-and-line traction. **a** Dissection of a 1-cm patch. **b** Clip-and-line traction of the dissected patch and deep section. **c** Clip closure of the resected area.

Chronic tracheoesophageal fistula is a rare disease presenting a therapeutic challenge. Unlike the case with most digestive fistulas, drainage with a pigtail stent [1] is not possible.

We present here the case of a 47-year-old man referred for a chronic 2-mm tracheoesophageal fistula (23 cm from mouth) of unknown cause. His past history revealed several pulmonary infections since childhood.

Several endoscopic treatments were attempted with clip closure and then hot biopsy forceps abrasion of the surrounding mucosa, but complete closure was not obtained. We therefore proposed endoscopic submucosal dissection (ESD) of the surrounding mucosa, namely a 1-cm mucosal patch (► **Fig. 1** and ► **Fig. 2**, ► **Video 1**) centered on the fistula, as previously described [2].

The patient underwent tracheal intubation with balloon placement just under the fistula. To allow deep dissection of the fistula tract we added a clip-and-line traction (► **Fig. 3**) [3] to pull the fistula

► **Fig. 2** Endoscopic submucosal dissection (ESD) of a refractory tracheoesophageal fistula. **a** Appearance of the fistula before ESD. **b** Incision of lower edge. **c** Incision of upper edge. **d** Lateral incisions.

Video 1 Treatment of a refractory tracheoesophageal fistula by endoscopic submucosal dissection of the surrounding mucosa, with clip-and-line traction, and clip closure

Fig. 3 Clip traction and closure of tracheoesophageal fistula. **a, b** Clip-and-line traction of the dissected patch. **c** Resected area with burnt fibrosis (black area). **d** Clip closure.

out of the wall. The fistula tract was then sectioned at the deepest possible point. Finally, we closed the resected area with four clips anchored into the submucosa (► **Fig. 4a, b**). A radiographic check was done at the end of the endoscopic procedure, and opacification showed no sign of fistula in the tracheal tract. The patient's course was rapidly favorable, without pain and with disappearance of aspiration coughing. At 2 months,

radiographic monitoring with opacification showed no residual fistula. The last clinical follow-up after 4 months confirmed complete resolution of the coughing and aspiration. Currently the management of tracheoesophageal fistula is not well codified [4], and the role of endoscopy is not defined. Endoscopic abrasion with argon plasma and stents are not very effective with a cure rate of only 53% [5]. ESD

might be a good option to reduce scarring and to allow definitive resolution of these fistulas.

Endoscopy_UCTN_Code_TTT_1AO_2AI

Competing interests

None

The Authors

Gaspard Bertrand¹, Jérémie Jacques², Jérôme Rivory¹, Florian Rostain¹, Jean-Christophe Saurin¹, Thierry Ponchon^{1,3}, Mathieu Pioche^{1,3}

- 1 Department of Endoscopy and Gastroenterology, Pavillon L, Edouard Herriot Hospital, Lyon, France
- 2 Department of Endoscopy and Gastroenterology, Dupuytren Hospital, Limoges, France
- 3 Inserm U1032 LabTau, Lyon, France

Corresponding author

Mathieu Pioche, MD

Endoscopy Unit – Digestive Disease
Department, Pavillon L – Edouard Herriot
Hospital, 69437 Lyon Cedex France
Fax: +33-4-72110147
mathieu.pioche@chu-lyon.fr

References

- [1] Donatelli G, Chiche R, Cereatti F et al. Endoscopic drainage of intra-abdominal collection after bariatric surgery. *Obes Surg* 2017; 27: 1635 – 1637
- [2] Gruner M, Heissat S, Pitiot V et al. Successful endoscopic closure of a refractory button cell tracheoesophageal fistula in a 3-year child using endoscopic submucosal dissection of the surrounding mucosa. *Endoscopy* 2017; E1 – E3. doi:10.1055/s-0043-113549
- [3] Jacques J, Legros R, Rivory J et al. The “tunnel+clip” strategy standardised and facilitates oesophageal ESD procedures: a prospective, consecutive bi-centric study. *Surg Endosc* 2017 Apr 4. doi:10.1007/s00464-017-5514-0. [Epub ahead of print]
- [4] Azoulay D, Regnard JF, Magdeleinat P et al. Congenital respiratory-esophageal fistula in the adult. Report of nine cases and review of the literature. *J Thorac Cardiovasc Surg* 1992; 104: 381 – 384
- [5] Silon B, Siddiqui AA, Taylor LJ et al. Endoscopic management of esophagorespiratory fistulas: a multicenter retrospective study of techniques and outcomes. *Dig Dis Sci* 2017; 62: 424 – 431

► **Fig. 4** End of the procedure for closure of tracheoesophageal fistula. **a** Endoscopic appearance, with four clips. **b** Radiographic view showing the clips. **c** Appearance at radiographic check 2 months later.

Bibliography

DOI <https://doi.org/10.1055/s-0043-119348>

Published online: 9.10.2017

Endoscopy 2017; 49: 1278–1280

© Georg Thieme Verlag KG

Stuttgart · New York

ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>