A retroperitoneal mass confirmed as a pancreatic adenosquamous carcinoma by endoscopic ultrasound-guided fine-needle aspiration

Adenosquamous carcinoma is a rare subtype of pancreatic adenocarcinoma, accounting for less than 4% of all exocrine pancreatic malignancies and representing a ductal adenocarcinoma mixed with at least 30% of malignant squamous cells [1–3]. Herein, we report a patient with a retroperitoneal mass in whom endoscopic ultrasound-guided fine-needle aspiration (EUS-FNA) and immunohistochemical evaluation confirmed a pancreatic adenosquamous carcinoma (► **Video 1**).

A 65-year-old white man with epigastric pain and weight loss was referred for investigation of a pancreatic mass detected on abdominal computed tomography. Results of clinical laboratory examinations were unremarkable. Sectorial endoscopic ultrasound (Olympus GF-UCT140-AL5 [Olympus America Inc., New York, USA], coupled to an ultrasound unit Aloka ProSound alpha-5 SX) detected a solid lesion with irregular borders measuring 4.9×3.5 cm in the pancreatic body. EUS-FNA was performed via a transgastric approach using a 19-gauge needle (Echo-Tip Ultra Echo-19; Cook Medical, Winston-Salem, North Carolina, USA) for a total of three passes. There was no on-site cytopathologist. Histopathology on cell blocks demonstrated a mixture of adenocarcinoma cells with malignant squamous epithelium. Immunohistochemistry panel was positive for S100p, cytokeratin 7 (both markers of glandular differentiation), p63, and cytokeratin 5/6 (both markers of squamous epithelium),

but negative for chromogranin A and mucin 5AC.

Optimal treatment remains undefined for the disease. The patient was offered chemotherapy with gemcitabine. After two cycles, liver metastases were detected. A new chemotherapy protocol was begun with oxaliplatin, folinic acid, and fluorouracil. There was no response and the patient died 6 months later.

Endoscopy_UCTN_Code_CCL_1AF_2AZ_3AB

► **Video 1** Adenosquamous carcinoma is an aggressive subtype of adenocarcinoma, accounting for less than 4% of all exocrine pancreatic malignancies. Endoscopic ultrasound-guided fine-needle aspiration can guarantee the diagnosis in the presence of a ductal adenocarcinoma mixed with malignant squamous cell carcinoma and specific immunohistochemical markers.

Competing interests

None

The authors

César Vivian Lopes¹, Antônio Dal Pizzol Júnior², Karen Margarita Rico Escamilla³, Antônio Atalíbio Hartmann³

- 1 Department of Gastroenterology and Digestive Endoscopy, Santa Casa Hospital, Porto Alegre, Brazil
- 2 Department of Clinical Oncology, Santa Casa Hospital, Porto Alegre, Brazil
- 3 Department of Pathology, Santa Casa Hospital, Porto Alegre, Brazil

Corresponding author

César Vivian Lopes, MD

Rua Prof. Cristiano Fischer 668/1001, C.E.P.
91.410-000 Porto Alegre-RS, Brazil
Fax: +55-51-33388054
drcvlopes@gmail.com

References

- [1] Boyd CA, Benarroch-Gampel J, Sheffield KM et al. 415 patients with adenosquamous carcinoma of the pancreas: a population-based analysis of prognosis and survival. *J Surg Res* 2012; 174: 12–19

- [2] Kardon DE, Thompson LD, Przygodzki RM et al. Adenosquamous carcinoma of the pancreas: a clinicopathologic series of 25 cases. *Mod Pathol* 2001; 14: 443–451
- [3] Borazanci E, Millis SZ, Korn R et al. Adenosquamous carcinoma of the pancreas: molecular characterization of 23 patients along with a literature review. *World J Gastrointest Oncol* 2015; 7: 132–140

Bibliography

DOI <https://doi.org/10.1055/s-0043-120519>
Published online: 25.10.2017
Endoscopy 2018; 50: E13–E14
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>