Long esophageal inlet patch as a rare cause of laryngopharyngeal symptoms

A 31-year-old woman presented to our department with nonspecific laryngopharyngeal symptoms, such as globus sensation, regurgitation, and heartburn, which she had experienced for a few years. Her medical history was irrelevant. An esophagogastroduodenoscopy revealed an irregular Z-line in the distal esophagus without evidence of Barrett's esophagus (► **Fig. 1 a**), and an area of ectopic gastric mucosa in the mid-proximal esophagus, 8 cm in length and at 35 to 27 cm from the incisors, involving more than half of the esophageal lumen (► **Fig. 1 b**, ► **Video 1**). Inspection with white-light endoscopy, narrow-band imaging (NBI), and acetic acid staining showed a columnar mucosal pattern with few discrete mucosal irregularities (► **Fig. 1 c, d**). Numerous biopsies revealed a fundic-type gastric epithelium with areas of moderate inflammation with no evidence of intestinal metaplasia or dysplasia (► **Fig. 2**).

Esophageal gastric heterotopia, also called inlet patch, is a rare, most likely congenital condition with prevalence ranging from 6.9% to 13.8% in recent endoscopic studies [1,2]. It is defined as areas of gastric mucosa in the esophagus – typically in its proximal part, although it can also be seen in other parts of the esophagus [3]. Proximal inlet patches are usually small [1], and can be easily missed on routine gastroscopy. The use of advanced imaging techniques, such as NBI, increases the detection rate [4]. Most inlet patches are asymptomatic; however, because of their acid secretion characteristics, they may cause a variety of laryngopharyngeal symptoms including regurgitation, globus sensation, heartburn, and chronic cough [1]. Although rare, complications such as strictures, fistulas, ulcerations, and malignant transformation have been reported. In symptomatic patients, the treatment options include gastric acid secretion inhibitors or endoscopic treatment, such as argon plasma coagulation or radio-

► **Fig. 1** Endoscopy images. **a** Irregular Z-line in the distal esophagus (40 cm from the incisors). **b** Proximal border of the inlet patch (27 cm from the incisors). **c** Distal border of the inlet patch seen in narrow-band imaging. **d** Central part of the inlet patch (30 cm from incisors) after acetic acid staining.

► **Video 1** Endoscopy showing an 8-cm-long inlet patch in the mid-proximal esophagus examined with white-light imaging, narrow-band imaging, and after acetic acid staining.

► **Fig. 2** Histopathological examination (× 100, hematoxylin and eosin staining): fundic-type gastric epithelium with gastric glands (arrowhead) and moderate inflammation (arrow).

frequency ablation [5]. The patient described here has been treated with long-term proton pump inhibitors, ranitidine and prokinetic drugs, and has not required any additional treatment in the year following presentation.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AH

Competing interests

None

The authors

**Władysław Januszewicz^{1,2}, Anna Pietrzak^{1,2},
Małgorzata Lenarcik³, Andrzej Mróz^{2,3},
Jarosław Reguła^{1,2}**

- 1 Department of Gastroenterological Oncology, The Maria Skłodowska-Curie Memorial Cancer Centre and Institute of Oncology, Warsaw, Poland
- 2 Department of Gastroenterology and Hepatology, Medical Centre for Postgraduate Education, Warsaw, Poland
- 3 Department of Pathology and Diagnostic Laboratory, The Maria Skłodowska-Curie Memorial Cancer Centre and Institute of Oncology, Warsaw, Poland

Corresponding author

Władysław Januszewicz, MD

Department of Gastroenterological Oncology, The Maria Skłodowska-Curie Memorial Cancer Center and Institute of Oncology, Roentgen 5 Str., 02-781 Warsaw, Poland
Fax: +48-22-5463035
w.januszewicz@tlen.pl

References

- [1] Govani SM, Metko V, Rubenstein JH. Prevalence and risk factors for heterotopic gastric mucosa of the upper esophagus among men undergoing routine screening colonoscopy. *Dis Esophagus* 2015; 28: 442–447
- [2] Ohara M. Incidence of heterotopic gastric mucosa in the upper esophagus in first time narrow banding image endoscopy of consecutive 900 patients. *Gastrointest Endosc* 2010; 71: AB316–317
- [3] Chong VH. Clinical significance of heterotopic gastric mucosal patch of the proximal esophagus. *World J Gastroenterol* 2013; 19: 331–338
- [4] Chung CS, Lin CK, Liang CC et al. Intentional examination of esophagus by narrow-band imaging endoscopy increases detection rate of cervical inlet patch. *Dis Esophagus* 2015; 28: 666–667
- [5] Dunn JM, Sui G, Anggiansah A et al. Radio-frequency ablation of symptomatic cervical inlet patch using a through-the-scope device: a pilot study. *Gastrointest Endosc* 2016; 84: 1022–1026

Bibliography

DOI <https://doi.org/10.1055/s-0043-123823>
Published online: 15.12.2017
Endoscopy 2018; 50: E61–E62
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>