Fibrotic submucosal scar after endoscopic submucosal dissection (ESD) or how to divert a negative adverse event into a positive endoscopic result

Authors

Marc Barthet¹, Jean-Michel Gonzalez²

Institutions

- 1 Hôpital Nord, Marseille, France
- 2 Assistance Publique Hopitaux de Marseille – Gastroenterology, Hôpital Nord Chemin des Bourrelys, Marseille, France

Bibliography

DOI <https://doi.org/10.1055/s-0043-125327> | Endoscopy International Open 2018; 06: E354–E355

© Georg Thieme Verlag KG Stuttgart · New York
ISSN 2364-3722

Corresponding author

Marc Barthet, Hôpital Nord – Gastroenterology, Chemin des Bourrelys, Marseille cedex 20 13915, France
Fax: 33 4 91 96 87 32
marc.barthet@ap-hm.fr

The paper from K Yamamoto about application of endoscopic submucosal dissection (ESD) to reduce an inappropriate flexure of the gastric conduit opens a new door. They used the extensive fibrotic scar following ESD to reduce the gastric lumen, subsequently straightening the gastric conduit. Until now, post-ESD fibrosis was considered as the most frequent adverse event (AE) leading to lumen stenosis, with incidence ranging between 68% and 90% after involving more than 50% to 75% of the esophageal circumference for instance [1–3]. The ability to prevent such an AE is still unclear despite several available tools such as steroids with or without local injections, repeated dilatations, epithelial cell sheet translation, and biodegradable stents [4–7]. Systemic treatment with early oral steroids has been shown to be negative with gastric ESD [6]. No targeted pharmacological agents are available to decrease post-ESD fibrosis whereas biotherapy such as an anti-TNF agent exists for inflammatory diseases [6]. Other recent surgical endoscopic procedures may lead to extensive and stenotic stricture, complicating patient outcome and leading to a need for esophageal reconstruction OR endoscopic gastrojejunal anastomosis [8,9].

The goal of the Japanese team presented by Yamamoto K et al was to use the extensive fibrotic scar induced by ESD for healing an abnormal flexure of the intrathoracic stomach after esogastric anastomosis. In most cases, eating troubles following esogastric anastomosis are due to anastomotic stricture, transdiaphragmatic stenosis, injury of vagal nerve resulting in pyloric spasm and gastroparesia. In the reported case, it was due to a long flexure of the ascended stomach, which acted like a U-bend. The authors performed an 8 cm ESD of the normal contra-lateral gastric side to straighten the gastric conduit. Despite a perforation, 10 days later, a straight gastric conduit was dem-

onstrated. Eighteen days later, the patient was eating and 35 days later, the patient was discharged from the hospital. Surgeons may argue that the 35-day hospital stay was very long, maybe longer than with uncomplicated surgery.

Healing approaches in this situation may be surgery or endoscopic alternatives such as suturing with overstitch suture devices, making a kind of sleeve gastropasty. The authors chose to divert the fibrotic side effects of ESD to a positive result by stomach retraction. Such diverting of fibrotic adverse event after endoscopic mucosal resection (EMR) is under assessment with EMR of esogastric junction for treating gastroesophageal reflux in Japan with the so-called ARMS procedure [10]. However, even if gastric ESD is mainly performed by Japanese experts, it is a more and more widespread technique in the West with a good efficiency (98% en bloc resection; 90–94% R0 rate) and low complication rate (perforation 1–4%) [11]. The example of this successful case added to an ARMS procedure or endoscopic antral plasty could lead one to wonder what could be in the future, new indications. Antireflux treatment is already under assessment with extended EMR of the esogastric junction, with promising results [10]. Endoscopic sleeve gastropasty usually is done with overstitch devices with increasing success compared to surgical sleeve gastrectomy, but one might think that multiple extended ESD could be an alternative. Endoscopic management of symptomatic diverticulum (but with high risk of perforation) or rectocele associated with dyschezia also could be interesting alternatives to surgery. Development of training models and animal experiments is required before this new endoscopic management will take off.

Competing interests

None

References

- [1] Ono S, Fujishoro M, Nimi K et al. Predictors of post-operative stricture after endoscopic submucosal dissection for superficial squamous cell neoplasms. *Endoscopy* 2009; 41: 661 – 665
- [2] Mizuta H, Nishimori I, Kuratani Y et al. Predictive factors for esophageal stenosis after endoscopic submucosal dissection for superficial esophageal cancer. *Dis Esophagus* 2009; 22: 626 – 631
- [3] Hanaoka N, Ishihara R, Takeuchi Y et al. Intralesional steroid injection to prevent stricture after endoscopic submucosal dissection for esophageal cancer: a controlled prospective study. *Endoscopy* 2012; 44: 1007 – 1011
- [4] Pioche M, Rivory J. Prevention of esophageal stenosis after large endoscopic submucosal dissection: is there a better way to use steroids. *Endosc Int Open* 2015; 3: E118 – E119
- [5] Shibagaki K, Ishimura N, Oshima N et al. Esophageal triamcinolone acetonide-filling method: a novel procedure to prevent stenosis after extensive endoscopic submucosal dissection. *Gastrointest Endosc* 24.08 2017: doi:10.1016/j.gie.2017.08.016 (pii: S0016-5107(17)32198-3)
- [6] Kishida Y, Kakushiman N, Takizawa M et al. Effects of steroids use for stenosis prevention after wide endoscopic submucosal dissection for gastric neoplasm. *Surg Endosc* 21.07 2017: doi:10.1007/s00464-017-5732-5
- [7] Jain D, Singhal S. Esophageal stricture prevention after endoscopic submucosal dissection. *Clin Endosc* 2016; 49: 241 – 256
- [8] Gonzalez JM, Vanbiervliet G, Gasmi M et al. Efficacy of endoscopic rendez-vous technique for the reconstruction of complete esophageal. *Endoscopy* 2015; 48: 179 – 183
- [9] Vanbiervliet G, Bonin E, Garces-Duran R et al. Gastrojejunal anastomosis using a tissue apposition stent : a feasibility study in live pigs. *Endoscopy* 2014; 46: 871 – 877
- [10] Inoue H, Minami H, Kobayashi Y et al. Anti-reflux mucosectomy for gastroesophageal reflux disease in the absence of hiatal hernia: a pilot study. *Ann Gastroenterol* 2014; 27: 346 – 351
- [11] Ma MX, Bourke M. Endoscopic submucosal dissection in the West: Current status and future directions. *Dig Endosc* 2017: doi:10.1111/den.12960 [Epub ahead of print]