



Mature Teratoma with Somatic-Type Malignancy: An Entity of Unacquaintance—A Case Report

Batuk Diyora¹ Kavini Devani¹ Sridhar Epari² Gauri Deshpande² Anup Purandare¹ Ravi Wankhade¹

¹Department of Neurosurgery, LTMG Hospital, Mumbai, Maharashtra, India

²Department of Pathology, Tata Memorial Centre, Homi Bhabha National Institute, Tata Memorial Hospital and ACTREC, Mumbai, Maharashtra, India

Address for correspondence Batuk Diyora, DNB, Department of Neurosurgery, Second Floor, College Building, LTMG Hospital, Sion, Mumbai, 400022, Maharashtra, India
(e-mail: bddiyora@gmail.com; batuk73@yahoo.co.in).

AJNS 2023;18:366–371.

Abstract

Keywords

- ▶ germ cell tumors
- ▶ teratoma
- ▶ malignant transformation
- ▶ adenocarcinoma
- ▶ cranial

Primary intracranial teratomas are nongerminomatous germ cell tumors. They are infrequent lesions along the craniospinal axis, with their malignant transformation extremely uncommon. A 50-year-old-male patient presented with one episode of generalized tonic–clonic seizure (GTCS), without any neurological deficit. Radiological imaging revealed a large lesion in the pineal region. He underwent gross total excision of the lesion. Histopathological examination was representative of teratoma with adenocarcinomatous malignant transformation. He underwent adjuvant radiation therapy and had an excellent clinical outcome. The present case highlights the rarity of malignant transformation of the primary intracranial mature teratoma.

Introduction

Central nervous system (CNS) teratomas, a group of non-germinomatous germ cell tumors (GCTs), are rare neoplasms containing tissues derived from all three germ cell layers. These account for 0.3 to 0.6% of all intracranial tumors,¹ with a slightly higher incidence in the pediatric age group and male predominance.^{2,3} They can be found at any location along the craniospinal axis, usually in the midline,⁴ and have a varied presentation depending on their location. They can further be categorized into three subtypes, including mature teratoma, immature teratoma, and teratoma with malignant transformation (TMT).⁵ Their prognosis is good, with 5-year survival rates ranging from 87 to 100% for mature teratomas and 33 to 71% for malignant teratomas.⁶ A teratoma with somatic-type malignancy

is an exceptionally rarely encountered lesion with only six cases reported so far (▶Table 1).

Case Report

A 50-year-old male patient presented with one episode of generalized tonic–clonic convulsion (GTCC) and drowsiness. He had a history of holocranial headaches for 4 months. There were no other neurological symptoms. His neurological examination revealed no abnormality. Computed tomography (CT) scan of the brain revealed approximately 8.1 × 9.7 × 9.6 cm size, well-defined, multilobulated, extra-axial, solid-cystic lesion with calcification in the region of the quadrigeminal cistern and left posterior parasagittal area (▶Fig. 1A,B). The lesion extended underneath the falx toward the right side. Magnetic resonance imaging (MRI)

article published online
June 7, 2023

DOI <https://doi.org/10.1055/s-0043-1768603>.
ISSN 2248-9614.

© 2023. Asian Congress of Neurological Surgeons. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

Table 1 A summary of previously reported cases of teratoma with somatic-type malignancy

Sl. no.	Study	Age & sex	Symptoms	Location	Radiological features	Approach	HPE	Results
1	Freilich et al ¹¹	Case 1 26/M	Persistent Headache, nausea, vomiting	Pineal region	Solid nodular tumor on CT with hydrocephalus	Not mentioned; subtotal resection + postoperative radiation	Yolk sac tumor with mucin secreting adenocarcinomatous transformation of teratoma	Recurrence after 6 mo
		Case 2 8/M	Persistent headache, nausea, vomiting	Pineal region	Cystic lesion initially; solid cystic lesion after 4 y	VP shunt followed by radiation; subtotal resection + postoperative radiation	Yolk sac tumor initially; enteric-type adenocarcinomatous transformation of teratoma	Recurrence after 6 mo
2	Matsutani et al ¹⁷	Cases 3-6 31 ± 14.6/M	Symptoms of raised intracranial pressure	Pineal region (1) Neurohypophysis (1) Other regions (2)	Solid-cystic lesion	Infratentorial approach (1) Pterional approach (3) Subtotal resection	Epidermal carcinoma (3) Sarcoma (1)	Intermediate prognosis
3	Current study	50/M	Headache, GTCS	Pineal region	Solid cystic lesion	Left parieto-occipital approach; gross total excision + postoperative radiation	Adenocarcinomatous transformation of mature teratoma	Good prognosis

Abbreviations: CT, computed tomography; GTCS, generalized tonic-clonic seizure; VP, ventriculoperitoneal.

revealed a similar-sized, well-defined, multilobulated, extra-axial, solid-cystic lesion in the region of the quadrigeminal cistern, with the pineal gland not separately visible from the lesion. The solid component of the lesion appeared isointense on T1-weighted images and heterogeneously hyperintense on T2-weighted images with multiple foci of calcification. The lesion was heterogeneously enhancing postcontrast administration. The peripheral cystic component appeared iso-hypointense on T1-weighted images and heterogeneously hyperintense on T2-weighted images (►Fig. 1C–G). MR spectroscopy of the solid component revealed an elevated choline peak with a high choline/creatinine ratio of 5.3 and a mildly raised choline/N-acetyl-aspartate (NAA) ratio of 1.7. Diffusion tensor imaging (DTI) revealed displacement of bilateral corticospinal tracts, left superior longitudinal fasciculus, left inferior longitudinal fasciculus, and splenium (►Fig. 1H). Blood and cerebrospinal fluid (CSF) obtained via lumbar puncture values of alpha-fetoprotein (AFP) and beta-human chorionic gonadotrophin (B-HCG) hormone were within normal limits. CSF cytology was negative.

The patient underwent a left parasagittal craniotomy in the sitting position. A large, solid-cystic, extra-axial, mildly vascular, easily suckable lesion was encountered (►Fig. 2). Gross total excision of the lesion was achieved. Convalescence was uneventful. Histopathological examination of the tissue sample showed a heterogeneous tumor (►Fig. 3A,B), predominantly composed of admixed mature adipose tissue, fascicles of mature skeletal muscle (►Fig. 3C), mature chondroid lobules (►Fig. 3D), and respiratory and glandular columnar epithelium-lined ducts with mucoid material (►Fig. 3D–F). This predominant component of the tumor was consistent with mature teratoma. In addition to this component, an area with distinctive morphology composed of back-to-back arranged glands endowed with architectural and cytomorphological atypia (►Fig. 3G, lower half, H,I), nuclear dysplasia and luminal necrosis (►Fig. 3I), which was consistent with intestinal-type adenocarcinoma. The foci of dystrophic calcification were noted both within the intraglandular component of the tumors and in the adjacent adherent, cortical brain parenchyma. On immunohistochemistry, the adenocarcinomatous component was positive for CK20 and p53 protein and negative for CK7 and glypican-3 (►Fig. 3J,K). The overall histomorphology features were of an adenocarcinoma arising in the background of a mature teratoma (i.e., teratoma with somatic-type malignancy). The tumors showed wild type for BRAFV600 (►Fig. 3L), KRAS (►Fig. 3M), and NRAS and HRAS sequence on Sanger sequencing, suggesting no mutations in the genes as mentioned earlier.

Postoperative MRI was consistent with the gross total excision of the lesion (►Fig. 4A,B). The patient was subjected to positron emission tomography (PET) to rule out systemic malignancy. There was no evidence of systemic lesion. However, a small residue was noted in the brain (►Fig. 4C). Postoperatively, he was administered 54-Gy radiation in 30 fractions.




Fig. 1 Computed tomography (CT) scan of the brain showing large heterogeneous midline mass containing irregular areas of high attenuation (calcification), low attenuation (fat), and intermediate attenuation (soft tissue). (A,B) It causes moderate asymmetric dilatation of the right lateral ventricle with a shift of the midline structures to the right. Magnetic resonance imaging (MRI) of the brain showing large well-defined, multilobulated extra-axial solid cystic lesion in the region of quadrigeminal cistern with the pineal gland not separately visible from the lesion. The solid component of the lesion appeared isointense on T1-weighted images in (C) axial view and (D) sagittal view, heterogeneously hyperintense on T2-weighted images in (E) axial view and (F) coronal view, with multiple foci of calcification. (G) The lesion was heterogeneously enhancing on postcontrast administration. The peripheral cystic component appeared iso-hypointense on T1-weighted images and heterogeneously hyperintense on T2-weighted images. (H) Diffusion tensor imaging (DTI) revealed displacement of bilateral corticospinal tracts, left superior longitudinal fasciculus, left inferior longitudinal fasciculus, and splenium.

Discussion

Extragenadal GCTs have no evidence of a primary tumors in the testis or ovary.⁷ Extragenadal GCTs can arise anywhere in the midline, including the retroperitoneum, anterior mediastinum, or along the craniospinal axis in the midline.⁴ CNS GCTs are grouped into three subgroups: germinomatous, nongerminomatous (yolk sac tumor, teratoma, embryonic carcinoma, endodermal sinus tumors, choriocarcinoma), and mixed GCTs.⁸ They comprise only 0.3 to 0.6% of all intracranial tumors.¹ The most commonly proposed mechanisms for these tumors are KIT, KRAS/NRAS, and CBAL mutation.⁹ These tumors show male predominance with an overall male-to-female ratio being 3:1 with approximately over 90% incidence in a young population.

Their pathophysiology is not clearly understood; however, there are two proposed theories: germ cell theory and embryonic theory.¹⁰ The germ cell theory hypothesizes that GCTs arise from primordial germ cells, which have migrated aberrantly during embryogenesis and later undergo malignant transformation. According to the embryonic theory, these tumors arise from mixed migrational pluripotent germ cells. Another school of thought proposes that germinomas arise from germ cells. In contrast, other nongerminomatous GCTs, including teratomas, occur due to misfolding or misplacement of embryonic cells into the lateral mesoderm

causing their entrapment in various parts of the CNS.^{10,11} The movement of the primordial germ cell migration, which coincides with the fetal hypothalamus maturation, is influenced by a few trophic molecules like growth factor β . It is believed that the hypothalamus may produce chemotropic factors like growth factor- β , which can cause mis-migration.¹¹ Teratomas are a subtype of nongerminomatous GCTs, which by definition are tumors of multipotential cells derived from all three germ cell layers. They are hypothesized to arise from the misplacement of pluripotent germ cells.¹² They are delineated into three histological variants by the World Health Organization (WHO): mature, immature, and TMT.⁸ According to the 2021 WHO CNS 5 classification, the term TMT has been revised to teratoma with somatic-type malignancy.¹³

Mature teratomas are composed of well-differentiated tissue consisting of tissues of all three germ layers and low mitotic activity without necrosis. In contrast, immature teratoma consists of components resembling fetal tissue, like hypercellular embryonic mesenchyme, or primitive neuroectodermal elements mixed with mature tissue elements.¹⁴ Teratoma with somatic-type malignancy is a generic designation of a teratomatous neoplasm with an additional malignant somatic tissue. The malignant component is usually a rhabdomyosarcoma or an undifferentiated sarcoma and, less commonly, a squamous cell carcinoma or an enteric




Fig. 2 Intraoperative photographs showing (A) a multiloculated cystic component with straw colored fluid and (B) heterogeneous soft tissues, including hemorrhagic foci, yellowish cheesy soft parts and calcification.

type adenocarcinoma. Yolk sac tumor elements have also been considered progenitors of enteric-type adenocarcinoma from intracranial GCTs. Immunohistochemical examination plays a pivotal role in the diagnosis of TMT. Stains like vimentin, desmin, SMA, S-100, CD99, and GFAP antibodies are usually performed in the cases with sarcomatous transformation,¹⁵ whereas CK20, CK7, and p53 are performed in patients with carcinomatous transformation.¹⁶ Glypican-3 is used to rule out any yolk sac element.¹⁷

Intracranial teratoma usually arises from midline structures, including the pineal region, quadrigeminal plate, wall of the third ventricle, suprasellar area, or cerebellar vermis. The pineal is the most common region where they are found.⁴ However, other less common locations from which intracranial GCT can arise include the cerebral hemispheres, ventricles, thalamus, basal ganglia, and medulla oblongata.²

There are a gamut of presenting features of these tumors, which are usually based on their location. They can present with only distressing localized or holocranial headaches or accompanied nausea and vomiting. They can cause visual disturbances like blurred vision, diplopia, or vertical gaze palsy. Not uncommonly, they can also present with GTCS or focal seizures. The ones in the suprasellar region can lead to dysfunction of the hypothalamus and pituitary gland, leading to features of diabetes insipidus, precocious puberty,

developmental retardation, sexual precocity, isolated growth hormone deficiency hypopituitarism, and visual field defects like bitemporal hemianopia.¹¹

A clinical diagnosis is often based on imaging, tumor marker detection, and CSF cytology. A CT and a contrast MRI with spectroscopy are required to reach a diagnosis and for operative planning. In patients later diagnosed with TMT, it is prudent to rule out primaries elsewhere in the body by getting a PET CT. Intracranial teratomas can be large at the time of their presentation. Due to extreme variability in their content, imaging findings are also heterogeneous. Most of these lesions contain fat and calcification that is usually solid/clumplike. On MRI, they typically reveal lesions of mixed signal intensity due to their various components. On T1-weighted images, the lesion shows hyperintense, isointense, and hypointense signals due to fat/proteinaceous/lipid-rich fluid, soft tissue, and calcification/blood product, respectively. On T2-weighted images, the lesion shows mixed signals. Soft-tissue component usually enhances on intravenous gadolinium administration.¹⁸

Detection of B-HCG and AFP in CSF and serum aids in detecting the tumor type and its further management. If possible, tumor markers should be detected in both CSF and serum, as it has been shown that the amount of β -hCG is substantially higher in the CSF, whereas that of AFP is slightly higher in the serum. A malignant teratoma would not present with a high concentration of AFP and β -hCG, which indirectly helps rule out other GCTs. CSF cytology should be routinely performed to rule out tumor dissemination into CSF. Patients with positive cytology have a poor prognosis.²

Management is aimed at radical resection. They are chemo-resistant, and recurrence is common. Complete microsurgical resection of residual or recurrent tumors provides the best chance of prolonging survival.¹⁹ The location of the tumor decides the approach to the tumors. Excision is considered complete if more than 90% tumor is resected.¹² An attempt should be made to excise as much tumor as possible because the tumor left with a malignant component leads to recurrence in the long term. In comparison to GCTs, NGGCTs are less radiosensitive. So chemotherapy and radiation therapy, with or without surgery, is usually required. In patients with teratoma with somatic-type malignancy, a search for the primary should be done. If no primary is found, the tumor bed should be irradiated to prevent a recurrence. When the tumors produce B-HCG or AFP markers, they can monitor treatment and early detection of recurrence.

The prognosis in a case of teratoma depends on its subtype. Mature teratomas containing well-differentiated elements of all three germ cell layers are associated with a good prognosis, with 10-year survival rates of more than 90%. In contrast, immature teratomas, with their undifferentiated components and the possibility of developing malignant tumors, have a less favorable prognosis with a 5-year survival rate of around 70%. Teratoma with somatic-type malignancy containing various components of conventional malignant tumors have a poor prognosis.^{20,21} Five-year survival rates are better for patients with mature teratoma




Fig. 3 Representative photomicrographs (A–I: HE; J,K: immunohistochemistry; L,M: electropherogram). The tumor shows heterogeneous morphological areas (A,B: HE, $\times 10$) composed predominantly of admixed areas of mature adipose tissue and skeletal muscles (C: HE, $\times 100$), columnar epithelium-lined structures (D,E: HE, $\times 200$), chondroid lobule (D: HE, $\times 200$), dilated epithelium-lined ducts with mucoid material (F: HE, $\times 100$). Additionally, a morphologically distinct cellular area (as represented in the lower half in the G: HE, $\times 100$) composed of packed glands (H: HE, $\times 100$) which are architectural and nuclear atypia (I: HE, $\times 200$). Luminal necrosis noted (I: HE, $\times 200$). Electropherograms showing wild-type BRAFV600 sequence (L) and exon 3 KRAS sequence (M).




Fig. 4 Postoperative magnetic resonance imaging (MRI) (A) T1-weighted and (B) postcontrast axial images showing gross total resection of the tumor with a small remnant near the mesial temporal lobe. (C) Postoperative position emission tomography (PET) scan showing a small remnant.

than patients with teratoma with somatic-type malignancy. In patients with malignant transformation, a search for primary should be done.¹⁵ If no primary is found, the tumor bed should be irradiated to prevent a recurrence. They have an aggressive clinical course than mature or immature teratoma.²²

Conclusion

CNS teratoma is an uncommon tumor, and adenocarcinomatous transformation has made the tumor extremely rare as only six cases have been reported so far to the best of our

knowledge. Owing to the rarity of these lesions, their treatment strategy remains controversial. Patient age, biochemical markers, and histological diagnosis aid in their management. Radical resection, followed by adjuvant radiation, forms the cornerstone of its management.

Informed Consent

Patient's informed consent was obtained for this study.

Conflict of Interest

None declared.

References

- Georgiu C, Opincariu I, Cebotaru CL, et al. Intracranial immature teratoma with a primitive neuroectodermal malignant transformation: case report and review of the literature. *Rom J Morphol Embryol* 2016;57(04):1389–1395
- Kong Z, Wang Y, Dai C, Yao Y, Ma W, Wang Y. Central nervous system germ cell tumors: a review of the literature. *J Child Neurol* 2018;33(09):610–620
- Lee YH, Park EK, Park YS, Shim KW, Choi JU, Kim DS. Treatment and outcomes of primary intracranial teratoma. *Childs Nerv Syst* 2009;25(12):1581–1587
- Ramamurthy B, Tandon PN. *Textbook of Neurosurgery*. New Delhi: BI Publications; 1992:1034–1036
- Romić D, Raguž M, Marčinković P, et al. Intracranial mature teratoma in an adult patient: a case report. *J Neurol Surg Rep* 2019;80(01):e14–e17
- Zygourakis CC, Davis JL, Kaur G, et al. Management of central nervous system teratoma. *J Clin Neurosci* 2015;22(01):98–104
- Ait Benali H, Lalya L, Allaoui M, et al. Extragonadal mixed germ cell tumor of the right arm: description of the first case in the literature. *World J Surg Oncol* 2012;10:69
- Rosenblum MK, Nakazato Y, Matsutani M. CNS germ cell tumors. In: Louis DN, Ohgaki H, Weistler OD, Cavenee WK, eds. *WHO Classification of Tumors of the Central Nervous System*. 4th ed. Lyon: IARC; 2007:198–204
- Kubota Y, Seki M, Kawai T, et al. Comprehensive genetic analysis of pediatric germ cell tumors identifies potential drug targets. *Commun Biol* 2020;3(01):544
- Echevarría ME, Fangusaro J, Goldman S. Pediatric central nervous system germ cell tumors: a review. *Oncologist* 2008;13(06):690–699
- Palmer RD, Foster NA, Vowler SL, et al. Malignant germ cell tumours of childhood: new associations of genomic imbalance. *Br J Cancer* 2007;96(04):667–676
- Agrawal M, Uppin MS, Patibandla MR, et al. Teratomas in central nervous system: a clinico-morphological study with review of literature. *Neurol India* 2010;58(06):841–846
- Mahajan S, Suri V, Sahu S, Sharma MC, Sarkar C. World Health Organization Classification of Tumors of the Central Nervous System 5th Edition (WHO CNS5): what's new? *Indian J Pathol Microbiol* 2022;65(Supplement):S5–S13
- Kobayashi T, Lunsford LD, eds. *Pineal Region Tumors. Diagnosis and Treatment Options. Progress in Neurological Surgery Vol 23*; Basel: Karger; 2009:59–75
- Freilich RJ, Thompson SJ, Walker RW, Rosenblum MK. Adenocarcinomatous transformation of intracranial germ cell tumors. *Am J Surg Pathol* 1995;19(05):537–544
- Selves J, Long-Mira E, Mathieu MC, Rochemaix P, Ilié M. Immunohistochemistry for diagnosis of metastatic carcinomas of unknown primary site. *Cancers (Basel)* 2018;10(04):108
- Zynger DL, McCallum JC, Luan C, Chou PM, Yang XJ. Glypican 3 has a higher sensitivity than alpha-fetoprotein for testicular and ovarian yolk sac tumour: immunohistochemical investigation with analysis of histological growth patterns. *Histopathology* 2010;56(06):750–757
- Buetow PC, Smirniotopoulos JG, Done S. Congenital brain tumors: a review of 45 cases. *AJR Am J Roentgenol* 1990;155(03):587–593
- Comiter CV, Kibel AS, Richie JP, Nucci MR, Renshaw AA. Prognostic features of teratomas with malignant transformation: a clinicopathological study of 21 cases. *J Urol* 1998;159(03):859–863
- Matsutani M, Sano K, Takakura K, et al. Primary intracranial germ cell tumors: a clinical analysis of 153 histologically verified cases. *J Neurosurg* 1997;86(03):446–455
- Ogawa K, Toita T, Nakamura K, et al. Treatment and prognosis of patients with intracranial nongerminomatous malignant germ cell tumors: a multiinstitutional retrospective analysis of 41 patients. *Cancer* 2003;98(02):369–376
- Kyritsis AP. Management of primary intracranial germ cell tumors. *J Neurooncol* 2010;96(02):143–149