



Postoperative Contralateral Spontaneous Epidural Hematoma

Abhijit Acharya¹ Ashok Kumar Mahapatra¹ Soubhagya Tripathy¹ Souvagya Panigrahi¹
Rama Chandra Deo¹ Satya Bhusan Senapati¹

¹ Department of Neurosurgery, Institute of Medical Sciences (IMS) and SUM Hospital, Siksha 'O' Anusandhan (SOA) University, Bhubaneswar, Odisha, India

Address for correspondence Abhijit Acharya, MCh, Department of Neurosurgery, Institute of Medical Sciences (IMS) and SUM Hospital, Siksha 'O' Anusandhan (SOA) University, Bhubaneswar, Odisha, India (e-mail: abhijitkirtika@gmail.com).

Indian J Neurosurg

Postoperative contralateral extradural hematoma (EDH) is a rare condition.¹ The dura lies closely adherent to the calvarium which prevents collection of any hematoma. However, due to trauma or nontraumatic causes, there can be acute collection of blood in the space resulting in EDH. The most common cause of EDH is trauma of various etiologies. Ten percent of all traumatic head injuries result in EDH. Traumatic injuries consist of road traffic accidents and violent collisions of the head, which induce bleeding due to the tearing of meningeal vessels like arteries, veins, or sinuses. Nontraumatic spontaneous EDH causes are rare. Spontaneous EDH patients are categorized into primary (cryptogenic) and secondary EDH. Nontraumatic causes of EDH are coagulation abnormalities, blood dyscrasias like Sickle cell disease and hemophilia, vascular malformations, dural arteriovenous fistulae, infective causes like meningitis, etc.²

Sarkari et al published an article "Delayed opposite frontal epidural hematoma due to bleeding from superior sagittal sinus without any cranial fracture; A case report" in the *Indian Journal of Neurotrauma* in 2012 about a 24-year-old patient who presented with an EDH due to trauma and was operated. He developed a spontaneous EDH on the opposite side following the first surgery after 3 hours.³

Spontaneous EDH intraoperatively or in the postop period is a fatal complication that may be missed many times. Although rare, multiple case reports have been published from time to time regarding the same (► **Table 1**). Few were detected early and managed surgically whereas the rest succumbed due to delayed diagnosis. Spontaneous EDH can result during surgery or even in the postoperative period. In spontaneous intraoperative, the preoperative computed tomography (CT) or magnetic resonance imaging suggests a normal scan on the

contralateral side. Spontaneous hematomas develop during the surgical period, which requires a keen clinical acumen to suspect such a possibility. Tense dura intraoperatively is a sign of suspect bleeds on the contralateral side (► **Fig. 1**).⁴ In suspected cases, an early postoperative CT scan is required to exclude EDH. Postoperative spontaneous epidural hematoma on the contralateral side of the brain develops due to loss of the tamponade effect on the contralateral side. Preoperative noncontrast CT brain may not reveal any bleeding; however, there may be a fracture of calvarial bone present on the opposite side, which may be seen only in 75% of the patients. There may be underlying torn vessels like artery, venous, or dural sinuses, which lie compressed due to the tamponade effect. As soon as the ipsilateral EDH is evacuated, the tamponade effect on the opposite side is over, which leads to the gradual bleeding and formation of a spontaneous EDH in the postop period.⁵ Early imaging (within 1–2 hours) may miss the pathology on the contralateral side. Risk factors like the rapid evacuation of EDH on the ipsilateral side, use of anticoagulants, etc. can lead to spontaneous EDH. Tense dura intraoperatively, with massive brain bulge on durotomy due to massive brain shift, bradycardia, irregular breathing pattern, and irregular pupils with development of dilatation of pupil on the contralateral side

Table 1 Causes of postoperative contralateral hematoma

Serial	Causes
1	Injury to superior sagittal sinus
2	Contralateral Skull Fracture
3	Bleeding disorder
4	Use of anticoagulant for other systemic causes

DOI <https://doi.org/10.1055/s-0044-1786974>.
ISSN 2277-954X.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India




Fig. 1 Spontaneous extradural hematoma (EDH) in a postoperative case on the contralateral side.

can lead to suspicion. Management is surgery in the form of craniotomy and EDH evacuation.⁶

Early detection and management can save the patient from complications and mortality. Sharma et al reported a case in 2015 where a 28-year-old patient developed spontaneous EDH intraoperatively, which was detected and managed early.⁷ Rarely EDH occur due to tear of the sagittal sinus and Mohindra et al have recommended routine postoperative CT immediately after surgery for a traumatic head injury to detect contralateral abnormality which will help in timely intervention.⁸

Spontaneous hematoma cannot be prevented; however, we can take some precautionary measures to avoid the development of contralateral hematomas:

Serial no.	Precautionary measures
1	Avoid giving anticoagulants
2	Avoid osmotic like mannitol
3	Avoid head raising if dura is too lax
4	PCO ₂ to maintain the upper limit
5	Valsalva

Conflict of Interest

None declared.

Acknowledgment

We acknowledge SOA University for the unconditional support.

References

- 1 Su TM, Lee TH, Chen WF, Lee TC, Cheng CH. Contralateral acute epidural hematoma after decompressive surgery of acute subdural hematoma: clinical features and outcome. *J Trauma* 2008;65(06):1298–1302
- 2 Rosenthal AA, Solomon RJ, Eyerly-Webb SA, et al. Traumatic epidural hematoma: patient characteristics and management. *Am Surg* 2017;83(11):e438–e440
- 3 Sarkari A, Satyarthee G, Mahapatra AK, et al. Delayed opposite frontal epidural hematoma due to bleeding from superior sagittal sinus with no fracture- a case report. *J. Ind J Neurot* 2012;8:7–10
- 4 Singh M, Ahmad FU, Mahapatra AK. Intraoperative development of contralateral extradural hematoma during evacuation of traumatic subdural hematoma; a rare cause of malignant brain bulge during surgery. *Indian J Neurotrauma* 2005; 2:139–140
- 5 Feuerman T, Wackym PA, Gade GF, Lanman T, Becker D. Intraoperative development of contralateral epidural hematoma during evacuation of traumatic extraaxial hematoma. *Neurosurgery* 1988;23(04):480–484
- 6 Wani, et al. Bilateral sequential developed asynchronous extradural hematoma. *Indian J Neurotrauma* 2010;7:79–80
- 7 Sharma A, Sharma A, Dewan Y. Intraoperative contralateral extradural hematoma during evacuation of traumatic acute extradural hematoma. *Rom Neurosurg* 2015;29(03):377–381
- 8 Mohindra S, Mukherjee KK, Gupta R, Chhabra R, Gupta SK, Khosla VK. Decompressive surgery for acute subdural haematoma leading to contralateral extradural haematoma: a report of two cases and review of literature. *Br J Neurosurg* 2005;19(06): 490–494