Small Neuroendocrine Carcinoma of the Rectum Entirely Covered by an Adenomatous Component

H. Yoshikane¹, D. Arakawa¹,
H. Kawashima¹, A. Sakakibara¹,
H. Hidano¹, H. Takahashi², T. Yokoi³

¹ Dept. of Internal Medicine,
Handa City Hospital, Handa, Japan

² Dept. of Pathology,
Handa City Hospital, Handa, Japan

³ Division of Pathology,
Aichi Medical University, Aichi, Japan

Corresponding Author

H. Yoshikane, M.D.
Dept. of Internal Medicine
Handa City Hospital
2-29 Toyo, Handa
Aichi, 475
Japan

Fax: + 81-569-243253

E-mail: winwin@cac-net.ne.jp

Figure 1 **a** Neuroendocrine carcinoma (NEC) of the large intestine is a rare condition with an extremely poor prognosis. At autopsy, a laterally spreading tumor (LST) 5 cm in diameter was found in the rectum in a 54-year-old woman who had multiple liver and bone metastases of unknown origin. **b** Cross section of the LST revealed, however, the coexistence of a whitish round tumor (arrow) beneath the brownish LST.

Figure 2 Pathological examination revealed the LST to be a tubulovillous adenoma and the round tumor to be NEC. The extensive metastases resulted from the NEC which was just 1 cm in diameter (haematoxylin and eosin stain, original magnification $\times 12$).