Two Cases of Small Non-Polypoid Neoplasia in an Elongated Colonic Polyp

K. Hizawa¹, T. Nakahara²,
S. Inuzuka², K. Akagi²,
R. Shimono², M. Kitamura³,
K. Iwai³, T. Matsumoto⁴

¹ Department of Internal Medicine,
Kyushu Central Hospital, Fukuoka, Japan

² Department of Surgery, Kyushu
Central Hospital, Fukuoka, Japan

³ Department of Anatomic Pathology,
Graduate School of Medical Sciences,
Kyushu University

⁴ Department of Medicine and Clinical
Science, Graduate School of Medical
Sciences, Kyushu University

Corresponding Author

K. Hizawa, M.D.
Dept. of Medicine and Clinical Science
Graduate School of Medical Sciences
Kyushu University
Maidashi 3-1-1
Higashi-ku
Fukuoka 812-8582
Japan

Fax: + 81-92-6425273
E-mail: hizawa@intmed2.med.
kyushu-u.ac.jp

Figure 1 Patient 1: colonoscopy shows a flabby potbelly-shaped pedunculated polyp (A) with a reddish and uneven head (B). The elongated polyp is composed of edematous submucosa and normal colonic mucosa (arrows), containing a depressed form of tubular adenoma (arrowheads)

Figure 2 Patient 1: the arrows indicate the tubular adenoma

Figure 3 Patient 2: colonoscopy shows an elongated polyp covered by wrinkled normal mucosa (A) with a reddish and slightly depressed surface (B). The polyp has wide areas of hyperplastic epithelium (arrows), and a well-differentiated adenocarcinoma with non-polypoid growth (arrowheads) invades the edematous submucosa

Figure 4 Patient 2: the arrows indicate the well-differentiated adenocarcinoma