Synthesis and Characterization of Annulene-Fused Pseudorotaxanes

Joshua J. Pak,^{a,1} Timothy J. R. Weakley,^a Michael M. Haley,^{*a} Danny Y. K. Lau,^b J. Fraser Stoddart^b

^a Department of Chemistry, University of Oregon, Eugene, OR 97403-1253 USA
Fax +1(541)3460487; E-mail: haley@oregon.uoregon.edu

^b Department of Chemistry, University of California – Los Angeles, 405 Hilgard Avenue, Los Angeles, CA 90095 USA
Fax +1(310)2061843; E-mail: stoddart@chem.ucla.edu

Received 1 April 2002

Abstract: A 24-crown-8 ether-fused dehydrobenzo[18]annulene has been prepared via a stepwise Pd/Cu-mediated strategy. This annulene-crown ether hybrid forms a pseudorotaxane complex in CD₂Cl₂ with dibenzylammonium hexafluorophosphate when they are combined in a 1:1 ratio, as indicated by ¹H NMR spectroscopy and mass spectrometry.

Key words: annulenes, crown compounds, macrocycles, pseudorotaxanes, supramolecular chemistry

Phenylacetylene macrocycles² and networks³ are potential building blocks for the assembly of tubular and/or porous molecular crystals.⁴ Key to such a transformation is preorganization of the precursor molecules so that they may undergo controlled oligomerization/polymerization to form ordered arrays. Unfortunately, such preorganization in the solid state⁵ is often rare and impossible to predict *a priori*. For example, tribenzo[14]annulene (**1**, Figure 1)⁶ packs in the solid state according to the parameters outlined by Enkelmann⁷ for a topochemical diacetylene polymerization. Indeed, **1** readily undergoes 1,4-polymerization when heated or exposed to UV light.⁶ Subsequent studies of a number of other dehydrobenzoannulenes (DBAs) show the reactivity displayed by **1** to be unique.⁸ More often than not the packing of the macrocycles in the crystal lattice precludes any type of ordered polymerization, which is reflected in the broader ($w^{1/2} \approx 5$ – 10 °C) than normal ($w^{1/2} < 1$ °C) DSC exotherms of the annulenes.

Figure 1 The structure of tribenzo[14]annulene (**1**)

A possible solution to the preorganization problem involves supramolecular chemistry.⁹ Over the last decade a tremendous number of ordered arrays and assemblies have been prepared utilizing various noncovalent bonding interactions (hydrogen bonding, π -stacking, dipole-dipole interactions, etc.).¹⁰ One of the more interesting and relevant examples was the recent use of supramolecular organization for the first 1,6-polymerization of a triacetylene.¹¹ Similar preorganization has been successfully used for the polymerization of a number of other diacetylenes.¹² Encouraged by these reports, we decided to target a new class of pseudorotaxanes using crown ether-dehydrobenzoannulene hybrids (e.g., **2**) in order to organize the DBA molecules. These systems can utilize external forces such as the interpenetrating guest-host interaction between secondary dialkylammonium ion centers in guests containing one or more binding sites for crown ether hosts.¹³ The resultant supramolecular complexes might thermally generate dimer fragments, which could provide insight into diacetylene polymerization for DBA systems. We describe herein our initial work to-

2

3

4

Figure 2 The structures of crown ether-dehydrobenzoannulene hybrid **2** and mono- and dicationic hexafluorophosphate salts **3** and **4**

wards the assembly of novel supramolecular architectures containing hybrid **2** and the mono- and dicationic hexafluorophosphate salts **3** and **4**, respectively (Figure 2).

Hybrid **2** was assembled using the same strategy as that reported previously for the construction of site-specifically functionalized DBAs (Scheme 1).^{8b,f,g} Cyclization of ditosylate **5**¹⁴ with 4,5-diiodocatechol¹⁵ using high dilution conditions gave diiododibenzo-24-crown-8 (**6**) in moderate yield. Cross-coupling **6** with two equivalents of 1-(4-trimethylsilylbuta-1,3-diynyl)-2-(triisopropylsilyl-ethynyl)benzene^{8g} using in situ desilylation/coupling conditions furnished the α,ω -polyne **7** in 81% yield. Desilylation of **7** with Bu_4NF and subsequent Cu-mediated intramolecular oxidative cyclization under pseudo-high dilution conditions generated the benzo-24-crown-8-fused annulene **2** as a light yellow solid in 55% yield.

Slow evaporation of a THF solution of **2** afforded yellow blocks suitable for X-ray diffraction. The crystal structure of macrocycle **2** is shown in Figure 3.¹⁶ The annulenic core is essentially planar with a mean deviation less than 0.07 Å. The bond lengths and bond angles are typical for those found in other dehydrobenzoannulenes.^{6,8} Interestingly, the crown ether curves around the end of a second annulene. Although there are no π -stacking interactions in the crystal lattice, there are short CH...O hydrogen bonds that help dictate the crystal packing. With such a curved arrangement, it might be possible to thread both the crown ether and annulene, a motif we are actively pursuing.

Mixing equimolar amounts of hybrid **2** and salt **3** in CD_2Cl_2 at room temperature resulted in a significant difference in the ^1H NMR spectra (Figure 4), thus indicating formation of the pseudorotaxane host-guest complex. The top spectrum, which represents free **2**, is relatively uncomplicated, reflecting the C_{2v} symmetry of the crown

Scheme 1 (a) 4,5-diiodocatechol, CsCO_3 , MeCN; (b) 1-(4-trimethylsilylbuta-1,3-diynyl)-2-(triisopropylsilyl-ethynyl)benzene, aq KOH, $\text{PdCl}_2(\text{PPh}_3)_2$, CuI, Et_3N , THF; (c) Bu_4NF , MeOH, THF; (d) $\text{Cu}(\text{OAc})_2$, CuCl, pyridine

Figure 3 Molecular structure of **2** (left); ellipsoids drawn at the 30% level. Side view of two molecules (right). THF solvate molecules omitted for clarity

Figure 4 ^1H NMR (300 MHz, CD_2Cl_2) spectra of the free hybrid **2** (top) and of the 1:1 pseudorotaxane **2·3** (bottom)

ether moiety. The bottom spectrum shows a number of changes upon addition of an equimolar amount of **3**, all of which support strong complexation. The most noticeable differences are in the crown ether region, where the central ethylene protons (H^g) change from a singlet in free **2** to two multiplets in pseudorotaxane **2·3** (Figure 5) with an upfield shift of ca. 0.25 ppm. Crown ether protons H^e and H^f show increased separations between the two sets of multiplets as well as slight upfield shifts. The resonances in the aromatic region also exhibit notable changes. The broad singlet attributable to the four H^b protons in **2** splits into the expected AA'BB' multiplet with an upfield shift of ca. 0.1 ppm once the complex is formed. The singlet for

H^a shifts upfield by 0.15 ppm. Resonances of the ammonium ion were observed at $\delta = 7.57$ (broad singlet), 7.32–7.13 (multiplets), and 4.74 (broad triplet) representing H^3 , H^1 , and H^2 , respectively. Based on the large number of previous pseudorotaxane studies,¹³ the observed chemical shift changes, as well as there being no indication of free **2**, are consistent with strong complexation between the host and guest molecules.

The supramolecular complex **2·3** was also analyzed by FAB mass spectrometry. Although the spectrum did not show the parent peak of m/z 1085, peaks corresponding to $\text{M}^+ - \text{PF}_6^-$ and hybrid **2** were observed at 940 and 742 mass

Figure 5 The structures of 1:1 pseudorotaxane **2·3** and 2:1 pseudorotaxane **2₂·4**

units, respectively. Attempts to obtain single crystals suitable for X-ray diffraction have been unsuccessful to date.

The complexation between DBA **2** and the bis-ammonium salt **4** was also performed in CD_2Cl_2 by mixing a 2:1 ratio of host and thread molecules (Figure 5). The results of this experiment are considerably more complicated. ^1H NMR spectra indicated the presence of a mixture of pseudorotaxane complexes and their free components in solution. The FAB mass spectrum showed very small peaks of the 2:1 complex corresponding to $\text{M}^+ - \text{PF}_6^-$, $\text{M}^+ - 2\text{PF}_6^-$, and crown ether **2** containing a sodium ion. Further efforts to decipher these results are in progress.

In conclusion, we have demonstrated an efficient synthesis of crown ether containing dehydrobenzoannulene **2** and its ability to complex secondary dialkylammonium ions. The formation of a pseudorotaxane from a 1:1 mixture of **2** and **3** is facile in CD_2Cl_2 , according to ^1H NMR spectroscopic data. The formation of the corresponding [3]pseudorotaxane from a 2:1 mixture of **2** and **4** was incomplete under analogous conditions. Synthetic efforts towards making bis- and tris-crown ether-fused annulenes are underway and experiments to probe the formation of more complicated supermolecules will be reported in due course.

Reagents and instrumentation used have been described previously.^{8g}

4,5-Diiododibenzo-24-crown-8 (**6**)

A suspension of CsCO_3 (18.4 g, 56.5 mmol) in anhyd MeCN (250 mL) was placed under N_2 and brought to reflux. To this was added ditosylate **5**¹⁴ (7.72 g, 11.3 mmol) and 4,5-diiodocatechol¹⁵ (4.09 g, 11.3 mmol) dissolved in anhyd MeCN (250 mL) over a 24 h period. The mixture was refluxed for an additional 48 h and then cooled. The suspension was filtered and concentrated in vacuo. The residue was dissolved in CH_2Cl_2 (150 mL) and washed with aq sat. NaHCO_3 solution (150 mL). The aqueous layer was extracted with CH_2Cl_2 (2×100 mL) and the combined organics were dried (MgSO_4), filtered, and concentrated. Chromatography on silica gel (hexanes–EtOAc, 1:1) gave **6** (3.09 g, 39%) as a white solid; mp 91.7–93.6 °C.

IR (CDCl_3): 3062, 2922, 2870, 1592, 1249, 1127 cm^{-1} .

^1H NMR (CDCl_3): δ = 7.24 (s, 2 H), 6.92–6.84 (m, 4 H), 4.16–4.12 (m, 4 H), 4.10–4.06 (m, 4 H), 3.93–3.86 (m, 8 H), 3.84–3.77 (m, 8 H).

^{13}C NMR (CDCl_3): δ = 149.44, 148.85, 123.89, 121.39, 113.97, 96.47, 71.35, 71.23, 69.91, 69.64, 69.56, 69.32.

MS (APCI): m/z (%) = 718 (100, $\text{M}^+ + \text{H}_2\text{O}$), 701 (5, $\text{M}^+ + \text{H}$).

Anal. Calcd for $\text{C}_{24}\text{H}_{30}\text{I}_2\text{O}_8$ (700.30): C, 41.16; H, 4.32. Found: C, 40.98; H, 4.23.

α,ω -Polyynes **7**

The diiodide **6** (981 mg, 1.4 mmol) and 1-(4-trimethylsilylbuta-1,3-diynyl)-2-(triisopropylsilylethynyl)benzene^{8g} (1.59 g, 4.2 mmol) were dissolved in H_2O –THF– Et_3N (0.01:1:5, 20 mL) in separate vessels. The solutions were degassed vigorously by three freeze-pump-thaw cycles. The acetylene solution was added via syringe pump to the solution of **6** charged with $\text{PdCl}_2(\text{PPh}_3)_2$ (35 mg, 0.05 mmol), CuI (15 mg, 0.08 mmol), and KOH (2.36 g, 42 mmol) over 12 h at 50 °C under N_2 . Upon completion, the mixture was concen-

trated in vacuo and redissolved in CH_2Cl_2 . The dark residue was filtered through a thin cake of silica gel using CH_2Cl_2 and concentrated. Chromatography on silica gel (hexanes–THF, 3:1) gave polyyne **7** (1.21 g, 81%) as a tan gum.

IR (neat): 2946, 2866, 2415, 2203 cm^{-1} .

^1H NMR (CDCl_3): δ = 7.54–7.45 (m, 4 H), 7.33–7.21 (m, 4 H), 6.92 (s, 2 H), 6.89 (br s, 4 H), 4.19–4.11 (m, 8 H), 3.96–3.91 (m, 8 H), 3.84 (br s, 8 H), 1.16 (s, 42 H).

^{13}C NMR (CDCl_3): δ = 149.39, 148.84, 132.66, 132.28, 128.53, 127.86, 127.18, 124.91, 121.37, 118.56, 116.85, 113.91, 104.56, 96.06, 81.31, 80.55, 78.06, 77.11, 71.47, 71.29, 69.94, 69.52, 69.43, 69.36, 18.70, 11.30.

HRMS: m/z Calcd for $\text{C}_{66}\text{H}_{80}\text{O}_8\text{Si}_2$ (M^+): 1056.5392. Found: 1056.5386.

Crown Ether/Annulene Hybrid **2**

Polyyne **7** (850 mg, 0.78 mmol) was dissolved in THF–MeOH (3:1, 20 mL) and treated with Bu_4NF (2 mL, 1 M THF solution, 2.0 mmol) at r.t. The reaction was monitored by TLC and was complete within 30 min. The mixture was diluted with Et_2O (30 mL), washed with H_2O (3×20 mL) and brine (2×20 mL), and then dried (MgSO_4). The organic layer was concentrated in vacuo and the resultant oil was redissolved in pyridine (10 mL). The pyridine solution was added over 12 h via syringe pump to a flask charged with $\text{Cu}(\text{OAc})_2$ (3.54 g, 19.5 mmol), CuCl (2.10 g, 15.6 mmol), and pyridine (200 mL) at 60 °C. The addition was done under house air and stirred further for 8 h. Upon completion, the mixture was concentrated in vacuo and then redissolved in CH_2Cl_2 . The mixture was filtered through a thin cake of silica gel using CH_2Cl_2 and concentrated. Chromatography on silica gel (hexanes–THF, 3:1) afforded **2** (320 mg, 55%) as a light yellow solid; mp 230 °C (dec.).

IR (KBr): 2926, 2866, 2211, 2192, 2140 cm^{-1} .

^1H NMR (CD_2Cl_2): δ = 7.74–7.66 (m, 4 H), 7.49–7.42 (m, 4 H), 7.13 (s, 2 H), 6.89 (br s, 4 H), 4.19–4.15 (m, 4 H), 4.12–4.08 (m, 4 H), 3.92–3.82 (m, 8 H), 3.77 (br s, 8 H).

^{13}C NMR (CD_2Cl_2): δ = 150.47, 129.54, 133.35, 133.15, 129.61, 129.40, 125.63, 125.29, 121.89, 118.97, 116.84, 114.69, 81.76, 81.32, 80.96, 78.49, 78.16, 77.00, 71.83, 71.63, 70.41, 70.00, 69.97, 69.76.

MS (FAB): m/z = 742.2 (100, M^+).

Anal. Calcd for $\text{C}_{48}\text{H}_{38}\text{O}_8 \cdot \text{C}_4\text{H}_8\text{O}$ (814.93): C, 76.64; H, 5.69. Found: C 76.54, H 5.65.

Pseudorotaxane **2:3**

CD_2Cl_2 (0.7 mL) was added to hybrid **2** (11 mg, 0.015 mmol) and dibenzylammonium hexafluorophosphate (4.8 mg, 0.014 mmol). The reaction mixture was sonicated for 3 min and then filtered through a plug of glass wool into an NMR tube.

^1H NMR (CD_2Cl_2): δ = 7.76–7.69 (m, 4 H), 7.58 (br s, 2 H), 7.50–7.44 (m, 4 H), 7.36–7.27 (m, 4 H), 7.25–7.17 (m, 6 H), 6.88 (BB'm, 2 H), 6.72 (AA'm, 2 H), 4.68–4.62 (m, 2 H), 4.16–4.13 (m, 2 H), 4.05–4.02 (m, 2 H), 3.89–3.86 (m, 2 H), 3.74–3.71 (m, 2 H), 3.60–3.55 (m, 2 H), 3.53–3.49 (m, 2 H).

MS (FAB): m/z = 940.3 (82, $\text{M}^+ - \text{PF}_6^-$), 742.2 (13, $\text{M}^+ - \text{Bz}_2\text{NH}_2\text{PF}_6^-$).

Acknowledgements

We thank the National Science Foundation (CHE-0104854 to M.M.H. and CHE-9910199 to J.F.S.) for support of this research. J.J.P. acknowledges the ACS Division of Organic Chemistry for a DOC Graduate Fellowship (1998–1999). We thank A. J. Boydston for his help with the acquisition of some characterization data.

References

- (1) Current address: Department of Chemistry, Idaho State University, Pocatello, ID 83209, USA.
- (2) Haley, M. M.; Pak, J. J.; Brand, S. C. *Top. Curr. Chem.* **1999**, *201*, 81.
- (3) Haley, M. M.; Wan, W. B. In *Advances in Strained and Interesting Organic Molecules*, Vol. 8; Halton, B., Ed.; JAI Press: New York, **2000**, 1–41.
- (4) *Inter alia*: (a) Baughman, R. H.; Galvao, D. S. *Nature* **1993**, *365*, 635. (b) He, C. B.; Liu, P. W.; Griffin, A. C. *Macromolecules* **1998**, *31*, 3145. (c) Grima, J. N.; Evans, K. E. *Chem. Commun.* **2000**, 1531.
- (5) For relevant acetylene-containing examples, *inter alia*: (a) Gardner, G. B.; Venkataraman, D.; Moore, J. S.; Lee, S. *Nature* **1995**, *374*, 792. (b) Coates, G. W.; Dunn, A. R.; Henling, L. M.; Dougherty, D. A.; Grubbs, R. H. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 248. (c) Kiang, Y.-H.; Gardner, G. B.; Lee, S.; Xu, Z.; Lobkovsky, E. B. *J. Am. Chem. Soc.* **1999**, *121*, 8204. (d) Kolotuchin, S. V.; Thiessen, P. A.; Fenlon, E. E.; Wilson, S. R.; Loweth, C. J.; Zimmerman, S. C. *Chem.–Eur. J.* **1999**, *5*, 2537. (e) Kiang, Y.-H.; Gardner, G. B.; Lee, S.; Xu, Z. *J. Am. Chem. Soc.* **2000**, *122*, 6871. (f) Xu, Z.; Lee, S.; Kiang, Y.-H.; Mallik, A. B.; Tsomaia, N.; Mueller, K. T. *Adv. Mater.* **2001**, *13*, 637. (g) See also Refs.^{11,12}
- (6) Baldwin, K. P.; Matzger, A. J.; Scheiman, D. A.; Tessier, C. A.; Vollhardt, K. P. C.; Youngs, W. J. *Synlett* **1995**, 1215.
- (7) Enkelmann, V. *Adv. Polym. Sci.* **1984**, *63*, 91.
- (8) (a) Boese, R.; Matzger, A. J.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1997**, *119*, 2052. (b) Haley, M. M.; Bell, M. L.; English, J. J.; Johnson, C. A.; Weakley, T. J. R. *J. Am. Chem. Soc.* **1997**, *119*, 2956. (c) Tovar, J. D.; Jux, N.; Jarroson, T.; Khan, S. I.; Rubin, Y. *J. Org. Chem.* **1997**, *62*, 3432. (d) Correction: Tovar, J. D.; Jux, N.; Jarroson, T.; Khan, S. I.; Rubin, Y. *J. Org. Chem.* **1998**, *63*, 4856. (e) Bunz, U. H. F.; Enkelmann, V. *Chem.–Eur. J.* **1999**, *5*, 263. (f) Pak, J. J.; Weakley, T. J. R.; Haley, M. M. *J. Am. Chem. Soc.* **1999**, *121*, 8182. (g) Bell, M. L.; Chiechi, R. C.; Johnson, C. A.; Kimball, D. B.; Matzger, A. J.; Wan, W. B.; Weakley, T. J. R.; Haley, M. M. *Tetrahedron* **2001**, *57*, 3507.
- (9) (a) Fyfe, M. C. T.; Stoddart, J. F. *Acc. Chem. Res.* **1997**, *30*, 393. (b) Lehn, J.-M. *Supramolecular Chemistry*; VCH: Weinheim, **1995**. (c) *Comprehensive Supramolecular Chemistry*; Atwood, J. L.; Davies, J. E. D.; MacNicol, D. D.; Vögtle, F., Eds.; Pergamon: Oxford, **1996**. (d) Schneider, H.-J.; Yatsimirsky, A. *Principles and Methods in Supramolecular Chemistry*; Wiley: Chichester, **2000**. (e) Desiraju, G. R. *Nature* **2001**, *412*, 397.
- (10) *Inter alia*: (a) Lindsey, J. S. *New J. Chem.* **1991**, 153. (b) Philp, D.; Stoddart, J. F. *Synlett* **1991**, 445. (c) Whitesides, G. M.; Mathias, J. P.; Seto, C. T. *Science* **1991**, *154*, 1312. (d) Lawrence, D. S.; Jiang, T.; Levett, M. *Chem. Rev.* **1995**, *95*, 2229. (e) Amabilino, D. B.; Stoddart, J. F. *Chem. Rev.* **1995**, *95*, 2725. (f) Philp, D.; Stoddart, J. F. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 1154. (g) Jäger, R.; Vögtle, F. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 930. (h) Stang, P. J.; Olenyuk, B. *Acc. Chem. Res.* **1997**, *30*, 502. (i) Conn, M. M.; Rebek, J. Jr. *Chem. Rev.* **1997**, *97*, 1647. (j) Linton, B.; Hamilton, A. D. *Chem. Rev.* **1997**, *97*, 1669. (k) Fujita, M. *Chem. Soc. Rev.* **1998**, *27*, 417. (l) Breen, T. L.; Tien, J.; Oliver, S. R. J.; Hadzie, T.; Whitesides, G. M. *Science* **1999**, *284*, 948. (m) Breault, G. A.; Hunter, C. A.; Mayers, P. C. *Tetrahedron* **1999**, *55*, 5265. (n) Lindoy, L. F.; Atkinson, I. M. *Self-Assembly in Supramolecular Systems*, In *Monographs in Supramolecular Chemistry*; Stoddart, J. F., Ed.; Royal Society of Chemistry: Cambridge, **2000**. (o) Balzani, V.; Credi, A.; Raymo, F. M.; Stoddart, J. F. *Angew. Chem. Int. Ed.* **2000**, *39*, 3348. (p) Zimmerman, S. C.; Lawless, L. J. *Top. Curr. Chem.* **2001**, *217*, 95.
- (11) Xiao, J.; Yang, M.; Lauher, J. W.; Fowler, F. W. *Angew. Chem. Int. Ed.* **2000**, *39*, 2132.
- (12) Sarkar, A.; Okada, S.; Matsuzawa, H.; Matsuda, H.; Nakanishi, H. *J. Mater. Chem.* **2000**, *10*, 819; and references cited therein.
- (13) *Inter alia*: (a) Ashton, P. R.; Chrystal, E. J. T.; Glink, P. T.; Menzer, S.; Schiavo, C.; Spencer, N.; Stoddart, J. F.; Tasker, P. A.; White, A. J. P.; Williams, D. J. *Chem.–Eur. J.* **1996**, *2*, 709. (b) Glink, P. T.; Schiavo, C.; Stoddart, J. F.; Williams, D. J. *Chem. Commun.* **1996**, 1483. (c) Glink, P. T.; Stoddart, J. F. *Pure Appl. Chem.* **1998**, *70*, 419. (d) Fyfe, M. C. T.; Stoddart, J. F. *Adv. Supramol. Chem.* **1999**, *5*, 1. (e) Cantrill, S. J.; Pease, A. R.; Stoddart, J. F. *J. Chem. Soc., Dalton Trans.* **2000**, 3715. (f) Kolchinski, A. G.; Busch, D. H.; Alcock, N. W. *J. Chem. Soc., Chem. Commun.* **1995**, 1289. (g) Hubin, T. J.; Kolchinski, A. G.; Vance, A. L.; Busch, D. H. *Adv. Supramol. Chem.* **1995**, *5*, 237. (h) Yamaguchi, N.; Gibson, H. W. *Angew. Chem. Int. Ed.* **1999**, *38*, 143. (i) Takata, T.; Kihara, N. *Rev. Heteroatom Chem.* **2000**, *22*, 197.
- (14) Cantrill, S. J.; Youn, G. J.; Stoddart, J. F.; Williams, D. J. *J. Org. Chem.* **2001**, *66*, 6857.
- (15) Kinder, J. D.; Youngs, W. J. *Organometallics* **1996**, *15*, 460.
- (16) Crystal data for **2**: C₄₈H₃₈O₈·C₄H₈O, *M_r* = 814.93, yellow block, 0.22 × 0.31 × 0.41 mm, monoclinic, space group I2/a, *a* = 31.788(4), *b* = 8.6951(13), *c* = 35.273(4) Å, β = 113.07(1)°, *V* = 8970(2) Å³, *Z* = 8, ρ_{calc} = 1.207 g cm⁻³, MoK_α radiation (λ = 0.71073 Å), μ = 0.82 cm⁻¹, *F*(000) = 3440, *T* = 23 °C, 2θ_{max} = 23.5°, 13240 independent reflections scanned, 6623 independent reflections in refinement, 550 parameters, *R*(*F*) = 0.050 [*I* σ(*I*)], *wR*(*F*²) = 0.068 (all data). Data were obtained on an Enraf-Nonius CAD-4 Turbo diffractometer. Structure refinement (C, O atoms anisotropic, H atoms riding) was accomplished with teXsan (v. 1.7 for SGI workstations). Crystallographic data (excluding structure factors) for **2** have been deposited with the Cambridge Crystallographic Data Center as supplementary publication no. CCDC-182774.