Gastrobronchial fistula is an extremely rare occurrence. Most cases result from postoperative complications of esophago-gastric surgery [1] and rarely from traumatic injury [2], subphrenic abscess [3], perforating peptic ulcer [4], or malignancies [5]. We observed the delayed development of a gastrobronchial fistula after massive caustic injury of the stomach due to suicidal lye ingestion.

A 32-year-old woman was admitted after suicidal ingestion of about 100 ml of a 25% ammonia solution. Plain films of the chest and abdomen revealed no perforation. From the 5th day chest roentgenograms consistently documented a left-sided opacification that was sonographically echogenic and was located subphrenically. Endoscopic surveillance revealed deep ulcerations in the midbody and fundus which were covered by fibrinoid membranes which sloughed incrementally. After 10 weeks the fundus appeared thin and translucent. Recurring high-grade fever occurred in the 11th

week. A computerized tomography scan revealed a subphrenic abscess. A study with water-soluble contrast material revealed a passage from the fundus into the subphrenic cavity and into the bronchial system (Figure 1). At this time the patient rejected the proposed surgical procedure. With antibiotic coverage and parenteral nutrition the subphrenic abscess completely regressed, while the fistula enlarged. In the 16th week bronchial structures could be identified on gastroscopy (Figure 2). In the 20th week the fistula was closed by a subphrenic net seal and a partial resection of the fundus.

Complications of lye ingestion are typically esophageal and antral strictures as well as perforations. In this case, massive necrosis of the fundus resulted in a subphrenic inflammation. Secondly, with a delay of 10 weeks, this inflammation caused perforation of the fundus and the diaphragm [3]. Because of the patient's refusal to undergo prompt surgical therapy, this case also demonstrates that gas-

trobronchial fistula due to subphrenic abscess can be treated conservatively without surgery over several weeks.

E. A. Purucker, S. Südfeld, S. Matern

Department of Internal Medicine III,
Medical Faculty, Technical University,
Aachen, Germany

References

- ¹ Stal JM, Hanly PJ, Darling GE. Gastrobronchial fistula: an unusual complication of esophagectomy. *Ann Thorac Surg* 1994; 58: 886–887
- ² Al-Qudah A. Traumatic gastrobronchial fistula: case report and review of the literature. *J Pediatr Surg* 1997; 32: 1798–1800
- ³ Angelillo VA, O'Donohue Jr WJ, Campbell JC et al. Gastrobronchial fistula secondary to a subphrenic abscess. *Chest* 1983; 84: 85–86
- ⁴ Matsuoka T, Nagai Y, Muguruma K et al. Liver penetration and gastrobronchial fistula: unusual complications of a peptic ulcer. *Am Surg* 1995; 61: 492–494
- ⁵ Cameron EW, Colby JM, Swanson RS. Gastrobronchial fistula in untreated lymphoma. *J Thorac Imaging* 1996; 11: 150–152

Corresponding Author

E. A. Purucker

Department of Internal Medicine III
Medical Faculty
Technical University
Pauwelsstrasse 30
52057 Aachen
Germany

Fax: +49-241-8082455

E-mail: epurucker@ukaachen.de

Figure 1 Upper gastrointestinal contrast study demonstrating a subphrenic cavity and bronchial structures.

Figure 2 Endoscopic view of bronchial structures through a wide fistula in the gastric fundus.