SYNLETT Spotlight 88

MgBr₂·OEt₂ – A Versatile Reagent in Organic Synthesis

Compiled by Suresh Kumar Tipparaju

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Suresh Kumar Tipparaju obtained his BSc from Osmania University and MSc in chemistry from Sri Sathya Sai Institute of Higher Learning, Prashanthinilayam in 1998. He has been working on 'stereoselective functionalizations on arene tricarbonylchromium template' in Prof. Amitabha Sarkar's group at National Chemical Laboratory. His research interests are in metal-mediated organic synthesis, asymmetric synthesis, and the chemistry of hypervalent silicon.

Division of Organic Chemistry (Synthesis), National Chemical Laboratory, Pune – 411 008, India
Fax +91(20)5893153; E-mail: tsuresh_kumar@yahoo.com

Introduction

Use of magnesium(II) species as Lewis acid catalysts for various functional group transformations is well documented.¹ Of these, magnesium halides are the most useful. Ready availability and ease of preparation prompted the frequent use of MgBr₂·OEt₂ in various organic transformations. The oxophilic and coordinating nature of MgBr₂·OEt₂ has been demonstrated through its use as a bidentate chelating Lewis acid in a number of chelation-controlled reactions such as cycloadditions,² asymmetric aldol reactions,³ rearrangements,⁴ radical additions,^{5,6}

hydrogen transfer reactions,⁷ stereoselective reductions,⁸ and anomerizations.⁹

MgBr₂·OEt₂ is commercially available as a grey solid (mp >300 °C, fp 35 °C). It can be readily prepared by reacting a slight excess of magnesium turnings with 1,2-dibromoethane in anhydrous diethyl ether.¹⁰ The solution can be stored at room temperature for several months and the solid can be stored in a vacuum desiccator for indefinite periods without any loss of activity.

Abstracts

(A) Condensation of 2,5-dimethoxy-2,5-dihydrofuran with ethyl vinyl ether in the presence of a catalytic amount of MgBr₂·OEt₂ resulted in 2-furylacetaldehyde diethyl acetal in 50% yield. The reaction is a formal acetal and ethyl vinyl ether condensation followed by aromatization. The protocol has been employed in the synthesis of various 2-(2-furo)tetrahydrofuranic or -pyranic moieties in good yields.¹¹

(B) A mild and practical N-acylation of amides was possible by the dual activation of both amides and acid anhydrides with MgBr₂·OEt₂.¹² The method was applicable to amides that can undergo O-acylation and are susceptible to racemization or O,N-acyl migrations.

(C) We have recently reported a mild method for the formation of cyclic siloxanes by the exchange of the Li counterion of an intermediate alkoxide with Mg, using excess MgBr₂·OEt₂.¹³ The reaction may formally be considered to be a *semi*-Brook rearrangement.

(D) MgBr₂·OEt₂, in combination with Bu₃SnH, was effective in a chelation-controlled reductive opening of methoxybenzylidene acetals.¹⁴ The reaction offers a mild and efficient method for selective mono-MPM ether protection of diols. High conversions, regioselectivity and tolerance to functional groups make this a very useful protocol in natural product synthesis.

(E) MgBr₂·OEt₂ effected deprotection of aliphatic SEM ethers under extremely mild and high yielding conditions in the presence of other sensitive groups like acetonides, TBS and TIPS ethers and O-silylated cyanohydrins.¹⁵ A variety of functionalities including alcohols, esters, benzyl groups, dithianes, and methoxy acetals are tolerated.

(F) The $\text{MgBr}_2 \cdot \text{OEt}_2$ - Me_2S system was used for a mild and chemo-selective deprotection of *p*-methoxybenzyl (PMB) ethers in the presence of 1,3-diene, *t*-butyldimethylsilyl (TBDMS) ether, benzoate, benzyl ether, and acetonides.¹⁶ The method is especially effective for 1,3-diene systems that tend to isomerize rapidly when other protocols are employed.¹⁷

(G) A stereodivergent opening of the oxirane ring with $\text{MgBr}_2 \cdot \text{OEt}_2$ was recently described.¹⁸ While $\text{MgBr}_2 \cdot \text{OEt}_2$ alone resulted in the *anti*-bromohydrin in high diastereomeric excess, use of $\text{MgBr}_2 \cdot \text{OEt}_2$ /Amberlyst 15 gave the *syn*-product in high yields.

(H) The efficacy of $\text{MgBr}_2 \cdot \text{OEt}_2$ as a chelating Lewis acid in highly diastereoselective addition of nucleophiles¹⁹ to $\text{Cr}(\text{CO})_3$ -complexed aryl aldehydes was recently demonstrated by us.^{19a} The results indicated that $\text{MgBr}_2 \cdot \text{OEt}_2$ can form an effective seven-membered chelate.

(I) $\text{MgBr}_2 \cdot \text{OEt}_2$ could afford 3-hydroxyazetidines by a highly regio- and stereoselective cyclization of 2,3-epoxy amines.²⁰

References

- (1) For some lead references see: (a) Li, W.-D. Z.; Zhang, X.-X. *Org. Lett.* **2002**, *4*, 3485. (b) Marshall, J. A. *Chem. Rev.* **1996**, *96*, 31.
- (2) (a) Tamura, O.; Kuroki, T.; Sakai, Y.; Takizawa, J.; Yoshino, J.; Morita, Y.; Mita, N.; Gotanda, K.; Sakamoto, M. *Tetrahedron Lett.* **1999**, *40*, 895; and ref. 8 therein. (b) Kashima, C.; Fukusaka, K.; Takahashi, K.; Yokoyama, Y. *J. Org. Chem.* **1999**, *64*, 1108. (c) Oblin, M.; Pons, J.-M.; Parrain, J.-L.; Rajzmann, M. *Chem. Commun.* **1998**, 1619. (d) Zemribo, R.; Romo, D. *Tetrahedron Lett.* **1995**, *36*, 4159.
- (3) (a) Fujisawa, H.; Sasaki, Y.; Mukaiyama, T. *Chem. Lett.* **2001**, 190. (b) Kiyooka, S.; Shahid, K. A.; Hena, M. A. *Tetrahedron Lett.* **1999**, *40*, 6447. (c) Swiss, K. A.; Choi, W. B.; Liotta, D. C.; Abdel-Magid, A. F.; Maryanoff, C. A. *J. Org. Chem.* **1991**, *56*, 5978.
- (4) Black, T. H.; McDermott, T. S.; Brown, G. A. *Tetrahedron Lett.* **1991**, *32*, 6501; and ref. 4 and 7 therein.
- (5) (a) Sibi, M. P.; Sausker, J. B. *J. Am. Chem. Soc.* **2002**, *124*, 984. (b) Hayen, A.; Koch, R.; Saak, W.; Haase, D.; Metzger, J. O. *J. Am. Chem. Soc.* **2000**, *122*, 12458.
- (6) (a) Enholm, E. J.; Lavieri, S.; Cordóva, T.; Ghiviriga, I. *Tetrahedron Lett.* **2003**, *44*, 531; and ref. 3 therein. (b) Nagano, H.; Toi, S.; Matsuda, M.; Hirasawa, T.; Hirasawa, S.; Yajima, T. *J. Chem. Soc., Perkin Trans. I* **2002**, 2525; and ref. 4 therein.
- (7) Guindon, Y.; Liu, Z.; Jung, G. *J. Am. Chem. Soc.* **1997**, *119*, 9289.
- (8) (a) Guanti, G.; Banfi, L.; Riva, R.; Zannetti, M. T. *Tetrahedron Lett.* **1993**, *34*, 5483. (b) Kawate, T.; Nakagawa, M.; Kakikawa, T.; Hino, T. *Tetrahedron: Asymmetry* **1992**, *3*, 227.
- (9) Mukaiyama, T.; Takeuchi, K.; Uchiro, H. *Chem. Lett.* **1997**, 625.
- (10) For crystal structure and other details see: Ecker, A.; Ueffing, C.; Schnoeckel, H. *Chem.-Eur. J.* **1996**, *2*, 1112.
- (11) Malanga, C.; Mannucci, S. *Tetrahedron Lett.* **2001**, *42*, 2023.
- (12) Yamada, S.; Yaguchi, S.; Matsuda, K. *Tetrahedron Lett.* **2002**, *43*, 647.
- (13) Tipparaju, S. K.; Mandal, S. K.; Sur, S.; Puranik, V. G.; Sarkar, A. *Chem. Commun.* **2002**, 1924.
- (14) Zheng, B.-Z.; Yamauchi, M.; Dei, H.; Kusaka, S.; Matsui, K.; Yonemitsu, O. *Tetrahedron Lett.* **2000**, *41*, 6441.
- (15) Vakalopoulos, A.; Hoffmann, H. M. R. *Org. Lett.* **2000**, *2*, 1447.
- (16) Onoda, T.; Shirai, R.; Iwasaki, S. *Tetrahedron Lett.* **1997**, *38*, 1443.
- (17) For use of $\text{MgBr}_2 \cdot \text{OEt}_2$ in deprotection of MOM, MTM, and SEM ethers see: Kim, S.; Kee, I. S.; Park, Y. H.; Park, J. H. *Synlett* **1991**, 183.
- (18) Lupattelli, P.; Bonini, C.; Caruso, L.; Gambacorta, A. *J. Org. Chem.* **2003**, *68*, 3360; and references cited therein.
- (19) (a) Tipparaju, S. K.; Puranik, V. G.; Sarkar, A. *Org. Biomol. Chem.* **2003**, *1*, 1720. (b) Ward, D. E.; Hrapchak, M. J.; Sales, M. *Org. Lett.* **2000**, *2*, 57. (c) Minassian, F.; Pelloux-Leon, N.; Vallee, Y. *Synlett* **2000**, 242. (d) Kornienko, A.; d'Alarcao, M. *Tetrahedron Lett.* **1997**, *37*, 6497. (e) Banfi, L.; Guanti, G.; Zannetti, M. T. *J. Org. Chem.* **1995**, *60*, 7870. (f) Panek, J. S.; Cirillo, P. F. *J. Org. Chem.* **1993**, *58*, 999.
- (20) Karikomi, M.; Arai, K.; Toda, T. *Tetrahedron Lett.* **1997**, *34*, 6059.