Endoclips as Nidus for Choledocholithiasis Presenting 5 Years After Laparoscopic Cholecystectomy

S. Khanna, J. C. Vij

Department of Gastroenterology,
Pushpawati Singhanian Research Institute
for Liver, Renal and Digestive Diseases,
New Delhi, India

Corresponding Author

S. Khanna, MD

Pushpawati Singhanian Research Institute
for Liver, Renal and Digestive Diseases,
Shiekh Sarai-Phase II
New Delhi 110019

India

Fax: + 91-11-29250548

E-mail: khannasudeep@hotmail.com

Figure 1 a,b A middle-aged lady presented to our unit with progressively worsening jaundice of 5 days' duration. There had been no preceding prodrome. Based on the clinical features and liver function tests she was diagnosed as having surgical obstructive jaundice. Ultrasound showed dilated intrahepatic biliary radicals and a common bile duct (CBD) dilated as far as the lower end. The patient had undergone laparoscopic cholecystectomy for gall-

stone disease 5 years previously. On endoscopic retrograde cholangiography it was found that she had a stone, for which surgical clips had served as a nidus, at the lower end of the CBD. The stone was removed endoscopically and the patient improved. The migration of surgical clips as a complication of laparoscopic cholecystectomy is a rare but increasingly diagnosed complication.