SYNLETT Spotlight 111

HOF·CH₃CN Complex

Compiled by Siddhartha Gogoi

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Siddhartha Gogoi was born in Sivasagar, Assam (India) in 1977. He completed his B.Sc. (1998) degree from Dibrugarh University and M.Sc. (2000) degree in chemistry from Gauhati University, Guwahati (India). He is currently working toward a Ph.D. in organic chemistry under the supervision of Dr. N. C. Barua at the Regional Research Laboratory in Jorhat, India. His research interests are the total synthesis of bioactive natural products and the development of new methodologies.

Natural Products Chemistry Division, Regional Research Laboratory (CSIR), Jorhat-785006, Assam, India
E-mail: siddhukaku_7@yahoo.co.in

Introduction

HOF was originally synthesized by Appelman¹ in 1971, but was found to be a very unstable substance and of little use in synthetic chemistry as it was difficult to prepare and could be generated only in very minute amounts. A few years ago, Rozen developed a new way to prepare and stabilize it through complexation with acetonitrile.² Whereas neat HOF can be unpredictably explosive at temperatures above -40 °C, its acetonitrile complex (HOF·CH₃CN) is much more stable.

HOF·CH₃CN complex is a potent oxygen-transfer reagent for a variety of reactions, some of which are occasionally difficult on certain substrates. It reacts very efficiently with a large variety of organic compounds, resulting in either an oxidation or a transfer of an oxygen atom from the reagent to the substrate. This reagent can be applied to the synthesis of epoxides from various olefins, ketones from methyl ethers, esters by the Baeyer–Villiger reaction, sulfones from sulfides, nitro compounds from amines and

amino acids, N-oxides from tertiary amines, and other conversions. More recently, HOF·CH₃CN complex has been used in the direct oxidation of aliphatic and selected aromatic azides to nitro compounds, both very efficiently and in excellent yields.

Its uniqueness is based on its oxygen, which becomes a strongly electrophilic species because of it being bonded to highly electronegative fluorine. Its reactions have been shown to be ionic in nature and are usually complete in a few minutes at temperatures ranging from 0 °C to 25 °C.

Preparation:

HOF·CH₃CN complex is simply prepared in a glass reactor by bubbling nitrogen-diluted fluorine (10–15%) into aqueous acetonitrile (10% H₂O in CH₃CN) at 0 °C. The product is stable at 0 °C for a few hours.²

Scheme 1 Preparation of HOF·CH₃CN complex

Abstracts

(A) Rozen and coworkers reported the first direct oxidation of aliphatic and aromatic azides into nitro compounds with HOF·CH₃CN complex in excellent yields.³

(B) As a strong oxidant, HOF·CH₃CN complex can oxidize aromatic amines to nitroarenes in high yields.⁴ It can also oxidize the amino group of α -amino esters to α -nitro esters without affecting the aromatic ring.⁵

(C) HOF·CH₃CN complex is an ecologically friendly oxidizing agent and can oxidize alcohols⁶ and methyl ethers to ketones. Ketones also undergo Baeyer–Villiger oxidation to give esters under the same reaction conditions.⁷ All these reactions are ionic in nature.

(D) With its strong electrophilic oxygen, HOF·CH₃CN complex is an excellent oxygen transfer agent⁸ that epoxidizes a wide variety of alkenes and alkynes, including very deactivated ones that cannot be directly epoxidized by other methods.

(E) HOF·CH₃CN complex can oxidize all types of sulfides to sulfones in excellent yields in only a few minutes at or below room temperature.⁹ The reaction proceeds through the formation of sulfoxides which, at low temperatures of around –78 °C, can be isolated in good yields.

(F) HOF·CH₃CN complex can also hydroxylate¹⁰ and otherwise oxidize many aromatic compounds, though yields are moderate.

References

- (1) Studier, M. H.; Appelman, E. H. *J. Am. Chem. Soc.* **1971**, *93*, 2349.
- (2) Rozen, S.; Brand, M. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 554.
- (3) (a) Rozen, S.; Carmeli, M. *J. Am. Chem. Soc.* **2003**, *125*, 8118. (b) Surya Prakash, G. K.; Etkorn, M. *Angew. Chem., Int. Ed. Engl.* **2004**, *43*, 26.
- (4) Kol, M.; Rozen, S. *J. Chem. Soc., Chem. Commun.* **1991**, 567.
- (5) (a) Rozen, S.; Kol, M. *J. Org. Chem.* **1992**, *57*, 7342. (b) Rozen, S.; Bar-Haim, A.; Mishani, E. *J. Org. Chem.* **1994**, *59*, 1208.
- (6) Rozen, S.; Bareket, Y.; Kol, M. *Tetrahedron* **1993**, *49*, 8169.
- (7) Rozen, S.; Dayan, S.; Bareket, Y. *J. Org. Chem.* **1995**, *60*, 8267.
- (8) (a) Rozen, S.; Kol, M. *J. Org. Chem.* **1990**, *55*, 5155. (b) Rozen, S.; Bareket, Y.; Kol, M. *Tetrahedron Lett.* **1996**, *37*, 531. (c) Dayan, S.; Ben-David, I.; Rozen, S. *J. Org. Chem.* **2000**, *65*, 8816.
- (9) Rozen, S.; Bareket, Y. *J. Org. Chem.* **1997**, *62*, 1457.
- (10) (a) Rozen, S.; Brand, M.; Kol, M. *J. Am. Chem. Soc.* **1989**, *111*, 8325. (b) Kol, M.; Rozen, S. *J. Org. Chem.* **1993**, *58*, 1593. (c) Rozen, S. *Acc. Chem. Res.* **1996**, *29*, 243. (d) Rozen, S. *Pure Appl. Chem.* **1999**, *71*, 481.