SYNLETT
Spotlight 112

Oxaziridines

Compiled by Jitendra Kumar Mishra

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Jitendra Kumar Mishra was born in 1980 in Gorakhpur (U.P.), India. He completed his B.Sc. (1999) and M.Sc. (2001) degrees in organic chemistry from D.D.U. Gorakhpur University, Gorakhpur. He is currently working as a DRDO-JRF for his Ph.D. degree under the supervision of Prof. M. P. Kaushik at the Defence Research and Development Establishment in Gwalior, India.

Process Technology Development Division, Defence Research and Development Establishment, Jhansi Road, Gwalior, 474002 (MP) India

E-mail: jitendramishra16@rediffmail.com

Introduction

Oxaziridines were first discovered in 1956¹ and have been widely investigated, principally for two reasons. The presence of an inherently weak N–O bond in a strained ring promised a group of compounds of unusually high reactivity. In addition, this system possesses the structural elements that seem to be required in order to observe stereochemical isomerism at nitrogen: ring strain and an atom with unshared electron pairs attached to the nitrogen.

Oxaziridines can be used as both oxygenating and aminating agents in their reactions with a wide variety of nucleophiles. Oxaziridines with small groups on N, for example hydrogen or methyl, act as aminating agents, whereas those with bulky or electron withdrawing groups on N preferentially transfer the oxygen atom. Oxaziridine-mediated processes are of interest due to the easy accessibility of the reagents and their potential for asymmetric induction.

Abstracts

(A) For asymmetric oxygenation of chiral imide enolates:

Evans and coworkers have found that the α -hydroxylation reactions of chiral enolates derived from oxazolidinone carboximide proceed with exceptional facility with the oxidant 2-(phenylsulfonyl)-3-phenyl oxaziridine. The following carboximides were transformed into their respective Z-sodium enolates (1.2 equiv of $\text{NaN}(\text{Me}_3\text{Si})_2$, THF, -78°C) with slight excess of oxaziridine.²

(B) For asymmetric oxidation of ketone enolates:

The reagent-controlled asymmetric oxidation of trisubstituted enolate anions can be done by enantiomerically pure camphorsulfonyl oxaziridine.³

(C) For epoxidation of alkenes:

Epoxidation of alkenes are also possible by using oxaziridine.⁴

(D) For epoxidation of alkenes (with perfluorinated oxaziridines):

Alkyl-substituted olefins are epoxidized by perfluoro-*cis*-2,3-dialkyl oxaziridines under mild conditions. Similarly electron deficient substrates (e.g. α,β -enones) can also be epoxidized.⁵

(E) For amination of N-nucleophiles:

Oxaziridines can be used as an aminating reagent for N-amination of 2° amines.⁶

(F) For N-acylamidation:

Direct N-acyl transfer as a method for protected hydrazine synthesis has also been shown to proceed in good yield with a wide variety of substrates.⁷

References

- (1) Emmons, W. D. *J. Am. Chem. Soc.* **1956**, 78, 6208.
- (2) Evans, D. A.; Morrissey, M. M.; Dorow, R. L. *J. Am. Chem. Soc.* **1985**, 107, 4346.
- (3) Davis, F. A.; Sheppard, A. C.; Chen, B.-C.; Serajul Haque, M. *J. Am. Chem. Soc.* **1990**, 112, 6679.
- (4) Davis, F. A.; Abdul-Malik, N. F.; Awad, S. B.; Harakal, M. E. *Tetrahedron Lett.* **1981**, 22, 917.
- (5) Arnone, A.; Desmarteau, D. D.; Novo, V.; Petrov, V. A.; Pregmolato, M.; Resnati, G. *J. Org. Chem.* **1996**, 61, 8805.
- (6) Andrae, S.; Schmitz, E. *Synthesis* **1991**, 327.
- (7) Schmitz, E.; Fechner-Simon, H.; Schramm, S. *Liebigs Ann. Chem.* **1969**, 725, 1.