


F. A. DAVIS,* J. DENG (TEMPLE UNIVERSITY, PHILADELPHIA, USA)

Asymmetric Total Synthesis of (–)-Agelastatin A Using Sulfonimine (*N*-Sulfinyl Imine) Derived Methodologies
Org. Lett. **2005**, 7, 621–623.

Total Synthesis of (–)-Agelastatin



Significance: The target molecule was isolated from the sponge *Agelas dedromorpha*. It is active against several tumour cell lines and it inhibits glycogen synthase kinase-3 β . It has potential for the treatment of Alzheimer's disease, the prevention of neuronal apoptosis after stroke and it could function as an insulin mimetic.

Comment: The asymmetric synthesis of the 4,5-diaminocyclopenten-2-one derivative **D** entailed (a) nucleophilic addition of the lithium enolate of *N,N*-dibenzylglycine (5 equiv) to the homochiral sulfonimine **A** and (b) a ring closing metathesis of **C**. The nucleophilic addition step gave three of the four possible diastereoisomers (dr 18:1:5:0) with the major diastereoisomer **B** being the major product, isolated in 73% yield.

SYNFACTS Contributors: Philip Kocienski

Synfacts 2005, 0, 0012–0012

DOI: 10.1055/s-2005-865348; Reg-No.: K00905SF

2005 © THIEME STUTTGART • NEW YORK