A. YANAGISAWA,* T. TOUGE, T. ARAI (CHIBA UNIVERSITY, JAPAN)

Enantioselective Protonation of Silyl Enolates Catalyzed by a Binap-AgF Complex

Angew. Chem. Int. Ed. **2005**, *44*, 1546-1548.

Enantioselective Protonation of Silyl Enolates Catalyzed by a Binap-AgF Complex

Significance: A catalytic enantioselective protonation of trimethylsilyl enol ethers using AgF·binap under very mild conditions (CH_2Cl_2 –MeOH, $-20\text{ }^\circ\text{C}$ to $0\text{ }^\circ\text{C}$) is reported. Generally good to excellent yields and ee's were obtained. This method allows for preparation of enantiomerically enriched α -chiral ketones from racemic ketones by silyl enol ether formation followed by enantioselective protonation.

Comments: Cyclic trimethylsilyl enol ethers gave generally good yields and ee's using methanol as the proton source.

In the cases when $\text{R}^3 = \text{aryl}$, excellent yields and up to 99% ee's were obtained. Both 6- and 7-membered ring systems work well. However, no acyclic silyl enol ethers were reported in this communication. It was rationalized that Ag(I) acts as a Lewis acid to activate MeOH and F^- interacts with the silyl group in the protonation process. Previously, Yamamoto has reported that a chiral Lewis acid SnCl_4 /BINOL-catalyzed enantioselective protonation of silyl enol ethers and ketene disilyl acetals using 2,6-dimethylphenol as the proton source (S. Nakamura, M. Kaneeda, K. Ishihara, H. Yamamoto *J. Am. Chem. Soc.* **2000**, *122*, 8120–8130). Nakai has reported cationic Pd(II)/BINAP systems to achieve the same reactions using H_2O as the proton source (M. Sugiura, T. Nakai *Angew. Chem. Int. Ed.* **1997**, *36*, 2366–2368).

SYNFACTS Contributors: Mark Lautens, Y. Eric Fang

Synfacts 2005, 0, 0040-0040

DOI: 10.1055/s-2005-869934; Reg-No.: L00205SF

2005 © THIEME STUTTGART • NEW YORK