M. MARIGO, J. FRANZEN, T. B. POULSEN, W. ZHUANG, K. A. JØRGENSEN* (AARHUS UNIVERSITY, DENMARK)

Asymmetric Organocatalytic Epoxidation of α,β -Unsaturated Aldehydes with Hydrogen Peroxide
J. Am. Chem. Soc. **2005**, 127, 6964-6965.

Asymmetric Organocatalytic Epoxidation of α,β -Unsaturated Aldehydes

Significance: Epoxidation has long been an important synthetic transformation in synthetic chemistry. This represents the first direct formation of α,β -epoxy aldehydes which, importantly, was achieved under simple and mild conditions. Hydrogen peroxide was found to be the oxidant of choice while *t*-BuOOH, cumene hydroperoxide, and urea hydrogen peroxide gave similar results, but *m*-CPBA gave almost no conversion (< 3%).

Comment: The proposed mechanism involved formation of the imminium ion, which is nucleophilically attacked by the peroxide in 1,4-fashion. The subsequent enamine attacks the electrophilic peroxide oxygen forming the α,β -epoxy aldehyde. This type of activation of α,β -unsaturated carbonyls prompts for promising future studies for asymmetric 1,4-additions (see also: *Angew. Chem. Int. Ed.* **2005**, 44, 3703-3706).

SYNFACTS Contributors: Hisashi Yamamoto, Matthew Boxer

Synfacts 2005, 0, 0076-0076

DOI: 10.1055/s-2005-869988; Reg-No.: H01405SF

2005 © THIEME STUTTGART • NEW YORK