A 20-year-old, married, primigravida presented with a history of epigastric discomfort, nausea, and vomiting for 3 days. On the day of admission, she gave a history of vomiting up a 3–4-cm, pear-shaped, reddish-brown worm, which moved for a few seconds after it was vomited and then died. On further enquiry, the patient gave a history of consumption of vegetables that had been washed in pond water.

Upper gastrointestinal endoscopy revealed a reddish-brown, flat worm stuck in the second part of the duodenum (Figure 1, Video 1). The worm was extracted with the help of biopsy forceps under endoscopic guidance. The worm was 4 cm long, 2 cm wide, and about 0.25 cm thick, reddish-brown, flattened anteroposteriorly, and pear-shaped, with no prominent or obvious cephalic cone, and resembled the *fasciolopsis buski* fluke (Figure 2). Histopathological examination of the specimen confirmed it as *F. buski*. Routine investigations revealed a normal peripheral-smear eosinophil count. Stool examination did not reveal any *F. buski* eggs. The patient was treated with praziquantel 25 mg/kg as a single dose. The patient was asymptomatic and doing well on follow-up.

To our knowledge, this is the first report describing and illustrating the endoscopic removal of *F. buski* from the duodenum using forceps. Fascioliasis and other food-borne trematode infections are included in the list of important helminthiasis that have a great impact on human development. Current changes in global weather patterns appear to be increasingly affecting snail-borne helmin-

Figure 1 Endoscopic view of *fasciolopsis buski* stuck in the second part of the duodenum.

thiasis, which are strongly dependent on environmental factors. Fascioliasis is a good example of a parasitic disease that is emerging (or re-emerging) in many countries as a consequence of changes in both environmental and human factors [1,2]. Unfortunately, despite control programs, *F. buski* still remains a public health problem in endemic areas and in areas where it was once thought to have been controlled [3,4].

Though the disease is seen predominantly in south-east Asia, the endoscopic image of *F. buski* we have included here may be of interest to the rest of the world because of immigration, globalization, and the increased frequency of intercontinental travel.

Endoscopy_UCTN_Code_CCL_1AB_2AZ_3AZ
Endoscopy_UCTN_Code_TTT_1AO_2AN

M. Muruges¹, S. Veerendra¹, S. Madhu¹, S. Kude¹, A. D. Amrapurkar², U. U. Rathi¹, P. M. Rathi¹

¹ Department of Gastroenterology and Hepatology, BYL Nair Charitable Hospital and TN Medical College, Mumbai, India

² Department of Pathology, BYL Nair Charitable Hospital and TN Medical College, Mumbai, India.

Figure 2 The endoscopically extracted *F. buski* worm was 4 cm long, 2 cm wide, and about 0.25 cm thick.

References

- Mas-Coma S, Bargues MD, Valero MA. Fascioliasis and other plant-borne trematode zoonoses. *Int J Parasitol* 2005; 35: 1255–1278
- Bhatti HS, Malla N, Mahajan RC, Sehgal R. fasciolopsiasis – a re-emerging infection in Azamgarh (Uttar Pradesh). *Indian J Pathol Microbiol* 2000; 43: 73–76
- World Health Organization. Foodborne trematode infections. *Bull WHO* 1995a; 73: 397–399
- World Health Organization. Control of food-borne trematode infections. *WHO Tech Rep Ser* 1995b; 849: 1–157

Corresponding author

P. M. Rathi, MD

Department of Gastroenterology and Hepatology
BYL Charitable Nair Hospital and TN Medical College
Dr. AL Nair Road
Mumbai Central
Mumbai
Maharashtra 400 008
India
Fax: +91-22-23021168
Email: rathipm@hotmail.com

DOI: 10.1055/s-2006-945154

Video 1

Upper gastrointestinal endoscopy, showing live, mobile *F. buski* in the second part of the duodenum. The worm was extracted using biopsy forceps.