SYNLETT Spotlight 167

Sodium Hydrogen Sulfate: Safe and Efficient

Compiled by Eskandar Kolvari

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Eskandar Kolvari was born in Garmsar, Iran, in 1977. He received his B.Sc. in Applied Chemistry (2000) from Sharif University of Technology, Iran and his M.Sc. in Organic Chemistry (2003) from Bu-Ali Sina University, Hamadan, Iran. He is currently working towards his Ph.D. under the supervision of Professor Mohammad Ali Zolfigol at Bu-Ali Sina University. His research interests include the application of new catalysts in organic reactions.

Chemistry Department, College of Science, Bu-Ali Sina University,
65174 Hamadan, Post Box No. 4135, Iran
E-mail: kolvari@basu.ac.ir

Introduction

Although NaHSO_4 has been known for a long time, only in recent years it emerged as an efficient catalyst in organic chemistry. The new interest in this salt is due to environmental and economical considerations that prompt urgent need to redesign important chemical processes using suitable catalysts. NaHSO_4 can be used alone or supported on alumina¹ or silica gel, in solvent or under solvent-free conditions. The most often used form of it is the silica gel supported form. This catalyst promotes

various transformations like selective and regioselective protection and deprotection,^{2–9} nitration,¹⁰ nitrosation,¹¹ oxidation,¹² Beckman rearrangement,¹ synthesis of halide derivatives,^{13,14} coupling of indoles,¹⁵ and synthesis of quinazolinones.¹⁶

The advantages of using NaHSO_4 include operational simplicity, selectivity, and availability, and it is inexpensive and ecologically friendly.

Abstracts

(A) Bis- and tris(1*H*-indol-3-yl)methanes are synthesized in high yields by an electrophilic substitution reaction of indoles with carbonyl compounds under mild reaction conditions using silica-supported NaHSO_4 and Amberlyst-15.¹⁵

(B) Silica gel supported sodium hydrogen sulfate was found to be an efficient catalyst for the selective removal of the *N*-Boc protecting group from aromatic amines, keeping aliphatic *N*-Boc intact.²

(C) A combination of NaHSO_4 and NaNO_2 in the presence of wet SiO_2 was used as an effective nitrosating agent for the nitrosation of secondary amines to their corresponding nitroso derivatives under mild conditions.¹¹

(D) Different *p*-hydroxybenzyl alcohols were subjected to $\text{NaHSO}_4 \cdot \text{SiO}_2$ and it was shown that this catalyst can transform *p*-hydroxybenzyl alcohols to the corresponding *p*-hydroxybenzyl ethers and thioethers efficiently and selectively.⁶

(E) Cyclic and acyclic ketones, amides, and β -keto esters were converted to their α -brominated derivatives using $\text{NaHSO}_4 \cdot \text{SiO}_2$ in the presence of NBS and with Et_2O or CCl_4 as solvent at room temperature.¹³

(F) 1,2-Dihydroquinolines were subjected to oxidation effectively and in short reaction times with $\text{Na}_2\text{Cr}_2\text{O}_7 \cdot \text{H}_2\text{O}$ and NaHSO_4 as catalyst. The reactions proceed under mild conditions and with dichloromethane as solvent.^{12b}

(G) The reaction of ethyl glyoxylate with different heteroaromatic compounds in the presence of sodium salts was investigated. It was shown that NaHSO_4 is effective and affords Friedel–Crafts addition products in good yield under aqueous conditions.¹⁷

References

- (1) Gopalakrishnan, M.; Sureshkumar, P.; Kanagarajan, V.; Thanusu, J. *Lett. Org. Chem.* **2005**, *2*, 444.
- (2) Ravindranath, N.; Ramesh, C.; Reddy, M.; Das, B. *Adv. Synth. Catal.* **2003**, *345*, 1207.
- (3) Das, B.; Mahender, G.; Kumar, V.; Chowdhury, N. *Tetrahedron Lett.* **2004**, *45*, 6709.
- (4) Ramesh, C.; Ravindranath, N.; Das, B. *J. Org. Chem.* **2003**, *68*, 7101.
- (5) Ramesh, C.; Mahender, G.; Ravindranath, N.; Das, B. *Tetrahedron Lett.* **2003**, *44*, 1465.
- (6) Ramu, R.; Nath, N.; Reddy, M.; Das, B. *Synth. Commun.* **2004**, *34*, 3135.
- (7) Zhang, Z. *Monatsh. Chem.* **2005**, *136*, 1191.
- (8) Mahender, G.; Ramu, R.; Ramesh, C.; Das, B. *Chem. Lett.* **2003**, *32*, 734.
- (9) Breton, G. W. *J. Org. Chem.* **1997**, *62*, 8952.
- (10) (a) Zolfigol, M. A.; Madrakian, E.; Ghaemi, E. *Ind. J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.* **2001**, *40*, 1191. (b) Zolfigol, M. A.; Madrakian, E.; Ghaemi, E. *Molecules* **2001**, *6*, 614.
- (11) Zolfigol, M. A.; Madrakian, E.; Ghaemi, E.; Kiani, M. *Synth. Commun.* **2000**, *11*, 2057.
- (12) (a) Shirini, F.; Zolfigol, M. A.; Torabi, S. *Lett. Org. Chem.* **2005**, *2*, 544. (b) Damavandi, J. A.; Zolfigol, M. A.; Karimi, B. *Synth. Commun.* **2001**, *31*, 3183. (c) Zolfigol, M. A.; Sadeghi, M. M.; Mohammadpoor-Baltork, I.; Ghorbani Choghmarani, A.; Taqian-nasab, A. *Asian J. Chem.* **2001**, *13*, 887.
- (13) Das, B.; Venkateswarlu, K.; Mahender, G.; Mahender, L. *Tetrahedron Lett.* **2005**, *46*, 3041.
- (14) Das, B.; Banerjee, J.; Ravindranath, N. *Tetrahedron* **2004**, *60*, 8357.
- (15) Ramesh, C.; Banerjee, J.; Pal, R.; Das, B. *Adv. Synth. Catal.* **2003**, *345*, 557.
- (16) Das, B.; Banerjee, J. *Chem. Lett.* **2004**, *33*, 960.
- (17) Zhuang, W.; Jorgensen, K. A. *Chem. Commun.* **2002**, 1336.