SYNLETT
Spotlight 171

Benzophenone Imine

Compiled by Abel Crespo

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Abel Crespo was born in Asturias, Spain in 1979. He studied pharmacy at the University of Santiago de Compostela, Spain and received his B.Sc. in 2003. Currently he is working on his Ph.D. thesis at the Instituto de Farmacia Industrial of this same university under the supervision of Dr. E. Sotelo. His current research focuses on the development of multicomponent reactions.

Instituto de Farmacia Industrial (IFI), Facultad de Farmacia,
Universidad de Santiago de Compostela, Santiago de Compostela
15782, Spain
E-mail: qoabelcg@usc.es

Introduction

Benzophenone imine or (diphenylmethylene)amine (DPMA-H, **1**) is a valuable reagent in organic synthesis.¹ It is a commercially available liquid which is easily prepared by addition of phenylmagnesium bromide to benzonitrile followed by hydrolysis with methanol² or by reaction of benzophenone with ammonia.³

Synthetic applications of **1** have been historically related to peptide chemistry, specifically as protecting group of primary amines during the preparation of optically active α -amino acids.⁴ Used in conjunction with other anion-stabilising groups, **1** provides activation for proton abstraction. More recently, the development of highly efficient tin-free palladium-catalysed amination methodologies by the groups of Buchwald⁵ and Hartwig⁶ increased its synthetic utility as convenient ammonia surrogate in catalysed coupling reactions.

Abstracts

(A) A. de Meijere et al.⁷ have published the base-catalysed reaction of benzophenone imine (**1**) with methyl-2-chloro-2-cyclopropylideneacetate (**2**) to give 1,4-adduct **4**, which is a valuable intermediate in the synthesis of cyclopropyl- β -amino acids. The formal [2+4] cycloaddition of **1** and **2** affords substituted quinolines.

(B) DPMA-H (**1**) serves as amino-protecting group in the enantioselective synthesis of functionalized α -amino acids^{6,8} and small peptides⁹ using a chiral quaternary ammonium salt as catalyst.

(C) Benzophenone imine of glycine Wang resin (**12**) [prepared from F-moc of glycine Wang resin (**11**) by treatment with piperidine–DMF and then DPMA-H in NMP and glacial acetic acid] can be alkylated with α,ω -dihaloalkanes affording the valuable reactive intermediate **13**. Synthetic manipulation at the living group (X), cleavage of protecting fragments and resin yield the side-chain-reactive unnatural amino acids **17–19**.¹⁰

(D) DPMA-H (**1**) is employed in the Buchwald–Hartwig reaction as a convenient ammonia surrogate in the palladium- and nickel-catalysed amination of organic electrophiles **20**. The benzophenone imine adducts **21** can be isolated in pure form or cleaved directly to the corresponding primary anilines **22** under mild conditions by catalytic hydrogenation, treatment with hydroxylamine hydrochloride or a catalytic amount of HCl in wet THF.¹¹ A new series of improved catalysts for this transformation was recently documented.¹²

(a) 1 mol% Pd(OAc)₂, 1.5 mol% BINAP, 1.4 equiv Cs₂CO₃, THF, 65 °C, 16 h. (b) 0.25 mol% Pd₂(dba)₃, 0.75 mol% BINAP, 1.4 equiv NaOt-Bu, toluene, 80 °C, 13 h. (c) 2 mol% Pd(OAc)₂, 3 mol% BINAP, 1.4 equiv Cs₂CO₃, toluene, 100 °C, 6 h.

(E) Benzophenone imine (**1**) has been used in the preparation of the trifluoromethylated homoallylamine **26**, a useful starting material to achieve the synthesis of α -trifluoromethylated nitrogen heterocyclic compound **27** through an alkylation–ring-closure-metathesis (RCM)¹³ sequence.

References

- (1) Paquette, L. A. *Encyclopaedia of Reagents for Organic Synthesis*, Vol. 1; John Wiley: Chichester, UK, **1995**, 293.
- (2) Pickard, P. L.; Tolbert, T. L. *Org. Synth., Coll. Vol. V* **1973**, 520.
- (3) Verardo, G.; Giumanini, A. G.; Strazzolini, P.; Poiana, M. *Synth. Commun.* **1988**, 18, 1501.
- (4) O'Donnell, M. J. *Aldrichimica Acta* **2001**, 34, 3.
- (5) Guram, A. S.; Rennels, R. A.; Buchwald, S. L. *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 1348.
- (6) Louie, J.; Hartwig, J. F. *Tetrahedron Lett.* **1995**, 36, 3609.
- (7) Wessjohann, L.; Skattebol, L.; de Meijere, A. *J. Chem. Soc., Chem. Commun.* **1990**, 574.
- (8) Corey, E. J.; Noe, M. C.; Xu, F. *Tetrahedron Lett.* **1998**, 39, 5347.
- (9) Ooi, T.; Tayama, E.; Maruoka, K. *Angew. Chem. Int. Ed.* **2003**, 42, 579.
- (10) O'Donnell, M. J.; Alsina, J.; Scott, W. L. *Tetrahedron Lett.* **2003**, 44, 8403.
- (11) Wolfe, J. P.; Ahman, J.; Sadighi, J. P.; Singer, R. A.; Buchwald, S. L. *Tetrahedron Lett.* **1997**, 38, 6367.
- (12) Shen, Q.; Shekhar, S.; Stambuli, J. P.; Hartwig, J. F. *Angew. Chem. Int. Ed.* **2005**, 44, 1371.
- (13) Guille, S.; Ferry, A.; Billard, T.; Langlois, B. R. *J. Org. Chem.* **2003**, 68, 8932.