SYNLETT Spotlight 176

Tetra-*n*-Butylammonium Tribromide

Compiled by Valdemar B. C. Figueira

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Valdemar B. C. Figueira studied Applied Chemistry – Organic Chemistry (1998–2003) at FCT/UNL in Lisbon, Portugal. He joined the research group of Prof. John Paul Jones at the University of North Wales, Bangor, UK in 2002 as an Erasmus student, working with quartz crystal microbalances.

In 2003 he joined Prabhakar and Lobo's research group under the project 'Development of Synthetic Methodologies for Natural Algicides' where he is currently with a doctoral fellowship from Fundação para a Ciência e Tecnologia (FC&T, Lisbon, Portugal).

Secção de Química Orgânica Aplicada, Departamento de Química, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Quinta da Torre, 2829 – Monte da Caparica, Portugal

E-mail: valdemar.figueira@dq.fct.unl.pt

Introduction

Tetra-*n*-butylammonium tribromide (TBATB) is one of the most widely used organic ammonium tribromides (OATB's) for bromination^{1–3} of several organic substrates.

Because it is a stable crystalline solid, easy to handle and maintains the desired reaction stoichiometry it is looked at as a substitute of Br₂. It can also be considered as a 'greener' brominating agent⁴ as well as an in situ generator of anhydrous HBr.

Other types of reactions like acylation, oxidation and protection/deprotection of several functional groups can be performed with TBATB in catalytic amounts and under solvent-free conditions.

Preparation

Properties

Yellow-orange crystalline ionic solid, mp 75 °C (acetonitrile). IR bands at 170 and 190 cm⁻¹. Monoclinic space group C2/c structure determined by X-ray.

Abstracts

(A) Preparation of a wide variety of flavones and aurones starting from 2'-acetoxychalcones **1** in high yields (36–55% and 65–85%, respectively) was accomplished in two steps, being the first selective bromination with TBATB (70–80% for **2** and 75–85% for **3**).⁵

(B) Direct condensation of various alcohols and carboxylic acids was efficiently achieved with a catalytic amount of TBATB under solvent-free conditions at reflux. Chemoselectivity for primary alcohols was observed when secondary and phenolic alcohols were also present.⁶

(C) Selective protection of various carbonyl compounds to the corresponding 1,3-oxathiolanes was performed with catalytic amount of TBATB. Chemoselective deprotection of synthesized 1,3-oxathiolanes to the parent carbonyl compounds was also obtained with TBATB. No bromination on α -keto position, aromatic ring, allylic position or double bond was observed.⁷

(D) Gosain and Sharma described the kinetics and mechanism for oxidation of vicinal and non-vicinal diols to the corresponding aldehydes and hydroxycarbonyl compounds, respectively. Excellent yields were obtained.⁸

(E) Solvent-free, chemoselective diacylation of aldehydes was accomplished using a catalytic amount of TBATB, without side reaction yielding brominated products. Chemoselective cleavage of diacylates was also reported with TBATB when the reaction was performed in different conditions.⁹

(F) TBATB promotes tetrahydropyranylation and detetrahydropyranylation of primary, secondary and tertiary alcohols when present in 10 mol%. Mild reaction conditions make both reactions compatible with other acid-sensitive groups such as OTs, nitro and Boc, among others.¹⁰

References

- (1) Mondal, E.; Bose, G.; Khan, A. T. *Synlett* **2001**, 785.
- (2) Bose, G.; Li, Y.; Bhujarbarua, P. M.; Kalita, D.; Khan, A. T. *Chem. Lett.* **2001**, 30, 290.
- (3) Bose, U.; Chaudhuri, M. K.; Dey, D.; Dhar, S. S. *Pure Appl. Chem.* **2001**, 73, 93.
- (4) Bora, U.; Chaudhuri, M. K.; Dehury, S. K. *Curr. Sci.* **2002**, 82, 1427.
- (5) Bose, G.; Mondal, E.; Khan, A. T.; Bordoloi, M. J. *Tetrahedron Lett.* **2001**, 42, 8907.
- (6) Naik, S.; Kavala, V.; Gopinath, R.; Patel, B. K. *ARKIVOC* **2006**, (i), 119.
- (7) Mondal, E.; Sahu, P. R.; Bose, G.; Khan, A. T. *Tetrahedron Lett.* **2002**, 43, 2843.
- (8) Gosain, J.; Sharma, P. K. *Indian Acad. Sci. (Chem. Sci.)* **2003**, 115, 135.
- (9) Kavala, V.; Patel, B. K. *Eur. J. Org. Chem.* **2005**, 441.
- (10) Naik, S.; Gopinath, R.; Patel, B. K. *Tetrahedron Lett.* **2001**, 42, 7679.