Endoscopic Balloon Dilation of Ileal Stricture Due to Tuberculosis

Balloon dilation of small-bowel strictures is an attractive alternative that has been successfully used to treat ileal strictures due to Crohn's disease (1). However, there are few data on balloon dilation of ileal strictures due to tuberculosis (2).

A 45-year-old man presented with a history of recurrent episodes of subacute intestinal obstruction over a period of 18 months. A barium meal follow-through and barium enema showed narrowing in the terminal ileum. Colonoscopy revealed multiple superficial ulcerations in the cecum, with narrowing in the terminal ileum. A colonoscope could not be passed through the narrowed area. Cecal biopsies and a crush smear of biopsy tissue revealed epithelioid granulomas and acid-fast bacilli (*Mycobacterium tuberculosis*). Despite two months of antitubercular therapy, the symptoms persisted. Colonoscopy after two months showed persistence of the stricture (Figure 1). At this stage, balloon dilation was performed. Under endoscopic vision, a deflated TTS balloon (Rigiflex, Microvasive) was passed through the biopsy channel of the colonoscope and positioned within the stricture in the terminal ileum (Figure 2). The balloon was inflated with 10 ml air. Successful dilation was confirmed by the ability to move the inflated balloon to and fro within the strictured area without difficulty. After dilation a colonoscope (outer diameter 13 mm) could be passed through the stricture area. A repeat barium enema showed dilation of the stricture. The patient has been free of symptoms for the last 11 months.

Balloon dilation of a tubercular stricture is an easy and safe alternative to surgery, although the procedure is technically demanding. Patients with short and fibrous strictures, as in the present case, are amenable to balloon dilation (3), obviating the need for surgery.

D. K. Bhasin, B. C. Sharma, S. Dhavan,
A. Sethi, S. K. Sinha, K. Singh
Dept. of Gastroenterology,
Postgraduate Institute of
Medical Education and Research,
Chandigarh, India

Figure 1: Stricture in the terminal ileum (arrow).

Figure 2: Inflated balloon in the stricture of the terminal ileum (arrow), positioned endoscopically.

References

1. Williams AJK, Palmer KR. Endoscopic balloon dilatation as a therapeutic option in the management of intestinal strictures resulting from Crohn's disease. *Br J Surg* 1991; 78: 453-4.
2. Vij JC, Ramesh GN, Choudhary V, Malhotra V. Endoscopic balloon dilatation of tuberculous duodenal strictures. *Gastrointest Endosc* 1992; 38: 510-1.
3. Elhence IP. Abdominal tuberculosis as observed by a surgeon. *Indian J Tuberc* 1970; 26: 59-61.

Corresponding Author

D. K. Bhasin, M.D.
H. No. 1037
Sector 24 B
Chandigarh 160023, India
Fax: +91-172 540401