Ileal Microadenomas in Familial Adenomatous Polyposis: Observation Under Magnifying Colonoscopy

Recent advances in endoscopic instrumentation provide the opportunity to observe the gastrointestinal mucosa at high magnification. It has been recently reported that characterization of surface features under magnifying colonoscopy has diagnostic value for colorectal neoplasia (1).

We performed magnifying ileoscopy in six patients with familial adenomatous polyposis (FAP) who were under surveillance after total colectomy with ileorectal anastomosis. After spraying 0.2% indigo carmine solution, the ileum was observed with a magnifying colonoscope (Olympus CF-200 Z). In three of the patients, this procedure identified small areas with the surface features which were distinctive from the adjacent ileal mucosa with round and small-sized villous structure (Figure 1). The areas were slightly elevated with tubular pattern. Histologic examination of the biopsy specimens obtained from these lesions revealed adenomatous tubules with mild epithelial atypia (Figure 2).

In both experimental animals treated with colon carcinogens (2) and colectomy specimens of patients with FAP (3), stereomicroscopic observation with methylene-blue staining revealed distinct foci of large crypts, termed as aberrant crypt foci (ACF). Because the ACFs contained dysplastic tubules, they are regarded as an extremely early stage of adenoma. The magnified view and histologic findings of the ileal lesions demonstrated herein were similar to ACF found in the resected colon of FAP (3).

Our experience suggests that magnifying colonoscopy, as well as stereomicroscopy (4), can detect early microadenomas in

Figure 1: Magnified view of the ileal mucosa identifies a slightly elevated lesion with the surface of tubular structure. The lesion is distinctive from the surrounding ileal villi.

patients with FAP. Although the in-vivo observation under the instrument has some limitation in interpretation of the data, this approach would be another procedure in investigating the histopathogenesis of adenomas in patients with FAP.

T. Matsumoto¹, M. Iida¹, M. Shimizu²

¹ Division of Gastroenterology, Dept. of Medicine, Kawasaki Medical School, Kurashiki City, Okayama, Japan

² Dept. of Pathology, Kawasaki Medical School, Kurashiki City, Okayama, Japan

References

1. Kudo S, Hirota S, Nakajima T, et al. Colorectal tumors and pit pattern. *J Clin Pathol* 1994; 47: 880-5.
2. Tudek B, Bird RR, Bruce WR. Foci of aberrant crypts in the colons of mice and rats exposed to carcinogens associated with foods. *Cancer Res* 1989; 49: 1236-40.
3. Roncucci L, Stamp D, Medline A, et al. Identification and quantification of aberrant crypt foci and microadenomas in the human colon. *Hum Pathol* 1992; 22: 287-94.
4. Kubota O, Kino I. Minute adenomas of the depressed type in familial adenomatous polyposis of the colon. A pathway to ordinary adenomas. *Cancer* 1993; 72: 1159-64.

Figure 2: Histologic examination of the biopsy specimen shows crowded adenomatous tubules (top) of mild dysplasia (bottom).

4. Kubota O, Kino I. Minute adenomas of the depressed type in familial adenomatous polyposis of the colon. A pathway to ordinary adenomas. *Cancer* 1993; 72: 1159-64.

Corresponding Author

T. Matsumoto, M. D.
Division of Gastroenterology
Dept. of Medicine
Kawasaki Medical School
Matsushima 577
Kurashiki City
Okayama 701-01
Japan
Fax: +81-864621199