Air Contrast Fluoroscopy as a Diagnostic Aid During Colonoscopy

It is controversial whether there is a need for a radiography facility where colonoscopy is performed. If the endoscopist relies unduly on fluoroscopic guidance, the procedure may be more difficult and prolonged (1), but selective use of it can facilitate completion of the examination, particularly when recurrent looping of the scope in a mobile left colon occurs (2,3). An additional advantage, which we have not seen described elsewhere, is the potential to obtain high-quality air contrast films, which may aid management, as a result of air insufflation during endoscopy. When it is not possible to teach the caecum, the demonstration of a lesion in this area radiologically is a compelling argument for persevering with colonoscopy or repeating it, rather than opting for a barium enema (Figure 1). Assessment of a stricture in two additional dimensions is also possible, which is particularly helpful if the en-face view at endoscopy is suboptimal (Figure 2).

Although we have not been able to assess the sensitivity of this technique, which is likely to be poor for small lesions, we would recommend its use in the above circumstances if a radiographic facility is available.

W. Dickey, K.G. Porter

Dept. of Medicine, Queen's University of Belfast, Dept. of Gastroenterology, Belfast City Hospital, Northern Ireland, United Kingdom

References

1. Waye JD. Colonoscopy without fluoroscopy. *Gastrointest Endosc* 1990; 36: 72-3.
2. Rogers BHG. Colonoscopy with fluoroscopy. *Gastrointest Endosc* 1990; 36: 71-2.
3. Saunders BP, Macrae FA, Williams CB. What makes colonoscopy difficult? *Gut* 1993; 34 (Suppl 1): S46.

Corresponding Author

W. Dickey, M. D.
Dept. of Medicine
Institute of Clinical Science
Grosvenor Road
Belfast BT126 BJ
United Kingdom

Figure 1: Because of looping, the scope could not be passed to the caecum. Fluoroscopy showed a caecal mass. Repeat colonoscopy reached the caecum, allowing visual confirmation and biopsy of a large adenocarcinoma.

Figure 2: This tight stricture of the transverse colon shows a typical malignant "apple-core" appearance on fluoroscopy.